

Table10

Pelagic fish eggs 1.00–2.00 mm diameter collected in plankton tows in the Northeast Pacific Ocean and Bering Sea.

0 mm	1.0 mm	2.0 mm	3.0 mm	4.0 mm	5.0 mm
	1----1	<i>Parophrys vetulus</i> (0.80–1.05 mm)			
	1---1	<i>Psettichthys melanostictus</i> (0.83–1.04 mm)			
	1-----1	<i>Isopsetta isolepis</i> (0.84–1.10 mm)			
	1-----1	<i>Platichthys stellatus</i> (0.88–1.30 mm)			
	1--1	<i>Bathylagus ochotensis</i> (0.92–1.10 mm)			
	1--1	<i>Gadus macrocephalus</i> (0.98–1.08 mm)			
	1--1	<i>Merluccius productus</i> (1.07–1.18 mm)			
	1----1	<i>Tetragonurus cuvieri</i> (1.10–1.30 mm)			
	1--1	<i>Sebastolobus</i> spp. (1.15–1.30 mm)			
	1-----1	<i>Pleuronichthys coenosus</i> (1.20–1.56 mm)			
	1-----1	<i>Gadus chalcogrammus</i> (1.20–1.77 mm)			
	1-----1	<i>Engraulis mordax</i> (1.23–1.55 x 0.65–0.82 mm, oval)			
	1--1	Macrouridae (1.38–1.48 mm)			
	1----1	<i>Tactostoma macropus</i> (1.38–1.55 mm)			
	1---1	<i>Nansenia candida</i> (1.39–1.56 mm)			
	1---1	<i>Bathylagus pacificus</i> (1.40–1.60 mm)			
	1----1	<i>Lyopsetta exilis</i> (1.47–1.71 mm)			
	1-----1	<i>Cololabis saira</i> (1.50–1.80 x 1.60–2.13 mm)			
	1-----1	<i>Icichthys lockingtoni</i> (1.52–1.80 mm)			
	1-----1	<i>Boreogadus saida</i> (1.53–1.90 mm)			
	1-----1	<i>Atheresthes stomias</i> (1.58–1.98 mm)			
	1----1	<i>Leuroglossus schmidti</i> (1.65–1.90 mm)			
	1-----1	<i>Pleuronectes quadrituberculatus</i> (1.67–2.21 mm)			
	1-----1	<i>Pleuronichthys decurrens</i> (1.80–2.10 mm)			
	1-----1	<i>Glyptocephalus zachirus</i> (1.80–2.20 mm)			
	1----1	<i>Anoplopoma fimbria</i> (1.85–2.10 mm)			
	1--1	Macrouridae (1.92–2.08 mm)			
		Collected off U.S. West Coast only			
		Collected in Bering Sea only			