

Table 13

Pelagic fish eggs with oil globules collected in plankton tows in the Northeast Pacific Ocean and Bering Sea.

0 mm	1.0 mm	2.0 mm	3.0 mm	4.0 mm	5.0 mm
	1--1	<i>Citharichthys stigmaeus</i> (0.62–0.66 mm; one oil globule 0.06–0.08 mm)			
	1---1	<i>Paralichthys californicus</i> (0.68–0.83 mm; one oil globule 0.12–0.16 mm)			
	1-1	<i>Citharichthys sordidus</i> (0.78–0.84 mm; one oil globule 0.08–0.11 mm)			
	1--1	<i>Bathylagus ochotensis</i> (0.92–1.10 mm; multiple oil globules coalescing to two)			
	1-1	<i>Merluccius productus</i> (1.07–1.18 mm; one oil globule 0.27–0.34 mm)			
	1----1	<i>Sebastolobus</i> spp. (1.15–1.40 mm; one oil globule 0.10–0.20 mm)			
	1----1	<i>Nansenia candida</i> (1.39–1.56 mm; one oil globule 0.41–0.49 mm)			
	1----1	<i>Tactostoma macropus</i> (1.39–1.54 mm; one oil globule 0.30–0.40 mm)			
	1----1	<i>Bathylagus pacificus</i> (1.40–1.60 mm; multiple oil globules coalescing to 10–15, 0.02–0.20 mm)			
	1-----1	<i>Icichthys lockingtoni</i> (1.52–1.80 mm; one oil globule 0.30–0.44 mm)			
	1----1	<i>Leuroglossus schmid</i> (1.65–1.90 mm; multiple oil globules coalescing to one, 0.47 mm after fusion)			
	1--1	Macrouridae (1.92–2.08 mm; one large orange oil globule 0.50–0.80 mm)			
	1-----1	<i>Microstoma</i> sp. (2.00–2.40 mm; one oil globule 0.49–0.82 mm)			
	1-----1	<i>Icosteus aenigmaticus</i> (2.69–3.28 mm; one oil globule 0.42–0.60 mm)			
	Collected off U.S. West Coast only				