

Northwest and Alaska Fisheries Center

National Marine Fisheries Service

U.S. DEPARTMENT OF COMMERCE

NWAFC PROCESSED REPORT 83-22

The Numbers and Distribution of Walleye Pollock Eggs and Larvae in the Southeastern Bering Sea

December 1983

This report does not constitute a publication and is for information only. All data herein are to be considered provisional.

NOTICE

This document is being made available in .PDF format for the convenience of users; however, the accuracy and correctness of the document can only be certified as was presented in the original hard copy format.

Inaccuracies in the OCR scanning process may influence text searches of the .PDF file. Light or faded ink in the original document may also affect the quality of the scanned document.

THE NUMBERS AND DISTRIBUTION OF WALLEYE POLLOCK EGGS AND LARVAE IN SOUTHEASTERN BERING SEA

by

Suam Kim

and

Arthur W. Kendall, Jr.

School of Fisheries University of Washington Seattle, Washington 98195

National Oceanic and Atmospheric Administration National Marine Fisheries Service Northwest and Alaska Fisheries Center 2725 Montlake Boulevard East Seattle, Washington 98112

December 1983

INTRODUCTION

Our interest and knowledge in walleye pollock, <u>Theragra chalcogramma</u>, the most abundant commercial fish in North Pacific Ocean has increased through last decade. At the same time, interest in petroleum development in the area has increased. Our present research is prepared to exam some possible effects of oil pollution on fish in southeastern Bering Sea. In walleye pollock the planktonic egg and larva is the life history stage that would most likely be severly impacted by oil pollution. Therefore we have examined the amount of spawning and its distribution in time and space in southeastern Bering Sea.

The main spawning ground of walleye pollock is located between Unimak Pass and the Probilof Islands. Spawning occurs during a fairly long season, primarily in spring.

METHOD AND DATA

The spatial and temporal distribution of spawning is inferred from the distribution of the planktonic eggs. For the spatial scales used in our analysis, and considering the generally low velocity of currents in the southeastern Bering Sea, differences in the spatial distribution of eggs and of spawning cannot be detected. The eggs require between 2-3 weeks to hatch at the temperatures at which they occur. Thus, the temporal distribution of spawning would be displaced 2-3 weeks earlier than the occurrence of eggs. Considering egg mortality the actual displacement would probably be less.

We reviewed information about walleye pollock egg and larval distribution from various ichthyoplankton surveys, and collected available data from them.

Our area of concern, the southeastern Bering Sea, has been examined by research teams from several different countries (U.S., U.S.S.R., Japan). However, because each team had different sampling strategies as well as different sampling gear, comparing their data is very difficult and dangerous. Therefore, we chose U.S. scientists' ichthyoplankton surveys (Waldron, 1978; Waldron and Vinter, 1978; and Walline, 1981) that used same gear and same sampling methods for surveys during 1977, 1978, and 1979.

From 1977 to 1979, the sea surface temperatures near Pribilof Islands were warmer than the average (Niebauer, 1980) while from 1973 to 1976 they were colder than the average. However, the average sea surface temperatures during March, which is assumed to be important time for pollock spawning, were nearly the same from 1977 to 1979. Therefore we feel it is reasonable to assume that there was no difference in pollock spawning pattern among the years 1977-79 (the temperature during April 1977 was below normal, although it was above normal in March and May).

In order to investigate the pattern of temporal and spatial spawning and egg and larval distribution, we divided our research area to 37 rectangles of 0.5 degree of latitude by 1 degree of longitude each (Fig. 1). We calculated the areas of the small rectangles based on Lafond (1957). For better temporal resolution, we divided the 1977 cruise into four surveys. We then arranged the U.S. ichthyoplankton surveys in seasonal sequence and assigned them Roman numbers as follows:

Survey I ; March 10-26, 1978; Waldron (1978)
Survey II : April 16-22, 1977; Waldron and Vinter (1978)
Survey III ; April 23-27, 1977; Waldron and Vinter (1978)
Survey IV ; May 3-9, 1977; Waldron and Vinter (1978)
Survey V ; May 10-15, 1977; Waldron and Vinter (1978)
Survey VI ; June 1-23, 1979; Walline (1981)

As metioned above, we divided our research area to 37 subareas. The total number of eggs or larvae in a subarea was calculated by multiplying their average number per square meter by the area of the subarea in square meters. Appendix III includes results of eggs and larval abundance calculations for each subarea.

To arrive at the temporal distribution of spawning, we assumed uniform spawning activity within the total sampling area. Therefore the total amount of eggs present in the entire samping area during each specific survey was calculated by multiplying the area occupied by each station by sum of numbers of eggs present per square meter through all stations.

Generally spawning activity with time approximates a normal distribution. Therefore we could make a normal distribution curve of pollock egg abundance with time from a polynomial equation (Tanaka, 1962) based on the above data sources:

The normal distribution curve is

$$F(X) = \frac{N}{S\sqrt{2\pi}}e^{-\frac{(x-m)^2}{2S^2}}$$

where N = total number of individuals, S = standard deviation of distribution m = mean of distribution, X = abscissa, Julian day, π = 3.14159

Taking natural logarithms, we get

$$\ln (F(X)) = (\ln \frac{N}{S/2\pi} - \frac{m^2}{2S^2}) + (\frac{m}{S^2}) X - \frac{1}{2S^2}X^2$$

This is the parabola, also called the second degree polynomial which has form of $Y=A + BX + CX^2$. Because we can get the values of the constants (A, B, and C) by use of a computer, N, m, and S can be calculated;

$$A = \ln \left(\frac{N}{S\sqrt{2\pi}}\right) - \left(\frac{m^2}{2S^2}\right)$$
$$B = \frac{m}{S^2}$$
$$C = -\frac{1}{2S^2}$$

For comparison with our study based on U.S. data, we made another normal distribution curve of pollock egg abundance with time from Moiseev and Bulatov's (1979) data. Because Moiseev and Bulatov (1979) did not include information about egg production during June, we used a digitizer to calculate egg production during June from Kendall (NWAFC, unpublished graph which was based on Bulatov (1979)).

Based on U.S. scientists' ichthyoplankton survey data, total egg production can be estimated as in Houde (1977), assuming there were no significant differences of spawning pattern during 1977-79, i.e., we combined data by month ignoring the year of the survey.

$$N = \Sigma((N_i \cdot D_i/a))$$

where N is the total eggs spawned during one spawning season,

N; is the number of eggs present in survey i

D; is the duration of survey i

a is egg hatching time

Fecundity and age composition data of pollock enable us to make an egg production estimation.

There are several data sources for age composition, fish fecundity-length or weight relationship, and von Bertalanffy parameters. After considering all of the data sources, we used data of Smith (1978), Shew (1978), Niggol (1982), and Niggol (1982), respectively because they appeared most reasonable.

In order to calculate the total biomass of eggs spawned, we used egg density data from Kanoh (1954) and egg diameter data from Nishiyama and Haryu (1981).

In general, fecundity of fish is expressed by

Fecundity =
$$A \cdot L^B$$
 (1)

where A and B are constants, and L is the length of fish.

Fecundity =
$$E \cdot W^F$$
 (2)

where E and F are constants, and W is the weight of fish.

And weight of fish is

$$W = C \cdot L^{D}$$
(3)

where C and D are constants.

Because the length and weight of fish is the function of time, we can express them with time from von Bertalanffy equation;

$$L_{t} = L_{\infty}(1 - e^{-K(t - t_{0})})$$
(4)

$$W(t) = W_{\infty}(1-e^{-K(t-t_0)})^D$$
 (5)

where L_t and Wt = the length and weight at age t L^{∞} and W^{∞} = the asymptotic value of length and weight K = a relative growth completion rate t_0 = a hypothetical age of zero size d = a dimensionless exponent reflecting absolute growth rate.

If we combine (1) and (4), and (2), (3) and (5) with each other, we get two fecundity equations with time as follow:

Formula I : Fecundity (t) = $A L_{\infty}(1-e^{-K(t-t_0)})^B$

Formula II: Fecundity (t) = E
$$(W_{\infty}(1-e^{-K(t-t_0)})^D)^F$$

From these we can get the total numbers of egg produced by multiplying fecundity by the total number of females spawning;

$$N = \Sigma(F_+ \cdot N_+)$$

where N is the total number of eggs sapwned

 F_t is the fecundity of age t fish N_t is the number of fish at age t.

RESULTS

As mentioned before, we divided our research area to 37 subareas and calculated the number of eggs and larvae present in each area during each survey (Table 1 and Fig. 2). It appears that spawning is active on upper slope area during March, it moves to middle and outer shelf during April, and moves to northwestward later in the spawning season. Roughly speaking, spawning is active at around the 100 m isobath in the southeastern Bering Sea although spawning activity probably depends on several abiotic factors in addition to depth. The larval distribution in Fig. 2 shows seaward movement of larvae after hatching. This pattern of larval distribution was also observed by Serobaba (1974).

After the numbers of egg present at each station during each survey were calculated (Appendix I), the total number of eggs present in our research area

during each survey was calculated (Table 2). Also egg numbers present in eastern Bering Sea from Moiseev and Bulatov (1979) and Kendall (NWAFC, pers. comm.) during 1978 spawning season were calculated (Table 3).

When we assumed that the spawning started on the middle of February, the normal distribution based on the above data indicates that 104.4th Julian day (14 April) is the peak of spawning with a standard deviation of 13.8 days $-\frac{(X-104.41)^2}{382.16}$ (Fig. 3: $F(X) = 27.08e^{-\frac{(X-104.41)^2}{382.16}}$). Also under the same assumption, the result derived from Moiseev and Bulatov (1979) and Kendall (pers. comm.) shows 107.9th Julian day (18 April) as the peak of spawning and 26 days as one $-\frac{(X-107.94)^2}{1428.82}$, although smaller egg production was indicated by the Soviets' data than by U.S. scientists'. These results are not significantly different from others which indicate that along the slope and outer shelf the peak spawning occurs in March and April (Lynde, 1983).

Larvae were found beginning in the middle of March in southeastern Bering Sea. Table 1 shows that the maximum larval abundance occurred at around second survey, which was conducted just after peak spawning, and that the number of larvae decreases with time. Walline (1983) calculated the egg hatching dates for pollock from 1979 survey data; hatching dates were distributed from 1 April to 15 July; the hatch was most pronounced during the last 2 weeks in April and the last 2 weeks in May.

Based on Table 2, (ichthyoplankton survey data), the total number of pollock eggs produced during the spawning season in southeastern Bering Sea was estimated. If development time from spawning to hatching is assumed to be

17.3 days which corresponds to duration of the pollock egg stage in 4° C seawater (Richard Bates, NWAFC pers. comm.), the total number becomes 3.6918 E+13 eggs (282.7 eggs/m²) (see Appendix II).

Also we calculated pollock egg production by knowing the fecundity-age composition relationship of the adult population, although more research is needed to estimate the parameters more precisely for pollock in the eastern Bering Sea. For calculations of number of eggs produced the following constants were used: von Bertalanffy constants of K=0.209, to = -0.315 year, $L^{\infty} = 65.01$ cm (Niggol, 1981); Shew's fecundity-length relationship data (1978); and Smith's age compositon data (1981). If we multiply fecundity by number of females at each age, the caculation resulted in 6.17 \cdot E+13 eggs in 159,100 km² which equals a density of 387.95 eggs/m². For transforming this number to total weight of egg (Kanoh, 1954), 1.021, and mean egg volume of 0.0027 cm³ which came from Nishiyama and Haryu (1981). The total weight of eggs produced in the spawning area of 159,100 km² is about 170,000 mt in one spawning season.

DISCUSSION

The egg density derived from fecundity data $(387.95 \text{ eggs/m}^2)$ is higher than that of the ichthyoplankton surveys (282.7 eggs/m^2) and that of Serobaba (1968) $(293.44 \text{ eggs/m}^2)$. This indicates that spawning probably occurs outside the areas sampled during these ichthyoplankton surveys. Also catchability of eggs may be less than 100% with the methods used in these surveys.

In performing this study, the biggest problem is in data variation. Because small numbers of ichthyoplankton surveys have been conducted in Bering Sea and data collecting methods have varied, we needed to use many assumptions, which should be reconsidered when more data becomes available. Also the basic biology of pollock is not well understood. We do not know the exact spawning time of pollock in southeastern Bering Sea nor do we understand the variation in size of pollock eggs, which influences our estimation of pollock egg biomass. In order to understand pollock in Bering Sea more exactly, we need to increase research activities to study their biology in relation to oceanographic conditions.

CONCLUSIONS

Spawning occurs over a large area of the southeastern Bering Sea, mainly between the Pribilof Islands and Unimak Pass, between the 100-200 m isobaths. Present data are insufficient to resolve the pattern of spawning and larval distribution adequately. The total spawning area was probably not sampled during the surveys considered in this report.

Spawning peaks on about Julian day 104-108 (14-18 April) with a standard deviation of between 14 and 26 days, based on two sets of ichthyoplankton surveys. Some spawning occurs from about 15 February through June.

Based on fecundity and size of the adult population, 6.17 • E+13 eggs are produced during the spawning season. Based on ichthyoplankton surveys 3.69 • E+13 eggs are produced during the spawning season. The difference between these two estimates may partially be a result of the plankton surveys not covering the entire spawning area. The density of egg spawned based on adult

population parameters was 388 eggs/m^2 while that based on plankton surveys was 283 eggs/m^2 . This difference may be due in part to problems in plankton sampling and egg mortality.

REFERENCES

- Bulatov, O. A. 1979. Distribution of eggs and larvae of walleye pollock and other fish species in the eastern part of the Bering Sea, 1978-1979. Pacific Research Institute Fisheries and Oceanography (TINRO).
- Houde, E. D. 1977. Abundance and potential yield of the round herring, <u>Etrumeus teres</u>, and aspects of its early life history in the eastern Gulf of Mexico. Fish. Bull. 75(1):61-89.
- Kanoh, Y. 1954. On the buoyancy of the egg of Alaska pollock, <u>Theragra</u> chalcogramma. Japan J. Icchthyol. 3:238-246.
- Lafond, E. C. 1957. Processing oceanographic data. H. O. Pub. No. 614, U.S. Hydrographic Office, Washington, D.C.
- Lynde, C. M. 1983. Juvenile and adult walleye pollock of the Eastern Bering Sea. Proc. of 1983 Eastern Bering Sea pollock ecosystem workshop. NWAFC.
- Moiseev, E. I., and O. A. Bulatov. 1979. The state of pollock stocks in Eastern Bering Sea. Pacific Research Inst. of Fisheries and Oceanography (TINRO).

- Niebauer, H. J. 1980. Sea ice and temperature variability in the eastern Bering Sea and the relation to atmospheric fluctuations. J. Geophys. Res. 85:7507-7515.
- Niggol, K. 1982. Data on fish species from Bering Sea and Gulf of Alaska. NOAA Tech. Memo. NMFS F/NWC-29.
- Nishiyama, T. and T. Haryu. 1981. Distribution of walleye pollock eggs in the upper most layer of the Southeastern Bering Sea. In: Hood, D. W. and J. A. Calder (eds.), The eastern Bering Sea shelf: Oceanography and resources Vol. 2 (Chap. 59), pp. 993-1012, U.S. Dept. of Comm. NOAA.
- Serobaba, I. I. 1968. Spawning of Alaskan pollock, <u>Theragra chalcogramma</u> (Pallas), in the Northeastern Bering Sea. Vopr. Ichthyol. 8(6), 789-798.
- Serobaba, I. I. 1974. Spawning ecology of the walleye pollock, <u>Theragra</u> chalcogramma, in the Bering Sea. J. Ichthyol. 14:544-552.
- Shew, P. M. 1978. Pollock fecundity study; Preliminary report. NMFS, NWAFC, unpub. MS.
- Smith, G. B. 1981. The biology of walleye pollock. In: Hood, D. W. and J. A. Calder (Eds.), The eastern Bering Sea Shelf: Oceanography and Resources. VI. (Ch. 33) pp 527-551. U.S. Dept. of Comm., NOAA.

- Tanaka, S. 1962. A method of analyzing a polymodal frequency distribution and its application to the length distribution of porgy, <u>Taius</u> tumifrons. J. Fish. Res. Bd. Canada, 19(6).
- Waldron, K. D. 1978. Ichthyoplankton of the eastern Bering Sea. NMFS, NWAFC, Proc. Rep.
- Waldron, K. D., and B. M. Vinter. 1978. Ichthyoplankton of the eastern Bering Sea. NMFS, NWAFC, Proc. Rep.
- Walline, P. D. 1981. Distribution of ichthyoplankton in the Eastern Bering Sea during June and July, 1979. NMFS, NWAFC, Proc. Rep. 81-03.
- Walline, P. D. 1983. Growth of larval and juvenile walleye pollock related to year-class strength. Ph.D. Thesis, Univ. of Washington.

	Survey						
Subarea	I	II	III	IV	۷	VI	
A(1,1)		0.0000			1.3560 (0.428)		
A(1,2)		0.0381 (0.114)			1.8716 (0.496)	0.0000 (0.000)	
A(1,3)						0.0000 (0.024)	
A(1,4)						0.0221 (0.092)	
A(1,5)						0.1158 (0.719)	
A(2,1)						0.0000 (0.000)	
A(2,2)			1.0005 (1.221)		0.0867 (0.737)	0.0000 (0.024)	
A(2,3)			1.2447 (0.181)		0.8949 (0.291)	0.0000 (0.000)	
A(2,4)			0.5220 (0.019)		0.6674 (0.000)	0.0079 (0.075)	
A(2,5)			12.9470 (0.000)	3.7147 (0.000)			
A(3,1)					0.0485 (0.388)		
A(3,2)	0.0193	0.0397 (0.218)			0.0413 (0.496)	0.0000 (0.000)	

Table 1.--The total numbers of eggs (E+11) and larvae (E+11) present in each sub area during each survey. Larval data are in parentheses.

	Survey						
Subarea	I	II	III	IV	٧	VI	
A(3,3)	0.0182	0.1532	0.1871		0.4421	0.0000	
		(0.153)	(0.122)	(1.118)		(0.103)	
A(3,4)	0.0000	0.1547	0.1793		0.2623	0.0000	
		(0.000)	(0.021)		(0.031)	(0.123)	
A(3 5)	0 0000	0 6378	0 3298	0 7144			
Λ(3,3)	0.0000	(0,000)	(0,000)	(0.675)			
		(0.000)	(0.000)	(0.075)			
A(3,6)		25.8220		0.7017			
		(0.000)		(0.000)			
10 m 10 10							
A(4,1)	0.1158						
A(A 2)	2 2111	0 0967			0 0212	0 0000	
A(4,2)	2.2444	(12 113)			(0.255)	(0.000)	
		(12.115)			(0.255)	(0.000)	
A(4,3)	0.0550	0.1239	0.3672	0.1437	0.2008	0.0000	
		(0.137)	(0.516)	(1.265)	(3.414)	(0.015)	
A(4,4)	0.0000	0.0186	0.0102	0.1648			
		(0.008)	(0.051)	(0.991)			
	0 0200	1 2577	2 0710	0.0004			
A(4,5)	0.0309	1.35//	3.9/12	2.2024			
		(0.0/1)	(0.135)	(0.822)			
A(4,6)	0.0000	2.5091	11,9090	7,6239			
		(0.000)	(0.000)	(0.590)			
		, , , , , , , , , , , , , , , , , , , ,	(,			
A(4,7)				0.4143			
				(0.000)			

A(5,1) 0.0000

(0.025)

	Survey							
Subarea	I	II	III	IV	۷	νı		
A(5,2)	0,2358	0.5192		0.0677		0.000		
		(16.817)		(0.293)		(0.000		
A(5,3)	34.3490	0.0160	0.0483	0.0990		0.000		
		(1.495)	(6.884)	(2.299)		(0.012		
A(5,4)		0.1257	0.5852	0.4044		0.000		
		(3.839)	(3.162)	(0.929)				
A(5,5)	0.0184	0.6430	0.6932	0.6790				
		(0.047)	(0.158)	(0.980)				
A(5,6)		0.0559	0.0000	0.1966				
		(0.000)	(0.000)					
A(6,1)						0.000		
A(6,2)		0.1382	0.1100	0.2351	0.0882			
		(0.737)	(2.134)	(2.962)	(0.639)			
A(6,3)	1.0763	0.0223	0.0206		0.0758			
		(2.053)	(5.615)		(0.162)			
A(6,4)	0.0000	0.4393	5.3085	0.7404	0.0468			
		(4.675)	(2.628)	(0.912)	(0.937)			
A(6,5)	0.0000	2.2694	13.9810	2.5505				
		(5.266)	(8.063)	(0.548)				
A(7,1)				0.2068				
				(3.392)				
A(7,2)			0.2745	0.1067				
			(5.108)	(1.328)				

	Survey						
Subarea	I	II	III	IV	۷	VI	
A(7,3)		0.4877			0.0843		
		(20.950)			(0.139)		
Total	38.163	35.620	53.689	21.027	6.188	0.146	
(E+11)		(68.686)	(36.018)	(19.104)	(8.413)	(1.212)	
Mean	2.245	1.781	2.557	1.168	0.413	0.009	
(E+11)		(3.122)	(1.801)	(1.061)	(0.601)	(0.076)	
No. animals/	87.55	69.46	99.71	45.56	16.09	0.34	
survey (E+11)		(109.27)	(63.03)	(37.14)	(21.04)	(2.65)	

Survey duration	Mid-day of survey (Julian day (X))	No. of station sampled	Total eggs during survey (E+11 eggs)	Number of survey days	No. of eggs sampled/day (E+11)
Feb. 14	45			1	0.001
Mar. 10-16	71	21	70.100	7	10.010
Apr. 16-22	108	35	119.832	7	17.119
Apr. 23-27	114	31	90.430	5	18.086
May 3-9	125	28	40.505	7	5.786
May 10-15	131.5	17	14.564	6	2.427
June 1-23	162	39	0.170	22	0.008

Table 2. Summary table of egg for U.S. scientists' data.

Survey duration	Mid-day of survey (Julian day (X))	Total eggs during survey (E+8)	No. of eggs sampled per day (E+8)
April 10 - May 3	110.5	294.7	12.28
May 10-20	134.0	151.0	13.73
June 1-30	166.0	40.2	1.34

.

Table 3. The number of eggs present in spawning areas from Soviet survey.

FIGURES

- Figure 1. Research area in southeastern Bering Sea. The size of each subarea is 0.5° latitude by 1° longitude. 100 m isobath is included.
- Figure 2. Spatial distribution of pollock eggs and larvae. The three subareas with the greatest densities during each survey are indicated. 100 m isobath is included. (A) Eggs (B) Larvae.
- Figure 3. The relative abundance of pollock egg with time. (A) Based on U.S. scientists' data. (B) Based on Moiseev and Bulatov (1978) and Kendall (pers. comm.).



Figure 1. Research area in southeastern Bering Sea. The size of each subarea is 0.5° latitude by 1° longitude. 100 m isobath is included.



Figure 2. Spatial distribution of pollock eggs and larvae. The three subareas with the greatest densities during each survey are indicated. 100 m isobath is included. (A) Eggs,



Figure 2. Spatial distribution of pollock eggs and larvae. The three subareas with the greatest densities during each survey are indicated. 100 m isobath is included. (B) Larvae.



Figure 3. The relative abundance of pollock egg with time. (A) Based on U.S. scientists' data. (B) Based on Moiseev and Bulatov (1978) and Kendall (pers. comm.).

Appendix I: The number of eggs present at each station during each survey.

Station	No. of eggs sampled during survey	SHF	No.	of eggs/m ²
16	203	4.252		86.3156
17	5	5.379		2.6895
18	11	5.980		6.5780
19	1858	5.158		958.3564
20	100	6.340		63.4000
21	6	5.450		3.2700
22	1	5.494		0.5494
23	4	4.876		1.9504
24	1	5.181		0.5181
28	2	5.791		1.1582
29	3	5.817		1.7451
32	1	5.147		0.5147
			Total	1127.0454

 The total number of eggs present at each station during March 10-16, 1978. (Survey I)

SHF (Standard Haul Factor) converts actual catch to numbers beneath 10 \mbox{m}^2 of sea surface.

Total number of egg present in area of concern = (130613.6*1000000/21)*1127.045 = 7.0099*E+12

		Surv	ev II		Surve	y 111		Surve	y 1V		Surve	y V
Station	2005	No. 0	if equa	0.115	No. o	f equs	CUE	No. of	eggs 2	eur.	No. of	eggs 2
	SHF	total	/m*	SHE	total	/m²	SHF	total	/m~	SHF	total	/៣-
1	6.55	30	19.635	6.69	427	285.834	6.62	58	38.367			
2	5.73	8	4.584	6.19	11	6.810	6.12	4	2.448	6.46	2	1.219
3	4.99	1	0.499	6.53	1	0.653	5.42	6	3.253			
4	3.92	1	0.392	6.80	3	2.041	5.68	4	2.272			
5	3.04	11	3.347	6.77	6	4.059				5.67	10	5.672
6	4.06	9	3.653	5.96	28	16.685	4.06	10	4.059			
7	5.53	2	1.106	6.54	24	15.698	4.86	19	9.234			
8	5.91	10	5.910	6.28	27	16.956	5.80	23	13.333	Bongo	tow aborte	ed
9	4.73	131	62.553	5.68	678	385.375	5.81	121	70.301	2		
10	4.77	12	5.725	5.49	9	4.945	4.78	55	26.301			
11	5.20	58	30.154	5.84	58	33.855	5.04	23	11.587			
12	4.34	-	-	6.38	-	_	4.81	15	7.221			
13	5.25	2	1.050	5.78	1	0.578	4.18	5	2.091			
14	5.50	3	1.650	5.96	4	2.385				5.75	16	9.197
15	5.04	14	7.062	5.50	15	8.252				5.50	29	15.936
16	6.00	7	4,202	5.59	14	7.829				5.99	22	13.174
17	5.10	9	4.594	5.91	4	2.363	5.79	3	1.736			
18	5.00	6	2.998	5.87	-	-	5.15	4	2.059			
19	5.38	137	73.706	6.03	372	224.353	5.54	227	125.758			
20	5.20	3	1.561	5.73	-	-	5.49	10	5.485			
21	4.34	2	0.868	4.54	121	54.922	4.52	-				
22	5.73	246	140.884	5,50	1123	617.875	4.90	879	430.710			
23	5.82	36	20.941	5.58	25	13.958	5.18	68	35.251			
24	6.38	24	15.314	5.99	8	4.790	5.36	10	5.360			
25	5.98	1741	1041.466				5.08	16	8.120			
26	6.96	682	474.536				5.12	62	31.769			
27	Bongo	tow abor	rted				4.07	54	21.967			
39	5.60	2	1.120							4.70	117	54.943
40	4.86	-	-							5.24	76	39.809
49	6.35	6	3.808				6.48	10	6.479	6.08	4	2.430
50	6.99	19	13.283							7.65	3	2.296
51	6.15	1	0.615							6.85	3	2.055
52	6.30	23	14.485				6.29	د	1.888			
53	6.12	4	2.448							6.01	1	0.601
54	5.64	2	1.128							5.87	2	1.174
55	Bongo	tow abor	ted							6.89	2	1.379
58				5.87	50	29.370				6.37	4	2.546
59				5.89	62	36.537				6.11	43	26.269
60				5.68	27	15.323				6.32	31	19.592
61				5.24	726	380.061	5.65	193	109.045			
62							4.19	142	59.427			
64				6.48	1	0.648				7.07	3	2.122
65				6.06	5	3.032	5.81	5	2.906			
66				6.23	12	7.476	5.63	10	5.633			
	Tot	al =	1965.277			2182,660			984.634			200.486

2) The number of eggs present at each station during April 16-May 15, 1977.*

Total number of equs present in area of concern: subarea 1; Station No. 1-24 subarea 2: Station No. 25-66 Survey II subarea 1: (4.404 E+10/24) (19.6350+4.5840+,,,+15.3144) = 7.5673°E+11 +) subarea 2: (5.166 E+10/10) (1041.4662+474.5356+,,,+1.1282) = 80.222 · E+11 87.789 E+11 Survey III subarea 1: (4.404 E+10/24) · (285.8338+6.8101+,,,+4.7896) = 31.382*E+11 subarea 2: (5.166 E+10/8) (29.370+36.537+,,,+7.476) = 34.866°E+11 +) 66.248°E+11 Survey IV subarea 1: (4.404 E+10/20) (38.367+2.448+,,,+5.360) = 17.546°E+11 +) subarea 2: (5.166 E+10/8) (8.120+31.769+,,,+5.633) = 12.128°E+11 29.674 E+11 Survey V = 3.9873°E+11 subarea 1: (4.404 E+10/5) (1.2912+5.672+,,+13.174) +) subarea 2: (5.166 E+10/12) (54.943+39.809+,,,+2.122) $= 6.6821 \cdot E + 11$ 10.6694 E+11

*They mentioned that the total area of subarea 1 and 2 is 95700 km^2 . Because we assumed equal production rate outside this area, we can corrected that number to 130613.6 km² by multiplying 1.365 to total egg number present in subarea 1 and 2;

	Survey							
	II	III	IV	V				
Amount of egg present (E+11)	119.832	90.430	40.505	14.564				

3) Total number of eggs present at each station during June 1-23, 1979 (Survey VI).

Station	No. of eggs/m ²
V01-8 S 46A	0.7
5 1ZA	Total = $\frac{3.4}{5.4}$

Total number of egg present in area of concern = $(130613.6^{\circ}1000000/40)^{\circ}5.4$ = $1.763^{\circ}E+10$

Survey	(F+11)	Di
····	eggs	days
I	70.10	34.5
II	119.83	22.5
III	90.43	7.5
IV	40.51	9.5
V	14.56	14.0
VI	0.17	31.0

Appendix II: The estimation of total egg number produced.

N = Σ((N_i •D_i/a) = (6386.805/17.3)E+11 = 369.18•E+11 (eggs)

Appendix III: Total number of pollock eggs and larvae in subarea A(i,j) during specific survey.

Subarea	No. of times sampled	No. of times egg caught	Survey (station number)	No. of eggs/ m ²	Total No. of eggs present /subarea
A(1,1)	2	1	II (40) V (40)	0 39.8088	0 1.3560 E+11
A(1,2)	5	3	II (39) V (39) VI (S47A, S48	1.1196 54.9432 BA) O	3.8140 E+9 1.8716 E+11 0
A(1,3)	1	1	VI (S45A)	0	0
A(1,4)	2	1	VI (S40A, S46	GA) 1.3000	2.2142 E+9
A(1,5)	1	1	VI (S12A)	3.4000	1.1582 E+10
A(2,1)	2	2	VI (S50A, S53	BA) O	0
A(2,2)	4	2	III (58) V (58) VI (S41A, S44	29.3700 2.5460 A) 0	1.0005 E+11 8.6732 E+9 0
A(2,3)	4	2	III (59) V (59) VI (S38A, S39	36.5366 26.2687 9A) 0	1.2447 E+11 8.9487 E+10 0
A(2,4)	5	3	III (60) V (60) VII (V01-8	15.3225 19.5920	5.2198 E+10 6.6742 E+10
			\$33A, \$34A)	0.2333	0.7949 E+9
A(2,5)	2	2	III (61) IV (61)	380.0610 109.0450	1.2947 E+12 3.7147 E+11
A(3,1)	2	1	II (55) V (55)	Bongo tow a 1.3788	borted 4.8509 E+9
A(3,2)	5	3	I (22) II (54) V (54) VI (S42A, S43	0.5494 1.1282 1.1740 BA) 0	1.9329 E+9 3.9692 E+9 4.1304 E+9 0

EGGS

Subarea	No. of times sampled	No. of times egg caught	Survey (station number)	No. of eggs/ m ²	Total No. of eggs present /subarea
A(3,3)	11	7	I (24) II (14, 15) III (14, 15) IV (14, 15) VI (SO8A, SO8B S29A, S35A)	0.5181 4.3557 5.3184 12.5662 0	1.8228 E+9 1.5324 E+10 1.8711 E+10 4.4210 E+10 0
A(3,4)	11	6	I (25) II (16,17) III (16, 17) V (16, 17) VI (SO9A, S3OA, S31A, S32A)	0 4.3979 5.0958 7.4546 0	0 1.5473 E+10 1.7928 E+10 2.6227 E+10 0
A(3,5)	7	6	I (26) II (23, 24) III (23, 24) IV (23, 24)	0 18.1278 9.3736 20.3056	0 6.3777 E+1(3.2978 E+1) 7.1439 E+1)
A(3,6)	4	4	II (25, 26) 7 IV (25, 26)	58.0009 19.9444	2.5822 E+1 7.0168 E+1
A(4,1)	1	1	I (21)	3.2700	1.1576 E+1
A(4,2)	4	3	I (20) II (53) V (53) VI (S13A)	63.4000 2.4484 0.6014 0	2.2444 E+1 8.6676 E+9 2.1290 E+9 0
A(4,3)	13	8	I (23, 28) II (5, 6) III (5, 6) IV (6) V (5) VI (S06A, S07A, S28A, S36A, S37A	1.5543 3.5002 10.3721 4.0590 5.6720	5.5024 E+9 1.2391 E+1 3.6719 E+1 1.4369 E+1 2.0080 E+1 0
A(4,4)	7	4	I (27) II (12, 13) III (12, 13) IV (12, 13)	0 0.5251 0.2890 4.6560	0 1.8589 E+9 1.0231 E+9 1.6483 E+1
A(4,5)	8	6	I (29, 30) II (18, 19)	0.8726 38.3521	3.0890 E+9 1.3577 E+1

	and sense the data set of the second second			and the second se	
Subarea	No. of times sampled	No. of times egg caught	Survey (station number)	No. of eggs/ m ²	Total No. of eggs present /subarea
			III (18, 19) IV (18, 19)	112.1766 63.9084	3.9712 E+11 2.2624 E+11
A(4,6)	7	5	I (31) II (21, 22) III (21, 22) IV (21, 22)	0 70.8759 336.3983 215.3550	0 2.5091 E+11 1.1909 E+12 7.6239 E+11
A(4,7)	3	2	II (27) IV (27)	Bongo tow a 21.9672	borted 7.7767 E+10
A(5,1)	1	0	VI (S14A)	0	0
A(5,2)	6	3	I (18) II (52) IV (52) VI (504A, 505	6.5780 14.4854 1.8882 A.	2.3577 E+10 5.1918 E+10 6.7676 E+9
			S27A)	n n	0
A(5,3)	10	7	I (19) II (3, 4) III (3, 4) IV (3, 4) VI (SO1A, SO2 SO3A)	958.3564 0.4456 1.3471 2.7623 A 0	3.4349 E+12 1.5971 E+9 4.8282 E+9 9.9005 E+9 0
A(5,4)	8	6	II (7, 8) III (7, 8) IV (7, 8) V (8) VI (S14A)	3.5078 16.3272 11.2836 Bongo tow 0	1.2572 E+10 5.8519 E+10 4.0442 E+10 aborted 0
A(5,5)	7	7	I (32) II (10, 11) III (10, 11) IV(10, 11)	0.5147 17.9397 19.3996 18.9442	1.8448 E+9 6.4298 E+10 6.9317 E+10 6.7899 E+10
A(5,6)	3	2	II (20) III (20) IV (20)	1.5609 0 55.4850	5.5945 E+9 0 1.9659 E+10
A(6,1)	1	0	VI (S26A)	0	0

Subarea	No. of times sampled	No. of times egg caught	Survey (station number)	No. of eggs/ m ²	Total No. of eggs present /subarea
A(6,2)	4	4	II (49) III (65) IV (49) V (49)	3.8082 3.0320 6.4790 2.4304	1.3816 E+10 1.1000 E+10 2.3506 E+10 8.8174 E+9
A(6,3)	7	6	I (15, 16, 1 II (51) III (64) V (51, 64)	7) 29.6684 0.6151 0.6476 2.0885	1.0763 E+11 2.2316 E+9 2.2061 E+9 7.5770 E+9
À(6,4)	8	7	I (12) II (1, 2) III (1, 2) IV (1, 2) V (2)	0 12.1095 146.3220 20.4077 1.2912	0 4.3933 E+10 5.3085 E+11 7.4038 E+10 4.6844 E+9
A(6,5)	4	3	I (13) II (9) III (9) IV (9)	0 62.5525 385.3752 70.3010	0 2.2694 E+11 1.3981 E+12 2.5505 E+11
A(7,1)	1	1	IV (66)	5.6330	2.0681 E+10
A(7,2)	2	2	III (66) IV (65)	7.4760 2.9060	2.7447 E+10 1.0669 E+10
A(7,3)	2	2	II (50) V (50)	13.2829 2.2962	4.8767 E+10 8.4303 E+9
LARVAE					
	No. of	No. of	Survey	No. of	Total No. of

Subarea	No. of times sampled	No. of times larvae caught	Survey (station number)	No. of laryae/ m ²	Total No. of larvae present (E+10)/subarea
A(1,1)	2	1	II (40) V (40)	0 12.5712	0 4.2822
A(1,2)	4	2	II (39) V(39) VI (S47A,S48A)	3.3588 14.5576 0	1.1441 4.9588 0

Subarea	No. of times sampled	No. of times larvae caught	Survey (station number)	No. of laryae/ m ²	Total No. of larvae present (E+10)/subarea
A(1,3)	1	1	VI (S45A)	0.7	0.2384
A(1,4)	1	1	VI (S40A)	2.7	0.9197
A(1,5)	1	1	VI (S12A)	21.1	7.1874
A(2,1)	2	0	VI (S50A,S53A)	0	0
A(2,2)	4	4	III (58) V (58) VI (S41A,S44A)	35.8314 21.6410 0.7	12.2060 7.3722 0.2385
A(2,3)	4	2	III (59) V (59) VI (S38A,S39A)	5.3037 8.5526 0	1.8068 2.9135 0
A(2,4)	5	3	III (60) V (60)	0.5675 0	0.1933 0
			S33A,S34A)	2.2	0.7495
A(2,5)	2	0	III (61) V (61)	0 0	0 0
A(3,1)	1	1	V (55)	11.0304	3.8807
A(3,2)	4	2	II (54) V (54) VI (5420	6.2051 14.0880	2.1831 4.9564
			S43A)	0	0
A(3,3)	9	7	II (14, 15) III (14, 15) IV (14, 15)	4.3557 3.4623 31.7734	1.5324 1.2181 11.1780
			S29A, S35A)	2.93	1.0308
A(3,4)	9	5	II (16, 17) III (16, 17) V (16, 17)	0 0.5907 0.8779	0 0.2079 0.3089
			S31A, S32A)	3.5	1.2314
A(3,5)	6	1	II (23, 24) III (23, 24) IV (23, 24)	0 0 19.1808	0 0 6.7482

Subarea	No. of times sampled	No. of times larvae caught	Survey (station number)	No. of larvae/ m ²	Total No. of larvae present (E+10)/subarea
A(3,6)	4	0	II (25, 26) IV (25, 26)	0 0	0 0
A(4,2)	3	2	II (53) V (53) VI (S13A)	342.1639 7.2168 0	121.13 2.5548 0
A(4,3)	11	7	II (5, 6) III (5, 6) IV (6) V (5) VI (S06A, S07A, S28A, S36A, S37A)	3.8558 14.5679 35.7192 96.4240 0.42	1.3650 5.1572 12.6450 34.1350 0.1487
A(4,4)	6	4	II (12,13) III (12, 13) IV (12, 13)	0.2171 1.4448 28.0048	0.0769 0.5115 9.9141
A(4,5)	6	4	II (18, 19) III (18, 19) IV (18, 19)	1.9988 3.8155 23.2091	0.7076 1.3507 8.2163
A(4,6)	6	1	II (21, 22) III (21, 22) IV (21, 22)	0 0 16.66	0 0 5.8979
A(4,7)	1	0	IV (27)	0	0
A(5,1)	1	1	VI (S14A)	0.7	0.2509
A(5,2)	5	2	II (52) IV (52) VI (SO4A, SO5A, S27A)	469.2010 8.1822 0	168.1700 2.9326 0
A(5,3)	9	7	II (3, 4) III (3, 4) IV (3, 4) VI (SO1A, SO2A, SO3A)	41.7090 192.0720 64.1443 0.3333	14.9490 68.8410 22.9900 0.1195
A(5,4)	6	5	II (7, 8) III (7, 8) IV (7, 8)	107.1016 88.2340 25.9135	38.3870 31.6240 9.2878

Subarea	No. of times sampled	No. of times larvae caught	Survey (station number)	No. of laryae/ m ²	Total No. of larvae present (E+10)/subarea
A(5,5)	6	5	II (10, 11) III (10, 11) IV (10, 11)	1.2998 4.4124 27.3296	0.4659 1.5815 9.7953
A(5,6)	3	1	II (20) III (20) IV (20)	0 0 2.7425	0 0 0.9830
A(6,2)	4	4	II (49) III (65) IV (49) V (49)	20.3104 58.8208 81.6354 17.6204	7.3685 21.3400 29.6170 6.3926
A(6,3)	4	4	II (51) III (64) V (51, 64)	56.5984 154.7764 4.4744	20.5340 56.1520 1.6233
A(6,4)	7	7	II (1, 2) III (1,2) IV (1, 2) V (2)	128.8466 72.4442 25.1509 25.8240	46.7450 26.2820 9.1246 9.3688
A(6,5)	3	3	II (9) III (9) IV (9)	145.1600 222.2444 15.1060	52.6630 80.6290 5.4804
A(7,1)	1	1	IV (66)	93.5078	33.9240
A(7,2)	2	2	III (66) IV (65)	140.7980 36.6156	51.0810 13.2840
A(7,3)	2	2	II (50) V (50)	577.4566 3.8270	209.5000 1.3884

