

## 1.B. Assessment of walleye pollock in the Bogoslof Island Region

James N. Ianelli, S. J. Barbeaux, D. McKelvey and T. Honkalehto

Alaska Fisheries Science Center  
National Marine Fisheries Service

### Executive Summary

Management recommendations from last year remain the same since no new information is available on this stock in 2013. The most recent acoustic trawl survey estimate, from March 2012, was 67,063 t. The resulting 2014 ABC and OFL values (under Tier 5) are thus summarized below:

Quantity	As estimated or <i>specified last year for:</i>		As estimated or <i>recommended this year for:</i>	
	2013	2014	2014	2015
$M$ (natural mortality rate)	0.2	0.2	0.2	0.2
Tier	5	5	5	5
Biomass (t)	67,063	67,063	67,063	67,063
$F_{OFL}$	0.200	0.200	0.200	0.200
$maxF_{ABC}$	0.150	0.150	0.150	0.150
$F_{ABC}$	0.150	0.150	0.150	0.150
OFL (t)	13,413	13,413	13,413	13,413
maxABC (t)	10,059	10,059	10,059	10,059
ABC (t)	10,059	10,059	10,059	10,059
Status	As determined <i>this year for:</i>		As determined <i>this year for:</i>	
	2011	2012	2012	2013
Overfishing	No	n/a	No	n/a

### Introduction

Alaska pollock (*Theragra chalcogramma*) are broadly distributed throughout the North Pacific with largest concentrations found in the Eastern Bering Sea. The Bogoslof region is noted for having distinct spawning aggregations that appear to be independent from pollock spawning in nearby regions. The Bogoslof management district (INPFC area 518) was established in 1992 in response to fisheries and surveys conducted during the late 1980s, which consistently found a discrete aggregation of spawning pollock in this area during the winter. The degree to which this aggregation represents a unique, self-recruiting stock is unknown but the persistence of this aggregation suggests some spawning site fidelity that called for independent management. The Bogoslof region pollock has also been connected with the historical abundance of pollock found in the central Bering Sea (Donut Hole) due to concentrations of pollock successively moving toward this region prior to spawning (Smith 1981).

Collectively, pollock found in the Donut Hole and in the Bogoslof region are considered a single stock, the Aleutian Basin stock. Currently, based on an agreement from a Central Bering Sea convention meeting, it is assumed that 60% of the Aleutian Basin pollock population spawns in the Bogoslof region. The actual distribution of Aleutian Basin pollock is unknown and likely varies depending on environmental conditions and the age-structure of the stock. The Bogoslof component of the Aleutian Basin stock is one of three management stocks of pollock recognized in the BSAI region. The other stocks include pollock found in the large area of the Eastern Bering Sea shelf region and those in the

Aleutian Islands near-shore region (i.e., less than 1000m depth; Barbeaux et al. 2004). The Aleutian Islands, Eastern Bering Sea and Aleutian Basin stocks probably intermingle, but the exchange rate and magnitude are unknown. The degree to which the Bogoslof spawning component contributes to subsequent recruitment to the Aleutian Basin stock also is unknown. From an early life-history perspective, the opportunities for survival of eggs and larvae from the Bogoslof region seem smaller than for other areas (e.g., north of Unimak Island on the shelf). There is a high degree of synchronicity among strong year-classes from these three areas, which suggests either that the spawning source contributing to recruitment is shared or that conditions favorable for survival are shared. From a biological perspective, the degree to which these management units are reasonable definitions depends on the active exchange among these stocks. If they are biologically distinct and have different levels of productivity, then management should be adjusted accordingly. Bailey et al. (1999) present a thorough review of population structure of pollock throughout the north Pacific region. They note that adjacent stocks were not genetically distinct but that differentiation between samples collected on either side of the N. Pacific was evident.

There are some characteristics that distinguish Bogoslof region pollock from other areas. Growth rates appear different (based on mean-lengths at age) and pollock sampled in the Bogoslof Island survey tend to be much older. For example, the average percentage (by numbers of fish older than age 6) of age 15 and older pollock observed from the Bogoslof AT survey since 1988 is 18%; in the EBS region (from model estimates), the average from this period is only 2%. The information available for pollock in the Aleutian Basin and the Bogoslof Island area indicates that these fish may belong to the same “stock”. The pollock found in winter surveys are generally older than age 4 and are considered distinct from eastern Bering Sea pollock. Although data on the age structure of Bogoslof pollock show that a majority of pollock originated from year classes that were also strong on the shelf, 1972, 1978, 1982, 1984, 1989, 1992, 1996, 2000, and 2006 there has been some indication that there are strong year classes appearing on the shelf that have not been as strong (in a relative sense) in the Bogoslof region (Ianelli et al., 2004). Strong year classes of pollock in Bogoslof may be functionally related to abundance on the shelf.

## **Fishery**

Prior to 1977, few pollock were caught in the Donut Hole or Bogoslof region (Low and Ikeda 1978). Japanese scientists first reported significant quantities of pollock in the Aleutian Basin in the mid-to-late 1970's, but large-scale fisheries in the Donut Hole only began in the mid-1980's. By 1987 significant components of these catches were attributed to the Bogoslof Island region (Table 1b.1); however, the actual locations were poorly documented. The Bogoslof fishery primarily targeted winter spawning-aggregations but in 1992, this area was closed to directed pollock fishing.

In 1991, the only year with extensive observer data, the fishery timing coincided with the open seasons for the EBS and Aleutian Islands pollock fisheries (the Bogoslof management district was established in 1992 by FMP amendment 17). However, after March 23, 1991 the EBS region was closed to fishing and some effort was re-directed to the Aleutian Islands region near the Bogoslof district. In subsequent years, seasons for the Aleutian Islands pollock fishery were managed separately. Bycatch and discard levels were relatively low from these areas when there was a directed fishery (e.g., 1991). Updated estimates of pollock bycatch levels from other fisheries were small in recent years (Table 1b.2). The increase in pollock bycatch in 2010 (9 t in 2008 to 73 in 2009 and 176 t in 2010) can be attributed to the non-pelagic trawl arrowtooth flounder target fishery. The majority of pollock bycatch in the Bogoslof region continues to be occurring in the non-pelagic trawl arrowtooth flounder target fishery. For all fisheries there were 79 t of pollock catch in 2012 and 57 t in 2013.

## **Analytical approach**

For the purposes of this year's assessment, like last year's, a strictly survey-based management approach was selected. Previous assessments (e.g., Ianelli et al. 2004) developed a full-age structure model. In

those Ianelli et al. (2005) examined refinements to an age-structured model for Bogoslof pollock which included exploring the effect of Donut Hole catches in the 1980s on the stock assessment results. They assumed that 75% of the Donut Hole catches came from the Bogoslof stock, which is in accord with past practices of international pollock workshops (which used a range from 60 to 80%). However, concerns about this assumption were raised due to the uncertain degree of interchange between Bogoslof fish and central BS fish. In the SSC's December 2006 minutes they noted that additional research is needed to better understand the extent of these linkages. In lieu of new information on this becoming available, they recommended using Tier 5 as a basis for management, the maximum permissible ABC value would be 10,059 t (assuming  $M = 0.2$  and  $F_{ABC} = 0.75M = 0.15$ ):

$$ABC = B_{2012} \times M \times 0.75 = 67,063 \times 0.2 \times 0.75 = 10,059 t .$$

### **ABC Recommendation**

Maximum permissible ABC and OFL estimates for 2014 and 2015 under Tier 5 relies exclusively on the NMFS biennial acoustic-trawl survey biomass estimate. Since 2000, the values have varied between 292,000 t and 67,063 t. The most recent AT survey of the Bogoslof spawning stock was conducted in March of 2012 (McKelvey and Stienessen, in prep; Table 1b.3) and resulted in a biomass estimate of 67,063 t.

Our recommended ABC is based a Tier 5 calculation which results in 10,059 t for the next two years. The OFL for the Tier 5 calculation is 13,413 t.

An alternative approach that was suggested by the Plan Team for Tier 5 stocks was to use a process error model for estimating a mean over a number of surveys. Using that method indicates that the process-error approach would give nearly identical estimates as using the most recent survey (67,600 t of biomass for the process error model compared to 67,063 t; Fig. 1).

### **Ecosystem considerations**

In general, a number of key issues for ecosystem conservation and management can be highlighted.

These include:

- Preventing overfishing;
- Avoiding habitat degradation;
- Minimizing incidental bycatch (via multi-species analyses of technical interactions);
- Controlling the level of discards; and
- Considering multi-species trophic interactions relative to harvest policies.

For the case of pollock, the NPFMC and NMFS continue to manage the fishery on the basis of these issues in addition to the single-species harvest approach. The prevention of overfishing is clearly set out as a main guideline for management. Habitat degradation has been minimized in the pollock fishery by converting the industry to pelagic-gear only. Bycatch in the pollock fleet is closely monitored by the NMFS observer program, and individual species caught incidentally are managed on that basis. Discarding rates have been greatly reduced in this fishery and multi-species interactions is an ongoing research project within NMFS with extensive food-habit studies and simulation analyses to evaluate a number of "what if" scenarios with multi-species interactions.

As reported in Loughlin and Miller (1989) pups of Northern fur seals, *Callorhinus ursinus*, were first observed on Bogoslof Island in 1980. By 1988 the population had grown at a rate of 57% per year to over 400 individuals, including 80+ pups, 159 adult females, 22 territorial males, and 188 sub-adult males. They noted that the rookery is in the same location where solitary male fur seals were seen in 1976 and

1979 and is adjacent to a large northern sea lion rookery. On July 22, 2005 NMFS surveys resulted in counts of 1,123 adult males, a substantial increase over this time period (L. Fritz, AFSC, pers. comm.). The estimated number of Northern fur seal pups born on Bogoslof Island increased from 5,096 (SE = 33) to 12,631 (SE = 335) (Angliss and Allen, 2007). This suggests that conditions in the ecosystem have changed and appear to favor Northern fur seals. The extent that this is due to environmental conditions is unknown. However, pollock abundance may play only a small role since during peak abundance levels, the Northern fur seal abundance was at very low levels. Also, pollock are most concentrated in this region during winter months when Northern fur seals have migrated to more southern areas.

## Literature cited

- Angliss, R.P. and B.M. Allen. 2007. Northern fur seal (*Callorhinus ursinus*): Eastern Pacific Stock. NOAA-TM-AFSC-193. Alaska Fish. Sci. Cent., Natl. Mar. Fish. Serv., NOAA, 7600 Sand Point Way NE, Seattle WA 98115
- Bailey, K.M., T.J. Quinn, P. Bentzen, and W.S. Grant. 1999. Population structure and dynamics of walleye pollock, *Theragra chalcogramma*. *Advances in Mar. Biol.* 37:179-255.
- Barbeaux, S. J. Ianelli, and E. Brown. 2004. Stock assessment of Aleutian Islands region pollock. *In*: Stock assessment and fishery evaluation report for the groundfish resources of the Bering Sea/Aleutian Islands regions for 2005. North Pac. Fish. Mgmt. Council, Anchorage, AK, Appendix Section 1A.
- Ianelli, J.N., S. Barbeaux, G. Walters, T. Honkalehto, and N. Williamson. 2004. Bering Sea-Aleutian Islands Walleye Pollock Assessment for 2004. *In*: Stock assessment and fishery evaluation report for the groundfish resources of the Bering Sea/Aleutian Islands regions. North Pac. Fish. Mgmt. Council, Anchorage, AK, section 1:37-126.  
<http://www.afsc.noaa.gov/refm/docs/2004/EBSpollock.pdf>
- Ianelli, J.N., T. Honkalehto, and N. Williamson. 2005. An age-structured assessment of pollock (*Theragra chalcogramma*) from the Bogoslof Island Region. *In*: Stock assessment and fishery evaluation report for the groundfish resources of the Bering Sea/Aleutian Islands regions. North Pac. Fish. Mgmt. Council, Anchorage, AK, section 181-218,  
<http://www.afsc.noaa.gov/refm/docs/2005/Bogpollock.pdf>
- Ianelli, J.N., T. Honkalehto, and N. Williamson. 2006. An age-structured assessment of pollock (*Theragra chalcogramma*) from the Bogoslof Island Region. *In*: Stock assessment and fishery evaluation report for the groundfish resources of the Bering Sea/Aleutian Islands regions. North Pac. Fish. Mgmt. Council, Anchorage, AK, section 181-218,  
<http://www.afsc.noaa.gov/refm/docs/2006/Bogpollock.pdf>
- Loughlin, T.R. Miller, R.V. 1989. Growth of the northern fur seal colony on Bogoslof Island, Alaska. *Arctic*, v. 42, no. 4, Dec. 1989, p. 368-372.
- Low, L.L., and J. Ikeda. 1978. Atlas of groundfish catch in the Northeastern Pacific Ocean, 1964-1976. Northwest and Alaska Fisheries Center Data Report. 7600 Sand Point Way NE. Seattle WA. 166p.
- McKelvey, D. 2009. Results of the March 2009 echo integration-trawl survey of walleye pollock (*Theragra chalcogramma*) conducted in the southeastern Aleutian Basin near Bogoslof Island, Cruise DY2009-03. AFSC Processed Rep. 2009-05, 32 p. Alaska Fish. Sci. Cent., NOAA, Natl. Mar. Fish. Serv., 7600 Sand Point Way NE, Seattle WA 98115.
- McKelvey, D. and S. Stienessen. 2012. Results of the March 2012 acoustic-trawl survey of walleye pollock (*Theragra chalcogramma*) conducted in the southeastern Aleutian Basin near Bogoslof Island, Cruise DY2012-02. <http://www.afsc.noaa.gov/Publications/ProcRpt/PR2012-08.pdf>.

Smith, G.B. 1981. The biology of walleye pollock. *In* Hood, D.W. and J.A. Calder, The Eastern Bering Sea Shelf: Oceanography and Resources. Vol. I. U.S. Dep. Comm., NOAA/OMP 527-551.

Wespestad, V. G. and J. M. Terry. 1984. Biological and economic yields for eastern Bering Sea walleye pollock under differing fishing regimes. *N. Amer. J. Fish. Manage.*, 4:204-215.

Table 1b.1 Catch in tons from the Donut Hole and the Bogoslof Island area, 1977-2013.

<b>Year</b>	<b>Donut Hole (t)</b>	<b>Bogoslof Island (t)</b>	<b>Total (t)</b>
1977		11,500	11,500
1978		9,600	9,600
1979		16,100	16,100
1980		13,100	13,100
1981		22,600	22,600
1982		14,700	14,700
1983		21,500	21,500
1984	181,200	22,900	204,100
1985	363,400	13,700	377,100
1986	1,039,800	34,600	1,074,400
1987	1,326,300	377,436	1,703,736
1988	1,395,900	87,813	1,483,713
1989	1,447,600	36,073	1,483,673
1990	917,400	151,672	1,069,072
1991	293,400	316,038	609,438
1992	10,000	241	10,241
1993	1,957	886	2,843
1994		556	556
1995		334	334
1996		499	499
1997		163	163
1998		8	8
1999		29	29
2000		29	29
2001		258	258
2002		1,042	1,042
2003		24	24
2004		<1	<1
2005		<1	<1
2006		<1	<1
2007		<1	<1
2008		9	9
2009		73	73
2010		176	176
2011		173	173
2012		79	79
2013		57	57

Table 1b.2. Estimated retained, discarded, and total pollock catch (t) from the Bogoslof region. Source: NMFS Regional office Blend database and catch accounting system.

Year	Discarded	Retained	Total
1991	20,327	295,711	316,038
1992	240	1	241
1993	308	578	886
1994	11	545	556
1995	267	66	334
1996	7	492	499
1997	13	150	163
1998	3	5	8
1999	11	18	29
2000	20	10	29
2001	28	231	258
2002	12	1,031	1,042
2003	19	5	24
2004	< 1		< 1
2005	< 1	< 1	< 1
2006	< 1	< 1	< 1
2007	< 1	< 1	< 1
2008	< 1	9	9
2009	6	67	73
2010	53	124	176
2011	23	150	173
2012	5	74	9
2013	< 1	56	57

Table 1b.3. Biomass (tons) of pollock as surveyed in the Bogoslof region, 1988-2012. For additional details see McKelvey and Stienessen (2012).

Year	Survey biomass estimates (t)	Survey area (nmi <sup>2</sup> )	Relative error
1988	2,395,737	NA	22%
1989	2,125,851	NA	22%
1990		No survey	
1991	1,289,006	8,411	12%
1992	940,198	8,794	20%
1993	635,405	7,743	9%
1994	490,077	6,412	12%
1995	1,104,124	7,781	11%
1996	682,277	7,898	20%
1997	392,402	8,321	14%
1998	492,396	8,796	19%
1999	475,311	NA	22%
2000	301,402	7,863	14%
2001	232,170	5,573	10%
2002	225,712	2,903	12%
2003	197,851	2,993	22%
2004		No survey	
2005	253,459	3,112	17%
2006	240,059	1,803	12%
2007	291,580	1,870	12%
2008		No survey	
2009	110,191	1,803	19%
2010		No survey	
2011		No survey	
2012	67,063	3,656	10%




Figure 1. Distribution of pollock biomass (t) observed along transects during the winter 2012 acoustic-trawl survey. Transect numbers are underlined; trawl haul locations are indicated by circles.






Figure 2. Bogoslof Island pollock survey estimates fitted to a process error model for averaging recruitment. The shade represents the approximate 90% confidence interval from the model.