

Assessment of the Grenadier Stock Complex in the Gulf of Alaska, Eastern Bering Sea, and Aleutian Islands

Cara Rodgveller and Pete Hulson

November 2014

Executive Summary

The Secretary of Commerce approved Amendments 100/91 on August 6, which added the grenadier complex into both FMPs as Ecosystem Components. Under this rule, they are not allowed to be targeted but there is an 8% Maximum Retainable Allowance (MRA) (Federal Register, Proposed Rules, Vol. 79, No. 93). The final rule will publish before the end of the year and so it may be effective for the start of the 2015 fishing year.

As an Ecosystem Component, a stock assessment is not required and there is no ABC or OFL. A full unofficial assessment report was prepared for grenadiers in even years since 2006, even though they were “nonspecified”. For 2015, we are presenting an abbreviated SAFE report for the BSAI and GOA combined for the purpose of tracking trends in abundance. This content of future reports is still being evaluated since a SAFE report is not required. This report contains a time series of catch and abundance estimates and unofficial ABC and OFL values based on Tier 5 calculations. These values are not used for management or for determining if overfishing is occurring for Ecosystem Component species/complexes. There is no definition of overfishing for an Ecosystem Component.

Summary of Changes in Assessment Inputs

Changes in the input data: New data inputs include: 1) updated catch data for 2003-2014; 2) updated 2000-2014 Aleutian Island (AI) biomass from 1-1,000 m using the estimation method presented in the 2012 SAFE; 3) NMFS longline survey results for 2013 and 2014; 4) updated GOA biomass using a random effects model. There was no EBS slope trawl survey in 2014.

Changes in assessment methodology: This year we use a random effects model (a similar method, a Kalman filter, was presented in the 2012 SAFE report (Rodgveller et al. 2012)), that utilizes trawl survey data from 1984-2013 to estimate the exploitable biomass in 2013. Since there was no trawl survey in the GOA in 2014, the estimate for 2013 is used as the most recent value of exploitable biomass.

Summary of Results

For 2015, the maximum allowable ABC for the BSAI is 75,274 t and for the GOA is 30,691 t. This ABC is a 12% increase for the BSAI and a 12% decrease for the GOA. The corresponding reference values for grenadier are summarized in the following tables, with the recommended ABC and OFL values in bold. Overfishing is not occurring in either the BSAI or GOA.

Gulf of Alaska Grenadiers

Quantity	As estimated or specified <i>last year for</i> ^a :		As estimated or recommended <i>this year</i> for:	
	2014	2015	2015	2016
<i>M</i> (natural mortality)	0.078	0.078	0.078	0.078
Specified/recommended Tier	5	5	5	5
Biomass (t)	597,884	597,884	524,624	524,624
F_{OFL} (F=M)	0.078	0.078	0.078	0.078
$maxF_{ABC}$ (maximum allowable = $0.75x F_{OFL}$)	0.0585	0.0585	0.0585	0.0585
F_{ABC}	0.0585	0.0585	0.0585	0.0585
OFL (t)	46,635	46,635	40,921	40,921
maxABC (t)	34,976	34,976	30,691	30,691
ABC (t)	34,976	34,976	30,691	30,691
Status	As determined <i>last year</i> for:		As determined <i>this year</i> for:	
	2012	2013	2013	2014
Overfishing	No	n/a	No	n/a

^aThe values for biomass, OFL, and ABC in these two columns are based on Rodgveller and Hulson 2013. They are an average of the last three trawl surveys that sampled down to 1,000 m. The current values (for 2015 and 2016) are from the random effects model fit to survey biomass by region and depth strata.

These are unofficial ABC and OFL values since grenadier are an Ecosystem Component, which do not have ABCs or OFLs.

Bering Sea and Aleutian Islands Grenadiers

Quantity	As estimated or specified <i>last year for</i> ^a :		As estimated or recommended <i>this year</i> for:	
	2014	2015	2015	2016
<i>M</i> (natural mortality)	0.078	0.078	0.078	0.078
Specified/recommended Tier	5	5	5	5
Biomass (t)	1,152,284	1,152,284	1,286,734	1,286,734
F_{OFL} (F=M)	0.078	0.078	0.078	0.078
$maxF_{ABC}$ (maximum allowable = $0.75x F_{OFL}$)	0.0585	0.0585	0.0585	0.0585
F_{ABC}	0.0585	0.0585	0.0585	0.0585
OFL (t)	89,878	89,878	100,365	100,365
maxABC (t)	67,409	67,409	75,274	75,274
ABC (t)	67,409	67,409	75,274	75,274
Status	As determined <i>last year</i> for:		As determined <i>this year</i> for:	
	2012	2013	2013	2014
Overfishing	No	n/a	No	n/a

^aThe values for biomass, OFL, and ABC in these two columns are based on Rodgveller and Hulson 2013.

These are unofficial ABC and OFL values since grenadier are an Ecosystem Component, which do not have ABCs or OFLs.

Tier 5 computations for giant grenadier OFL and ABC are summarized as follows (AI = Aleutian Islands, EBS = Eastern Bering Sea, GOA = Gulf of Alaska; biomass, OFL, and ABC are in mt) for 2015:

BSAI and GOA grenadiers						
Area	Biomass	Natural mortality M	OFL definition	OFL	ABC definition	ABC
EBS	553,557	0.078	biom x M	43,177	OFL x 0.75	32,383
AI	733,177	0.078	biom x M	57,188	OFL x 0.75	42,891
BSAI total	1,286,734			100,365		75,274
GOA	524,624	0.078	biom x M	40,921	OFL x 0.75	30,691
Grand total	1,811,358			141,286		105,965

These are unofficial ABC and OFL values since grenadier are an Ecosystem Component, which do not have ABCs or OFLs.

The specifications in the GOA for 2015 differ from last year because a random effects model fit to the survey biomass was used as a proxy for the exploitable biomass in this year's assessment (following the recommendation of the Survey Averaging Working group). In the BSAI the ABC and OFL include the AI biomass estimated using the method presented in the 2012 SAFE report (Rodgveller et al. 2012). Further discussion of this method is below under SSC comments and under the Survey Data section. Catches are not approaching unofficial OFLs.

Summaries for Plan Team

Species	Year	BSAI Biomass	BSAI ABC	BSAI Catch ¹	GOA Biomass	GOA ABC	GOA Catch ¹	Total Catch ¹
grenadiers	2013	1,733,797	101,427	4,164	597,884	34,976	11,339	15,504
	2014	1,152,284	89,878	2,627	597,884	34,976	5,236	7,863
	2015	1,286,734	75,274		524,624	30,691		
	2016	1,286,734	75,274		524,624	30,691		

¹Current as of October 7, 2014. Source: NMFS Alaska Regional Office Catch Accounting System via the Alaska Fisheries Information Network (AKFIN) database (<http://www.akfin.org>).

SSC and Plan Team Comments on Assessments in General

There were no comments on assessments in general that pertains to this assessment.

SSC and Plan Team Comments Specific to this Assessment

Here we report comments from the SSC in 2012 and 2013, since there was an executive summary in 2013. We also present comments from 2013 regarding the EA/RIR/IRFA on grenadier that pertain to

stock assessment. Responses to these comments and additions to the document were also made in the final EA/RIR/IRFA document (NMFS 2014.)

“The authors introduced a new method for determining AI biomass and variance estimates. The SSC cautions that this is an uncertain extrapolation method. The catchability and size selection of longline surveys is known to differ from the trawl survey. This method assumes that the ratio between longline and trawl surveys in shallow water will be the same for the ratio of longline and trawl surveys in deep water. The SSC encourages the authors to verify whether this assumption is valid.” (SSC, 2012)

The primary problem with using the AI trawl survey biomass estimates for giant grenadier is that the survey does not sample deeper than 500 m; where the majority of the giant grenadier population can be found. To account for the missing biomass from the trawl survey an expansion method is needed, for which we use the AFSC longline survey data, the only survey that samples deeper than 500 m in the AI.

The primary uncertainty associated with this method centers on the use of a ratio estimator between trawl survey biomass and longline survey RPWs. The ratio between trawl survey biomass and longline survey RPWs is assumed to be the same in shallow depths (1-500 m, for which we have trawl survey data) and deep depths (500-1000 m, for which we do not have trawl survey data), an assumption that must be made due to the available data. There may be uncertainty associated with extrapolating trawl survey biomass in this manner. Our opinion is that it is important to present estimates of deep-water biomass so that a better reflection of the potential grenadier biomass in the AI can be presented.

A comparison of the ratio of longline and trawl survey data in shallow- and deep-waters in the Gulf of Alaska is presented in an attempt to verify the assumption of constant catchability in each survey in shallow- and deep-water (See “Biomass estimation in the AI” under the “Trawl Surveys” section). Ratios between the two surveys in shallow- and deep-water were almost identical in the GOA, indicating that the assumption is valid.

The trends in RPWs and biomass were compared in the EBS and the GOA to determine if the two surveys are sampling the same population. Trends in the EBS tracked well, but trends in the GOA were not consistent among the two surveys (See “RPWs” under the “Longline surveys” section), so this did not validate nor contradict the effect of the differing selectivities of the two surveys.

“In response to SSC comments, the authors included a Kalman filter model for estimating biomass. The Kalman filter estimates miss the most recent trawl biomass estimate in the GOA resulting in a substantially lower biomass estimate. For future assessments, the SSC encourages continued exploration of the Kalman filter method and we ask the authors to consider the recommendations in the Plan Team survey averaging work group (SSC, Dec. 2012).”

The survey averaging working group has recommended that for Tier 5 stocks authors should compare biomass estimates using random effects models to the standard calculations. We compared status quo to a Kalman filter in 2012 (Rodgveller et al. 2012). Here we use the random effects model to estimate GOA biomass for 2013. Approximately ½ of the grenadier biomass was in the deepest stratum (701-1,000 m) and this stratum was not sampled in 2011 or 2013. It is likely that it will not be sampled in 2015. The random effects model provides a method to incorporate all available trawl survey data since 1984. See the section titles “Biomass Estimation in the GOA” under the “Surveys” heading for details.

“As a non-specified species complex the Plan Teams and the SSC are not required to provide harvest specifications for this species group. Given the potential that grenadier management may change in 2014, the SSC requests a full assessment next year.” (SSC, Dec. 2013)

Because grenadiers were put into the FMPs as an Ecosystem Component in 2014, there are still no harvest specifications and no requirement to produce SAFE reports. Therefore, instead of presenting a full assessment, we are presenting an abbreviated assessment that contains a time series of catch and survey data and descriptions of new methodologies used for estimating biomass.

Comments on the EA/RIR/IRFA in 2013

“The SSC reviewed the document and concluded that it is very well done and ready for release for public review. However, the SSC identified several areas where the document could be improved and requests that staff strive to make these improvements prior to release.” (SSC, Dec. 2013)

“It would be useful to develop a food web for the slope regions as part of the ecosystem concerns chapter...” (SSC, Dec. 2013)

Little information is available on food web and habitat interactions between grenadiers and other groundfish. The information that is available indicates that in the Aleutian Islands, the diet of grenadiers is comprised mostly squid and bathypelagic fish (myctophids) (Yang 2003), whereas in the Gulf of Alaska, squid and pasiphaeid shrimp predominated as prey (Yang et al. 2006). Thus, other groundfish do not appear to compose the prey field of grenadiers.

“The 2012 appendix revealed strong spatial partitioning of the sexes by depth. The SSC requests the author to estimate the sex ratio for survey biomass estimates in the assessment. The SSC requests that, if possible, the document should provide trawl and longline survey biomass estimates by sex and depth.” (SSC, Dec. 2013)

In response to these comments, an appendix was added to the EA/RIR/IRFA with tables and figures that broke out catch, biomass, and RPWs by sex and depth (NMFS, 2013; see pages 77-82). We include a summary of that analysis as well as the data tables and figures presented in the EA/RIR/IRFA as an appendix to this SAFE report.

“With respect to depth, the SSC requests that the document includes a short discussion of the potential uncertainty associated with the expansion method used to estimate grenadier biomass at deeper depths in the AI.” (SSC, Dec. 2013)

See SSC comment from 2012 (above).

“The SSC also encourages the author to address comments and suggestions made by the non-target CIE review team if they are relevant to the grenadier appendix.” (SSC, Dec. 2013)

See below.

Center of Independent Expert Review Comments

In May, 2013 there was a Center of Independent Expert (CIE) review of non-target assessments at the AFSC. Three reviewers participated and each produced a report without collaboration from NMFS or other reviewers. Here I will summarize the comments pertaining to grenadier. General comments about the tier system are excluded.

- 1) The three CIE reviewers did not consider the estimates of absolute biomass to be reliable for grenadier, or for other reviewed species. The catchabilities of grenadier by the bottom trawl surveys in the EBS, AI, and GOA are unknown, but are currently assumed to be 1. One reviewer recommends using expert knowledge to estimate the availability, vulnerability, and density in trawlable and untrawlable grounds to explore converting the trawl survey data to absolute biomass.**

There is some evidence, based on diets and anecdotal evidence of higher catch rates when longline gear is held off off-bottom by rocky topography, that grenadier spend time off-bottom. If this is true, q is likely <1 for trawl gear. There have been no studies to estimate q for grenadier.

- 2) There were concerns over the ratio that is used to extrapolate the relative population numbers and weights in the western AI from a ratio of western to eastern AI data from the 1980's.**

The western AI has not been sampled by the domestic NMFS longline survey. Data from Japanese surveys are used to extrapolate western AI abundance. This method is used for all species, including sablefish, since it is the only available data.

- 3) The reviewers supported using models for estimating biomass, such as a Kalman filter or random effects model.**

This year we use a random effects model to estimate GOA biomass.

- 4) A reviewer said that maximum age methods used for estimating natural mortality are acceptable, but recommended for swapping otoliths with other labs.**

The AFSC lab is the only lab that ages giant grenadier. A method was developed by Charles Hutchinson in the Age and Growth Laboratory for a previous age at maturity study (Rodgveller et al. 2010). Ages could not be validated using C14. This is likely because young fish are not in shallower waters (<200 m). Young fish are needed to confirm annuli at age 1-5; however, no young fish have been found in surveys or fisheries.

Introduction

Grenadiers (family Macrouridae) are deep-sea fishes related to hakes and cods that occur world-wide in all oceans. Also known as “rattails”, they are especially abundant in waters of the continental slope, but some species are found at abyssal depths. At least seven species of grenadier are known to occur in Alaskan waters, but only three are commonly found at depths shallow enough to be encountered in commercial fishing operations or in fish surveys: giant grenadier (*Albatrossia pectoralis*), Pacific grenadier (*Coryphaenoides acrolepis*), and popeye grenadier (*Coryphaenoides cinereus*) (Mecklenburg et al. 2002). Of these, giant grenadier has the shallowest depth distribution and the largest apparent biomass, and hence is by far the most frequently caught grenadier in Alaska. Because of this importance, this report will emphasize giant grenadier, but it will also discuss the other two species.

Distribution: Giant grenadier range from Baja California, Mexico around the arc of the north Pacific Ocean to Japan, including the Bering Sea and the Sea of Okhotsk (Mecklenburg et al. 2002), and they are also found on seamounts in the Gulf of Alaska and on the Emperor Seamount chain in the North Pacific (Clausen 2008). In Alaska, they are especially abundant on the continental slope in waters >400 m depth.

These fish are the largest in size of the world's grenadier species (Iwamoto and Stein 1974); maximum weight of one individual in a Bering Sea trawl survey was 41.8 kg¹.

Speciation: Previous publications (Clausen 2006 and 2008) speculated that more than one species of giant grenadier may exist in Alaska because two morphs of the fish have been observed based primarily on the relative size of the eye to the head, as well as three very different patterns of otolith morphology. Tissue and otoliths samples were collected on the AFSC longline survey in 2013 for a more definitive analysis of speciation, stock structure, and otolith morphometrics.

Biology: There is some known biological information on adult giant grenadier, but data on larvae and juvenile grenadiers is nonexistent. The spawning period is thought to be protracted and may even extend throughout the year (Novikov 1970; Rodgveller et al. 2010). Two papers provide purported descriptions of larvae of giant grenadier in the North Pacific (Endo et al. 1993; Ambrose 1996), but Busby (2004) points out that these descriptions appear so different that they probably represent separate species. At any rate, no larvae have ever been collected in Alaska that correspond to either of these descriptions or to the description of a third form (Busby 2004) that is also giant grenadier-like². Small, juvenile fish less than ~15-20 cm pre-anal fin length (PAFL) are virtually absent from bottom trawl catches (Novikov 1970; Ronholt et al. 1994; Hoff and Britt 2009, 2011), and juveniles may be pelagic in their distribution. (Because the long tapered tails of grenadiers are frequently broken off when the fish are caught, PAFL is the standard unit of length measurement for these fish. PAFL is defined to be the distance between the tip of the snout and the insertion of the first anal fin ray). Bottom trawl studies indicate that females and males have different depth distributions, with females inhabiting shallower depths than males. For example, both Novikov (1970) in Russian waters and Clausen (2008) in Alaskan waters found that nearly all fish <600 m depth were female, and the Novikov study was based on trawl sampling throughout the year. Presumably, some vertical migration of one or both sexes must occur for spawning purposes; Novikov (1970) speculates that females move to deeper water inhabited by males for spawning.

Ecology: The habitat and ecological relationships of giant grenadier are likewise little known and uncertain. Clearly, adults are often found in close association with the bottom, as evidenced by their large catches in bottom trawls and on longlines set on the bottom. However, based on a study of the food habits of giant grenadier off the U.S. west coast, Drazen et al. (2001) concluded that the fish feeds primarily in the water column. Most of the prey items found in the stomachs were meso- or bathypelagic squids and fish, and there was little evidence of benthic feeding. Smaller studies of giant grenadier food habits in the Aleutian Islands (Yang 2003) and Gulf of Alaska (Yang et al. 2006) showed similar results. In the Aleutian Islands, the diet comprised mostly squid and bathypelagic fish (myctophids), whereas in the Gulf of Alaska, squid and pasiphaeid shrimp predominated as prey. The hypothesis regarding the tendency of the fish to feed off bottom is supported by observations of sablefish longline fishermen, who report that their highest catches of giant grenadier often occur when the line has been inadvertently "clothes-lined" between two pinnacles, rather than set directly on the bottom³. Pacific sleeper sharks (*Somniosus pacificus*) and Baird's beaked whales (*Berardius bairdii*) have been documented as predators on giant grenadier (Orlov and Moiseev 1999; Walker et al. 2002). Sperm whales (*Physeter macrocephalus*) are another likely predator, as they are known to dive to depths inhabited by giant

¹ G. Hoff, National Marine Fisheries Service, Alaska Fisheries Science Center, RACE Division, 7600 Sand Point Way NE, Seattle WA 98115. Pers. comm. March 2005.

² M. Busby, National Marine Fisheries Service, Alaska Fisheries Science Center, RACE Division, 7600 Sand Point Way NE, Seattle WA 98115. Pers. comm. October 2006.

³ D. Clausen, National Marine Fisheries Service, Alaska Fisheries Science, Auke Bay Laboratories, 17109 Point Lena Loop Rd., Juneau, AK 99801. Pers. observ. October 2004.

grenadier on the continental slope and have been observed in Alaska depredating on longline catches of giant grenadier⁴.

Distribution of Pacific and popeye grenadier: Pacific grenadier have a geographic range nearly identical to that of giant grenadier, i.e., Baja California, Mexico to Japan. Popeye grenadier range from Oregon to Japan. Compared to giant grenadier, both species are much smaller and generally found in deeper water. They appear to be most abundant in waters >1,000 m, which is deeper than virtually all commercial fishing operations and fish surveys in Alaska. For example, in a recent experimental longline haul in the western Gulf of Alaska at a depth of 1400-1500 m, 56% of the hooks caught Pacific grenadier⁵. This indicates that at least in some locations in deep water, abundance of Pacific grenadier in Alaska can be extremely high. Few popeye grenadier are caught on longline gear, apparently because of the relatively small size of these fish, and most of the information on this species comes from trawling. Food studies off the U.S. West Coast indicate that Pacific grenadier are more benthic in their habitat than are giant grenadier, as the former species fed mostly on bottom organisms such as polychaetes, mysids, and crabs (Drazen et al. 2001).

Evidence of stock structure: Stock structure and migration patterns of giant grenadier in Alaska are unknown, as no genetics studies have been done (except for brief genetic investigation of the two morphs of this species that was previously mentioned), and the fish cannot be tagged because all individuals die due to barotrauma when brought to the surface. One study in Russian waters, however, used indirect evidence to conclude that seasonal feeding and spawning migrations occur of up “to several hundred miles” (Tuonogov 1997).

Natural mortality: In the 2014 assessment we continue to use the natural mortality estimate (M) of 0.078, calculated using Hoenig’s (1983) longevity equation with a maximum age of 58 from a study of age at maturity for giant grenadier (Rodgveller et al. 2010). A discussion of the four methods employed by Rodgveller et al. (2010) and the reason for choosing Hoenig’s (1983) method can be found in the 2010 grenadier SAFE (Clausen and Rodgveller 2010). Giant grenadier greater than 60 cm PAFL have been caught on the AFSC longline survey, whereas the greatest length in the age samples was 53 cm (Rodgveller et al. 2010). Therefore, it is probable that fish older than 58 exist. An older maximum age would result in a decrease in M . Because fish older than 58 years may exist, we suggest revisiting the determination of M for giant grenadier if more age samples become available in the future.

Fishery

Catch History

Catches since 1997 have been estimated for the eastern Bering Sea (EBS), Aleutian Islands (AI), and GOA based largely on data from the Alaska Fishery Science Center’s Fishery Monitoring and Analysis program. The estimates for 1997-2002 were determined by simulating the catch estimation algorithm used for target species by the NMFS Alaska Regional Office in what was formerly called their “blend catch estimation system” (Gaichas 2002 and 2003). Although these estimates may not be as accurate as the official catch estimates determined for managed groundfish species, they are believed to be the best possible based on the data available. They do not appear unreasonable compared to the official catches of other species caught along with giant grenadier on the continental slope in Alaska, such as sablefish and

⁴ C. Lunsford, National Marine Fisheries Service, Alaska Fisheries Science Center, Auke Bay Laboratories, 17109 Point Lena Loop Rd., Juneau, AK 99801. Pers. comm. October 2006.

⁵ D. M. Clausen and C. J. Rodgveller, 2010. Deep-water longline experimental survey for giant grenadier and sablefish in the western Gulf of Alaska, August 2008. National Marine Fisheries Service, Alaska Fisheries Science Center, Auke Bay Laboratories, 17109 Point Lena Loop Rd., Juneau, AK 99801. Unpubl. manuscr. 23p.

Greenland turbot. The estimates for 2003-2014 were computed by the NMFS Alaska Regional Office based on their Catch Accounting System, which replaced the “blend” system in 2003. All the data are presented as “grenadiers, all species combined”.

In 2013 the observer program underwent restructuring. Implementation of this program is considered an improvement over the previous observer system and an analysis of the first year under the restructured program was presented at the June 2014 council meeting (Faunce et al. 2014). It is too early to determine if there are any changes in the time series due to this restructuring. More years of data are needed. There is now observer coverage of vessels <60 ft and of the IFQ Pacific halibut fleet. A description of changes in bycatch rate calculations and changes that occurred in CAS that relate to non-target species can be found in Tribuzio et al. (2014).

Overall, the estimate of total catch of grenadier in 2013 (15,504 mt) was almost the same as in 2012 (15,119 mt) (Table 1). Catch in 2013 was up 13% from the 2003-2012 average. Even though the great majority of grenadier catch occurs by Oct. 1, the catch estimate in 2014 is only 7,863 mt. For example, by Oct. of 2013 95% of the catch was taken. Thus we expect that the final catch estimate for 2014 will be much lower than average. It is possible that this is related to observer restructuring.

In the BSAI catch was down 21% (1,092 mt) from average in 2013 (Table 1). This was primarily in the Greenland turbot fishery in the BS and the Kamchatka flounder fishery in the AI (Table 2). In the GOA, grenadier catch was higher in 2013 than in previous years (2,890 mt above average; 34%; Table 1). The majority of this increase was in rockfish (461 mt; 84% above average), sablefish (1,476 mt; 22% above average) (Table 2), and the deep-water flatfish fisheries (1,063 mt; 581% above average) (Table 2). Surprisingly, the estimate of grenadier catch in the halibut target group was 47% below average in 2013; although, catch estimates in this target group have always been variable.

Survey Data

Trawl Surveys

Biomass estimation in GOA: The Plan Team Survey Averaging Plan Team working group and the SSC suggested that for tier 5 species authors should compare biomass estimates using a random effects model to standard calculations. For grenadier a Kalman filter model was presented in 2012 (Rodgveller et al. 2012). This year we used a random effects model to estimate exploitable biomass for unofficial ABCs and OFLs (Table 3). The only GOA trawl surveys that extended to 1,000 m include the surveys in 1984, 1987, 1999, 2005, 2007, and 2009. In 1990, 1993, 1996, and 2001 the trawl survey only sampled depths down to 500 m, and in 2003 and 2011 the trawl survey sampled depths up to 700 m. Due to the differences in the depth sampled among the various trawl surveys, and the distribution of giant grenadier biomass across depth strata, we applied the random effects model to biomass estimates for the 1-500 m, 501-700 m, and 701-1000 m depth strata by region separately. This resulted in three time series of biomass estimates: one for each depth stratum. The full time series of biomass estimates in the GOA from the random effects model were then obtained by summing the biomass estimates across the three depth strata (Figure 1). Biomass in the GOA, estimated using the random effects model, increased until 2005 and then decreased slowly and then increased in 2013 (Figure 1). Compared to status quo, biomass estimates from the random effects model are lower (Table 3) (ratios of random effects biomass/status quo biomass range from 0.76 to 0.97).

Biomass estimation in AI: The Aleutian Islands have presented a special problem for biomass estimation because no trawl surveys since 1986 have sampled waters deeper than 500 m, where most giant grenadier biomass is found. In previous SAFEs (Clausen 2006; Clausen and Rodgveller 2008, Clausen and

Rodgveller 2010) an AI biomass was estimated by using a combination of data from other areas and surveys: the GOA and EBS slope trawl surveys and the AFSC longline survey (Clausen and Rodgveller, 2010).

In 2012, a new method was used to estimate giant grenadier biomass that utilizes only AI survey data (Rodgveller et al. 2012; Appendix 1A and in the “survey data” section), and we continue to use this method. The AI trawl survey biomass estimates from the “shallow” depths, which are regularly sampled (1-500 m), and AI longline survey RPWs from “shallow” (200-500 m) and “deep” depths (501-1000 m) are used to estimate the total AI biomass using the following equation:


$$(1) B_y = \bar{r}W_y$$

where B_y is the total biomass in year y , \bar{r} is the ratio of the sum of bottom trawl survey biomass estimates to the sum of longline RPWs in the shallow depth stratum for years when both surveys occurred (2000, 2002, 2004, 2006, 2010, 2012, 2014), and W_y is the total RPW in year y . \bar{r} of “shallow” biomass to “shallow” RPWs for these years was 0.223. For those years when the AFSC longline survey occurs in the AI, an AI biomass estimate is now available (Table 4). Estimated biomass increased in 2014 by 53%. It is now similar to the estimated biomass in 2010.

When using this method, there is an assumption that the ratio between longline and trawl surveys in shallow water is the same as the ratio of longline and trawl surveys in deep water. In an attempt to validate this assumption, we examined the ratios of the two surveys in shallow- and deep-water in the GOA, where the trawl survey sampled down to 700 m in 2003 and 2011 and to 1,000 m in 1999, 2005, 2007, and 2009. In the GOA, the shallow ratio was nearly the same as the deep ratio (using the same method employed for calculating the AI shallow ratio), indicating that the assumption that the shallow- and deep-water ratios in the AI are similar is likely valid.

max depth	years	shallow ratio	deep ratio
700 m	2003,2011	0.60	0.62
1,000 m	1999,2005,2007,2009	0.52	0.51
all years		0.60	0.52

There is some evidence that trawl and longline survey abundance trends are similar. This may indicate that these surveys are sampling the same population and lend credence to the method we use to extrapolate AI biomass from longline survey data. Trawl and longline survey abundance trends can be compared in the GOA and EBS, since some trawl surveys sampled to at least 1,000 m in these areas. Longline and trawl surveys did not occur in the same years in the EBS (trawl survey was in even years and longline was in odd). The trends in the two surveys in the EBS tracked well. The trends in the GOA were not similar; however, there were not many trawl surveys in the GOA that sampled down to 1,000m within a short time frame (4 surveys over 10 years) vs. 5 years in the EBS.



Biomass in the EBS: There was no slope trawl survey in 2014. Biomass point estimates have ranged from 426–660 thousand mt between 2002 and 2012 (Table 4). Biomass was almost identical to the average in 2012 (550,266 mt), which was 17% lower than the estimate in 2010.

Longline Surveys

RPWs: RPWs of giant grenadier in the GOA had a general decreasing trend from 1999 through 2004, increased through 2007, and have been somewhat decreasing since then (Table 5). In 2014 the RPW in the GOA was 19% below average, which is well within the range of values in other years. RPWs in the Bering Sea have been increasing since 2007. In 2013 the biomass was 7% above average. The biomass of giant grenadier in the AI is larger than in other areas. This is because there is a large population estimated to be in the western Aleutians. This area is not currently sampled, but a ratio of eastern to western areas from previous surveys is used to extrapolate RPWs to these areas. In 2014, the AI RPW was 26% above average.

Analytic Approach

Modeling Approach

The tier 5 computations have been based only on giant grenadier because virtually none of the other species are caught in the commercial fishery or surveys. The exploitable biomass in the GOA was previously based on averaging the biomass estimates in the last three trawl surveys that extended to 1,000 m. The deepest stratum (701–1,000 m) was not sampled in 2011 or 2013, therefore, since 2009 the same estimate of biomass has been used for ABC/OFL calculations. This year we use a random effects model, presented in the 2012 SAFE report (Rodgveller et al. 2012), which utilizes trawl survey data from 1984–2013 to estimate the exploitable biomass in 2013 (see section above under Survey data, trawl surveys). Since there was no trawl survey in the GOA in 2014, the estimate for 2013 is used as the most recent value of exploitable biomass. To estimate exploitable biomass in the BS and AI, we continue to use an average of the three most recent trawl surveys with data available down to 1,000 m. For the BS that is 2008, 2010, and 2012. For the AI that is 2010, 2012, 2014. In the future we may use a random effects model in the Bering Sea to estimate biomass.

Parameter Estimates

Maximum Age: The most recent aging studies for giant grenadiers (Burton 1999 and Rodgveller et al. 2010) found the maximum age to be 56 and 58 years, respectively, based on specimens from the GOA. There have been no aging studies for Pacific grenadier in Alaska, but Andrews et al. (1999) found a maximum age of 73 years for this species off the U.S. west coast.

Natural mortality: In the 2014 assessment we continue to use the natural mortality estimate (M) of 0.078, calculated using Hoenig's (1983) longevity equation with a maximum age of 58 from a study of age at maturity for giant grenadier (Rodgveller et al. 2010).

Results

Harvest Recommendations

Parameters used: In the previous stock assessment for grenadiers (Rodgveller et al. 2013), the NPFMC’s tier 5 approach for determining the OFL and ABC was recommended, and this approach was supported by both the GOA Groundfish Plan Team and the NPFMC’s Scientific and Statistical Committee. We again use tier 5 unofficial ABC and OFL calculations.

Methods: Current biomass estimates in this assessment for giant grenadier in the EBS and GOA were calculated based on the average of the three most recent deep-water trawl surveys that sampled down to 1,000 or 1,200 m. In the EBS, these are now the 2008, 2010, and 2012 surveys. In the AI a method used in the 2012 SAFE was used to calculate biomass down to 1,000 m, even when trawl surveys sampled only to 500 m. Details are in Rodgveller et al. (2012). Estimates of AI biomass used for calculations of ABC and OFL are now based on 2010, 2012, and 2014.

The current GOA biomass was estimated using a random effects model (see section above under “Trawl Surveys”); therefore a single estimate is used as an estimate of exploitable biomass. This method is preferable to averaging the last three trawl surveys that sampled down to 1,000 m because trawl surveys have not extended this deep since 2009. The deepest stratum (700-1,000 m) will not likely be sampled in the near future.

The NPFMC’s tier 5 definitions for OFL and ABC are: $OFL = M \times Biomass$, where M is the estimated natural mortality rate, and $ABC \leq (0.75 \times OFL)$. Based on our discussion above, tier 5 recommendations for OFL and ABC of grenadiers are listed below (biomass, OFL, ABC, and mean catch are in mt).

BSAI and GOA grenadiers						
Area	Biomass	Natural mortality M	OFL definition	OFL	ABC definition	ABC
EBS	553,557	0.078	biom x M	43,177	OFL x 0.75	32,383
AI	733,177	0.078	biom x M	57,188	OFL x 0.75	42,891
BSAI total	1,286,734			100,365		75,274
GOA	524,624	0.078	biom x M	40,921	OFL x 0.75	30,691
Grand total	1,811,358			141,286		105,965

Gulf of Alaska Grenadiers

Quantity	As estimated or specified <i>last year for</i> ^a :		As estimated or recommended <i>this year</i> for:	
	2014	2015	2015	2016
<i>M</i> (natural mortality)	0.078	0.078	0.078	0.078
Specified/recommended Tier	5	5	5	5
Biomass (t)	597,884	597,884	524,624	524,624
F_{OFL} (F=M)	0.078	0.078	0.078	0.078
$maxF_{ABC}$ (maximum allowable = $0.75x F_{OFL}$)	0.0585	0.0585	0.0585	0.0585
F_{ABC}	0.0585	0.0585	0.0585	0.0585
OFL (t)	46,635	46,635	40,921	40,921
maxABC (t)	34,976	34,976	30,691	30,691
ABC (t)	34,976	34,976	30,691	30,691
Status	As determined <i>last year</i> for:		As determined <i>this year</i> for:	
	2012	2013	2013	2014
Overfishing	No	n/a	No	n/a

^aThe values for biomass, OFL, and ABC in these two columns are based on Rodgveller and Hulson 2013. They are an average of the last three trawl surveys that sampled down to 1,000 m. The current values (for 2015 and 2016) are from the random effects model fit to survey biomass by region and depth strata.

Bering Sea and Aleutian Islands Grenadiers

Quantity	As estimated or specified <i>last year for</i> ^a :		As estimated or recommended <i>this year</i> for:	
	2014	2015	2015	2016
<i>M</i> (natural mortality)	0.078	0.078	0.078	0.078
Specified/recommended Tier	5	5	5	5
Biomass (t)	1,152,284	1,152,284	1,286,734	1,286,734
F_{OFL} (F=M)	0.078	0.078	0.078	0.078
$maxF_{ABC}$ (maximum allowable = $0.75x F_{OFL}$)	0.0585	0.0585	0.0585	0.0585
F_{ABC}	0.0585	0.0585	0.0585	0.0585
OFL (t)	89,878	89,878	100,365	100,365
maxABC (t)	67,409	67,409	75,274	75,274
ABC (t)	67,409	67,409	75,274	75,274
Status	As determined <i>last year</i> for:		As determined <i>this year</i> for:	
	2012	2013	2013	2014
Overfishing	No	n/a	No	n/a

^aThe values for biomass, OFL, and ABC in these two columns are based on Rodgveller and Hulson 2013.

Not subject to over fishing: The recommended OFLs and ABCs in the above tables are much larger than the mean catches for grenadiers and also much larger than the catch in any single year, which indicates catches could increase without endangering the stocks. This is especially true for the EBS and AI, where the exploitation rate appears to be quite low.

Ecosystem Considerations

A determination of ecosystem considerations for grenadiers in Alaska is hampered by the extreme lack of biological and habitat information for these species and by limited knowledge in general on the deep slope environment inhabited by these fish.

Ecosystem Effects on the Stocks

Prey availability/abundance trends: The only food studies on grenadiers in the northeast Pacific have been on adults. One study of giant grenadier off the U.S. west coast concluded that the fish fed primarily off-bottom on bathy- and mesopelagic food items that included gonatid squids, viperfish, deep-sea smelts, and myctophids (Drazen et al. 2001). Smaller studies of giant grenadier food habits in Alaska showed generally similar results. In the Aleutian Islands, the diet comprised mostly squid and myctophids (Yang 2003), whereas in the Gulf of Alaska, squid and pasiphaeid shrimp predominated as prey (Yang et al. 2006). Research on these deep-sea prey organisms in Alaska has been virtually non-existent, so information on prey availability or possible variations in abundance of prey are unknown. Very few juvenile giant grenadier have ever been caught, so nothing is known about their food preferences.

In contrast to giant grenadier, a study of Pacific grenadier food habits off the U.S. west coast found a much higher consumption of benthic food items such as polychaetes, cumaceans, mysids, and juvenile Tanner crabs (*Chionoecetes* sp.), especially in smaller individuals (Drazen et al. 2001). Carrion also contributed to its diet, and larger individuals consumed some pelagic prey including squids, fish, and bathypelagic mysids.

Predator population trends: The only documented predators of giant grenadier are Pacific sleeper sharks (Orlov and Moiseev 1999) and Baird's beaked whales (Walker et al. 2002). According to Orlov's and Moiseev's study, giant grenadier was ranked third in relative importance as a food item in the diet of these sharks. Sperm whales are another potential predator, as they are known to dive to depths inhabited by giant grenadier on the slope and have been observed depredating on longline catches of giant grenadier⁶. Giant grenadier is a relatively large animal that is considered an apex predator in its environment on the deep slope (Drazen et al. 2001), so it may have relatively few predators as an adult. Predation on larval and juvenile giant grenadiers would likely have a much greater influence on the ultimate size of the adult population size, but information on predators of these earlier life stages is nil.

Changes in habitat quality: Little or no environmental information has been collected in Alaska for the deep slope habitat in which grenadiers live. This habitat is likely more stable oceanographically than shallower waters of the upper slope or continental shelf. Regime shifts on the continental shelf and slope in Alaska in recent decades have been well documented, but it is unknown if these shifts also extend to the deep slope. Regime shifts could have a pronounced effect on giant grenadier if their larvae or post-larvae inhabited upper portions of the water column. However, no larvae or post-larvae for this species have ever been collected in Alaska. The absence of larvae or post-larvae giant grenadier in larval surveys in Alaska, which have nearly all been conducted in upper parts of the water column, implies that larval giant grenadier may reside in deeper water, where they may be less affected by regime shifts since water temperatures in deep water tend to be more stable.

Fishery Effects on the Ecosystem

⁶ C. Lunsford, National Marine Fisheries Service, Alaska Fisheries Science Center, Auke Bay Laboratories, 17109 Point Lena Loop Rd., Juneau, AK 99801. Pers. comm. Oct 2012.

Because there has been virtually no directed fishing for grenadiers in Alaska, the reader is referred to the discussion on Fishery Effects in the sablefish SAFE report. The sablefish longline fishery is the main fishery that takes giant grenadier as bycatch, so the Fishery Effects section in the sablefish report is applicable to giant grenadier and is an indication of what the effects might be if a directed fishery for giant grenadier were to develop. It should be noted that because all grenadiers presently caught in the sablefish and Greenland turbot fisheries are discarded and do not survive, this constitutes a major input of dead organic material to the ecosystem that would not otherwise be there.

Data Gaps and Research Priorities

Research priorities

- 1) Because early life history information for giant grenadier is nil, studies are also needed to investigate where larvae and young juveniles reside.
- 2) Evaluation of the catchability of giant grenadier in the bottom trawl surveys, which would affect the accuracy of subsequent biomass estimates. Studies are needed on whether this fish is a completely benthic species or if individuals sometimes move off-bottom.
- 3) Validation of the AFSC REFM Division aging methodology for giant grenadier.
- 4) Further analysis and study of competition for hooks that may affect giant grenadier catch rates on the AFSC longline survey.
- 5) Continue a study to examine if the three different shapes of otoliths found in giant grenadier represent separate species or subpopulations. This is an ongoing cooperative project between the Marine Ecology and Stock Assessment program at Auke Bay Laboratories (ABL), REFM Age and Growth Lab, and the ABL genetics lab.

Current Research

Three otolith shapes were previously identified by Charles Hutchinson, AFSC Age and Growth Lab, in giant grenadier. A review of the literature revealed that this level of variability in otolith shape is extremely unusual for an individual fish species. These three otolith types may indicate that genetic subspecies or subpopulations exist for giant grenadier in Alaska that are not apparent based on the external morphology of the fish. Tagging studies are a traditional way to determine migration patterns and spatial stock structure for fish. However, these studies are not possible for giant grenadier because the fish do not survive the pressure difference when caught at depth and brought to the surface. Genetic and otolith microchemistry studies are an alternative means for determining patterns of stock structure.

In 2013, otolith and tissue samples were collected from giant grenadier in the eastern, central and western GOA and the EBS. Otoliths will be aged and photographed for morphometric measurements in 2015. Tissues are currently being analyzed by Scott Vulstek and Charles Guthrie, AFSC Auke Bay Laboratories genetics program. After genetic and morphometric data is collected, we will compare results from the two studies.

Literature cited

Ambrose, D. A. 1996. Macrouridae: grenadiers. *In*: Moser, H. G. (ed.). The early stages of fishes in the California Current region. CALCOFI Atlas No. 33. Calif. Coop. Oceanic Fish. Invest., La Jolla, CA, pp. 483-499.

- Andrews, A. H., G. M. Cailliet, and K. H. Coale. 1999. Age and growth of Pacific grenadier (*Coryphaenoides acrolepis*) with age estimate validation using an improved radiometric ageing technique. *Can. J. Fish. Aquat. Sci.* 56: 1339-1350.
- Burton, E. J. 1999. Radiometric age determination of the giant grenadier (*Albatrossia pectoralis*) using ^{210}Pb : ^{226}Ra disequilibria. Master's thesis, San Francisco State University, 91 p.
- Clausen, D. M. 2006. Grenadiers in the Gulf of Alaska, eastern Bering Sea, and Aleutian Islands. In Stock assessment and fishery evaluation report for the groundfish resources of the Gulf of Alaska and Bering Sea/Aleutian Islands regions, Appendix F, p. 563-600. North Pacific Fishery Management Council, 605 W 4th Ave., Suite 306, Anchorage AK 99501.
- Clausen, D. M. 2008. The giant grenadier in Alaska. *In* A. M. Orlov and T. Iwamoto (Editors), Grenadiers of the world oceans: biology, stock assessment, and fisheries, p. 413-450. *Amer. Fish. Soc. Symp.* 63. (Published by Amer. Fish. Soc., Bethesda, MD).
- Clausen, D. M., and C. J. Rodgveller. 2010. Assessment of grenadiers in the Gulf of Alaska, eastern Bering Sea, and Aleutian Islands. In Stock assessment and fishery evaluation report for the groundfish resources of the Gulf of Alaska and Bering Sea/Aleutian Islands regions, Appendix 1, p. 797-846. North Pacific Fishery Management Council, 605 W 4th Ave., Suite 306, Anchorage AK 99501.
- Rodgveller, C. J., D. M. Clausen, and P.-J. Hulson. 2012. Assessment of grenadiers in the Gulf of Alaska, eastern Bering Sea, and Aleutian Islands. In Stock assessment and fishery evaluation report for the groundfish resources of the Gulf of Alaska and Bering Sea/Aleutian Islands regions, Appendix 1, p. 797-846. North Pacific Fishery Management Council, 605 W 4th Ave., Suite 306, Anchorage AK 99501.
- Drazen, J. C., T. W. Buckley, and G. R. Hoff. 2001. The feeding habits of slope dwelling macrourid fishes in the eastern North Pacific. *Deep-Sea Res.* I 48:909-935.
- Endo, H., M. Yabe, and K. Amaoka. 1993. Occurrence of the macrourid alevins genera *Albatrossia* and *Coryphaenoides* in the northern North Pacific Ocean. *Japan J. Ichthyol.* 40(2): 219-226.
- Faunce, C., J. Cahalan, J. Gasper, T. A'mar, S. Lowe, F. Wallace, and R. Webster. 2014. Deployment performance review of the 2013 North Pacific Groundfish and Halibut Observer Program. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-AFSC-281, 74 p.
- Gaichas, S. 2002. Squid and other species in the Bering Sea and Aleutian Islands. *In*: Stock assessment and fishery evaluation report for the groundfish resources of the Bering Sea/Aleutian Islands Regions November 2002, p. 669-699. North Pacific Fishery Management Council, 605 W. 4th Ave., Suite 306, Anchorage AK 99501.
- Gaichas, S. 2003. Squid and other species in the Bering Sea and Aleutian Islands. *In*: Stock assessment and fishery evaluation report for the groundfish resources of the Bering Sea/Aleutian Islands region, November 2003, p. 777-808. North Pacific Fishery Management Council, 605 W. 4th Ave., Suite 306, Anchorage AK 99501.

- Hoening, J. M. 1983. Empirical use of longevity data to estimate mortality rates. *Fish. Bull.* 82(1): 898-902.
- Hoff, G. R., and L. L. Britt. 2009. Results of the 2008 eastern Bering Sea upper continental slope survey of groundfish and invertebrate resources. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-AFSC-197, 294 p.
- Hoff, G. R., and L. L. Britt. 2011. Results of the 2010 eastern Bering Sea upper continental slope survey of groundfish and invertebrate resources, 300 p.
- Iwamoto, T., and D. L. Stein. 1974. A systematic review of the rattail fishes (Macrouridae: Gadiformes) from Oregon and adjacent waters. *Occasional Papers of the California Academy of Sciences* No. 111, 79 p.
- Mecklenburg, C. W., T. A. Mecklenburg, and L. K. Thorsteinson. 2002. *Fishes of Alaska*. Amer. Fish. Soc., Bethesda, Maryland, 1,037 p.
- NMFS. 2014. Environmental Assessment, Regulatory Impact Review, and Initial Regulatory Flexibility Analysis for Amendment 100 to the Fishery Management Plan for Groundfish of the Bering Sea and Aleutian Islands, and Amendment 91 to the Fishery Management Plan for Groundfish of the Gulf of Alaska: Inclusion of Grenadiers (Family Macrouridae) in Groundfish Management. February 2014, U.S.DOC, NOAA, NMFS; Alaska Region, P.O. Box 21668, Juneau, AK 99802-1668.
- Novikov, N. P. 1970. Biology of *Chalinura pectoralis* in the North Pacific. *In*: P. A. Moiseev (Editor), Soviet fisheries investigations in the northeastern Pacific, Part V (In Russian). All-Union Scientific Research Institute of Marine Fisheries and Oceanography (VNIRO), Proceedings Vol. 70, and Pacific Scientific Research Institute of Fisheries and Oceanography (TINRO), Proceedings Vol. 72. (Translated by Israel Program for Scientific Translations, Jerusalem, 1972, p. 304-331).
- Orlov, A. M., and S. I. Moiseev. 1999. Some biological features of Pacific sleeper shark, *Somniosus pacificus* (Bigelow *et* Schroeder 1944) (Squalidae), in the northwestern Pacific Ocean. Polish Academy of Sciences, National Scientific Committee on Oceanic Research, Institute of Oceanography, University of Gdansk. *Oceanological Studies* FVIII No. 1-2: 3-16.
- Rodgveller, C. J., D. M. Clausen, J. J. Nagler, and C. Hutchinson. 2010. Reproductive characteristics and mortality of female giant grenadiers in the northern Pacific Ocean. *Mar. Coast. Fish.: Dynamics, Management, and Ecosystem Sci.* 2:73-82.
- Rodgveller, C. J., D. M. Clausen, P. Hulson. 2012. Assessment of the grenadier stock complex in the Gulf of Alaska, Eastern Bering Sea, and the Aleutian Islands. *In* Stock assessment and fishery evaluation report for the groundfish resources of the Gulf of Alaska for 2013. North Pacific Fishery Management Council, 605 W 4th Ave, Suite 306, Anchorage, AK 99501. <http://www.afsc.noaa.gov/REFM/Docs/2012/GOAgrenadier.pdf>
- Ronholt, L. L., K. Teshima, and D. W. Kessler. 1994. The groundfish resources of the Aleutian Islands region and southern Bering Sea 1980, 1983, and 1986. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-AFSC-31, 351 p.
- Tribuzio, C. A., C. Rodgveller, and P.-J. Huslon. 2014. GOA and BSAI shark assessments. September

Plan Team document.

http://www.afsc.noaa.gov/REFM/stocks/plan_team/2014_sharks_SeptPT.pdf

Tuponogov, V. N. 1997. Seasonal migrations of the grenadier *Coryphaenoides pectoralis* in the Sea of Okhotsk and contiguous waters. *Russ. J. Mar. Bio.* 23(6):314-321.

Walker, W. A., J. G. Mead, and R. L. Brownell, Jr. 2002. Diets of Baird's beaked whales, *Berardius bairdii*, in the southern Sea of Okhotsk and off the Pacific coast of Honshu, Japan. *Marine Mammal Science* 18(4): 902-919.

Yang, M-S. 2003. Food habits of the important groundfishes in the Aleutian Islands in 1994 and 1997. AFSC Processed Rep. 2003-07, 233 p. (Available from National Marine Fisheries Service, Alaska Fisheries Science Center, 7600 Sand Point Way NE, Seattle WA 98115).

Yang, M-S., K. Dodd, R. Hibpshman, and A. Whitehouse. 2006. Food habits of groundfishes in the Gulf of Alaska in 1999 and 2001. U.S. Department of Commerce, NOAA Technical Memorandum NMFS-AFSC-164.

Table 1.—Updated catch data (mt) for grenadiers, nearly all of which are thought to be giant grenadier, as of October 7, 2014 (NMFS Alaska Regional Office Catch Accounting System via the Alaska Fisheries Information Network (AKFIN) database, <http://www.akfin.org>). The estimates for 2003-2012 were computed by the NMFS Alaska Regional Office based on their Catch Accounting System, which replaced the “blend” system in 2003. Observer restructuring began in 2013 so the mean from 2003-2012 is presented for comparison to 2013 and 2014.

	Eastern Bering Sea	Aleutian Islands	Gulf of Alaska	Total
1997	2,964	2,887	12,029	17,881
1998	5,011	1,578	14,683	21,272
1999	4,505	2,883	11,388	18,776
2000	4,067	3,254	11,610	18,931
2001	2,294	1,460	9,685	13,439
2002	1,891	2,807	10,479	15,177
2003	2,641	3,036	10,843	16,520
2004	2,225	1,251	10,471	13,946
2005	2,633	1,795	6,606	11,034
2006	2,068	2,225	8,515	12,808
2007	1,645	1,817	9,629	13,091
2008	1,687	1,800	11,167	14,654
2009	2,983	3,681	6,696	13,360
2010	2,928	3,690	5,564	12,181
2011	4,333	2,578	7,437	14,349
2012	2,926	4,625	7,568	15,119
2013	1,629	2,535	11,339	15,504
2014	876	1,751	5,236	7,863
Mean 2003- 2012	2,607	2,650	8,450	13,706

Table 2.—Estimated catch (mt) of grenadiers (all species combined) in the eastern Bering Sea, Aleutian Islands, and Gulf of Alaska, by target species/species group, 2003-2011. Arrow = arrowtooth flounder; DW flat = deep-water flatfish; GT = Greenland turbot; halibut = Pacific halibut; Kam = Kamchatka flounder; cod = Pacific cod; rex = rex sole; sable = sablefish; other sp. = other species combined (including yellowfin sole, rock sole, shallow-water flatfish, “other flatfish”, flathead sole, and all other species). Source: Regional Office Catch Accounting System accessed through the Alaska Fisheries Information Network (AKFIN), October 7, 2014.

Year	Target species/species group									
	arrow	DW flat	GT	halibut	Kam	cod	rex	rockfish	sable	other
<u>Aleutian Islands</u>										
2003			113	1,376		46		6	1,494	0
2004			14	414		13		60	748	1
2005			161	617		2		21	979	16
2006	341		328	172		121		154	1,109	0
2007	108		342	69		41		21	1,161	76
2008	397		67	229		26		59	746	276
2009	1,377		414			12		152	1,642	84
2010	1,674		210	44		259		168	1,127	206
2011	51		83	13	723	18		292	1,292	105
2012	264			113	2,56	55		38	1,167	428
2013	278		44	239	406	3		215	1,139	212
2014	254			63	295	23		218	842	56
<u>Eastern Bering Sea</u>										
2003	38		1,452	355		240		9	370	164
2004	24		1,315	254		240		22	287	83
2005	11		1,977	143		333		32	108	31
2006	125		1,192	174		130		12	420	16
2007	2		1,073	89		179		17	215	70
2008	69		708	392		163		3	127	226
2009	243		1,823			212		6	692	8
2010	186		2,036	36		390		126	145	8
2011	807		1,799	7	241	1,13		17	316	5
2012	673		1,464	61	5	514		3	179	27
2013	272		533	321	12	274		47	166	4
2014	120		377	143	10	100		2	113	10

Table 2.—continued.

Year	Target species/species group									
	Arrow	DW flat	GT	Halibut	Kam Fl	Cod	Rex	Rockfish	Sable	Other
	<u>Gulf of Alaska</u>									
2003	27	474		710		5	325	613	8,464	223
2004	171	178		156		0	5	2,231	7,657	74
2005	103			488			4	212	5,743	56
2006	18			766		22	4	338	7,243	124
2007	90	20		530		79	5	198	8,702	5
2008	3			1,918		97	89	164	8,651	244
2009				1,430		79	102	227	4,816	43
2010	40	60		243		149	140	511	4,359	62
2011	114			172	723	69	229	529	6,208	116
2012	155			18	2,561	173	2	438	6,666	116
2013	161	1,246		338	406	169	4	1,008	8,327	87
2014	387			259	295	162	5	602	3,803	19

Table 3.—Biomass estimates for grenadier in the Gulf of Alaska using a random effects model. Estimates are for 1-1,000 m in all years. Left: estimates from using a random effects model. Center: estimates from NMFS trawl surveys that sampled down to 1,000 m. Right (status quo): biomass estimates used in SAFE reports for specifications of unofficial ABCs and OFLs since 2006. Biomass in 2006 was calculated as the average of the last two trawls surveys that sampled down to 1,000 m. Years after that the last three trawl surveys were averaged.

Year	Random Effects			Surveys down to 1,000 m			Status quo
	Biomass	U 95% CI	L 95% CI	Biomass	U 95% CI	L 95% CI	
1984	175,388	227,242	135,366	169,708	228,015	111,401	
1985	170,807	225,204	129,548				
1986	166,557	218,599	126,905				
1987	162,617	208,284	126,963	135,971	188,211	83,731	
1988	167,761	226,612	124,193				
1989	173,766	244,626	123,432				
1990	180,602	262,528	124,242				
1991	193,747	286,163	131,177				
1992	209,763	311,007	141,477				
1993	229,936	337,265	156,763				
1994	243,428	356,531	166,204				
1995	257,863	371,900	178,794				
1996	273,315	381,925	195,591				
1997	299,036	404,975	220,811				
1998	331,011	428,314	255,814				
1999	371,879	443,349	311,930	389,908	466,030	313,786	
2000	397,273	505,803	312,029				
2001	427,155	544,558	335,063				
2002	453,259	588,879	348,873				
2003	482,870	598,383	389,656				
2004	493,499	626,562	388,695				
2005	507,214	609,381	422,176	587,346	754,202	420,489	
2006	477,897	596,180	383,081				488,627
2007	463,065	568,252	377,349	487,987	629,173	346,802	488,414
2008	452,765	624,873	328,061				488,414
2009	455,461	749,608	276,738	718,320	1,484,296	0	597,884
2010	460,223	818,702	258,709				597,884
2011	473,685	898,965	249,595				597,884
2012	496,996	996,834	247,789				597,884
2013	524,624	1,099,917	250,229				597,884
2014	524,624	1,194,419	230,431				597,884


Table 4.—Biomass estimates (mt) and associated 95% confidence bounds (mt), variances, and coefficients of variation (cv) for giant grenadier in recent NMFS surveys in Alaska that sampled the upper continental slope. The Gulf of Alaska surveys included depths to 1,000 m, whereas the eastern Bering Sea slope surveys included depths to 1,200 m. Aleutian Islands biomass was estimated from trawl survey biomass estimates from 1-500 m and AFSC longline survey relative population weights from 200-1000m (see section titled “survey data”).

Region	Year	Biomass	95% Conf. bounds		Variance	cv (%)
			Lower	Upper		
Aleutian Islands	2000	560,200	290,106	830,294	18,989,690,223	24.6
Aleutian Islands	2002	570,239	295,651	844,828	19,626,898,754	24.6
Aleutian Islands	2004	575,396	297,497	853,295	20,103,051,745	24.6
Aleutian Islands	2006	721,531	373,113	1,069,950	31,600,203,801	24.6
Aleutian Islands	2008	365,940	189,744	542,136	8,081,265,243	24.6
Aleutian Islands	2010	688,251	356,757	1,019,745	28,604,825,470	24.6
Aleutian Islands	2012	478,991	246,867	711,115	14,025,825,748	24.7
Aleutian Islands	2014	733,177	379,987	1,086,366	32,471,605,527	24.6
Eastern Bering Sea	2002	426,397	344,922	507,871	1,659,519,194	9.6
Eastern Bering Sea	2004	666,508	527,524	805,491	4,829,084,657	10.4
Eastern Bering Sea	2008	449,777	353,902	545,652	2,298,003,647	10.7
Eastern Bering Sea	2010	660,528	521,035	800,021	4,864,588,623	10.6
Eastern Bering Sea	2012	550,366	433,097	667,635	3,437,997,235	10.6

Table 5.—Giant grenadier relative population weight, by region, in AFSC longline surveys in Alaska, 1990-2014. Dashes indicate years that the eastern Bering Sea or Aleutian Islands were not sampled by the survey. Gulf of Alaska values include data only for the upper continental slope at depths 201-1,000 m and do not include continental shelf gullies sampled in the surveys. Note: relative population weight, although an index of biomass (weight), is a unit-less value. NA indicates that length data is not available for calculations of RPWs. AFSC longline survey database query, October 2014.

Year	Eastern Bering Sea	Aleutian Islands	Gulf of Alaska
1992	-	-	686,827
1993	-	-	1,041,508
1994	-	-	1,018,292
1995	-	-	1,264,245
1996	-	2,281,815	1,121,058
1997	762,639	-	1,266,800
1998	-	2,268,918	1,066,477
1999	571,852	-	1,277,141
2000	-	3,039,523	1,143,980
2001	398,950	-	1,067,335
2002	-	3,093,994	904,922
2003	538,190	-	1,058,570
2004	-	3,121,973	801,271
2005	694,456	-	826,495
2006	-	3,914,871	857,510
2007	437,268	-	1,242,833
2008	-	1,985,511	919,083
2009	521,179	-	1,063,104
2010	-	3,734,301	1,236,692
2011	574,349	-	829,476
2012	-	3,230,202	911,728
2013	605,727	-	896,776
2014	-	3,978,057	848,321
mean	567,179	3,151,928	1,012,140

Figure 1.—Biomass estimates of giant grenadier from NMFS bottom trawl surveys and from a random effects model that utilizes trawl survey biomass estimates from all years (with 95% confidence intervals). The estimates of exploitable biomass used in previous assessments are also presented. From 1987-1996 this was the average of the last two trawl surveys that extended down to 1,000 m. From 1999-2013, it was the average of the biomass from the most recent three surveys that sampled down to 1,000 m.



Appendix A. Giant grenadier depth distribution by sex in fisheries and surveys.

Method and Results

Catch

Observed grenadier catch, not estimated total catch, was split by sex using sex ratios from observer specimen data, i.e., fish that had their lengths taken from 2003-2013 (Table 1; Figure 1). This timeframe was chosen because catch estimates are available for grenadier since 2003. Length frequencies by sex, stratum, and FMP area were converted to weights using area (BS, AI, and GOA) and sex specific growth curves from AFSC trawl surveys. The percent males by weight were used to split the observed catch for the Bering Sea (BS), Aleutian Islands (AI), and the Gulf of Alaska (GOA) by stratum (Table 2). The same percentages were used for splitting BS and AI observed catch (Table 1).

Total estimated grenadier catch from the Catch Accounting System (CAS) was split by sex using sex ratios of weight from observer specimen data (Table 3), as described above, except a single proportion was used for all depth strata combined because catch is not available by depth from CAS. The percent male was 13% in the BSAI and 15% in the GOA.

AFSC Longline Survey

The AFSC longline survey stations are spaced systematically (~20-30 km apart) along the slope from the eastern Gulf of Alaska west to the Aleutian Islands and north into the eastern Bering Sea. At each station, depths from ~150-1000 m are sampled. Giant grenadier are caught in great numbers throughout the survey range, primarily in depths from 400-1,000 m. The Aleutian Islands are sampled in even years, the Bering Sea in odd years, and the Gulf of Alaska is sampled annually. Because the area that is sampled by the longline cannot be defined, an index of abundance in weight is calculated, called relative populating weight (RPW), but is not a measure of absolute biomass. The index is used for tracking trends in abundance.

Giant grenadier length frequencies are available since 2006. Length frequencies by sex, stratum, and area (AI, BS, GOA) were converted to weights using area and sex specific growth curves from AFSC trawl surveys. The percent males, by weight, for each depth strata and area were calculated (Table 4) and used to split the RPWs by sex and stratum (Table 5, Figure 2).

AFSC Gulf of Alaska Trawl Survey

The AFSC GOA trawl survey samples the continental shelf and slope where stations are randomly chosen within depth strata. Only surveys that sampled down to 1,000 m were included in this analysis (1999, 2005, 2007, and 2009); surveys in 1984 and 1987 were not included because survey methodology changed in 1996. In other years, surveys sampled down to only 500 or 700 m and are not reflective of the extent of grenadier distribution by sex.

Giant grenadier population length frequencies are available split by sex for each depth stratum. We converted these population length frequencies to weight (biomass) using sex specific growth curves from GOA trawl survey data (Table 6). The biomass split by sex, year, and strata, as well as the percent of giant grenadier biomass that is male, is presented in Table 6. For comparison to the longline survey and observed catch, the average biomass by sex and strata are shown in Figure 3. Bering Sea trawl survey biomass estimates split by sex are not currently available and will be examined in the future.

Discussion

The observed catch is primarily between depths of 201-400 m; however, this is not where the bulk of giant grenadier biomass is found (e.g., see figures 1 and 2 for AFSC longline and trawl survey data). Observer length data shows that the percent of the catch that is male, by weight, increases with depth in the GOA, but there is the opposite trend in the BSAI. Although, the decreasing trend in male abundance is not dramatic in the BSAI and sample sizes for several depths strata are small. Due to small sampled sizes, the apparent trend in the BSAI may not be representative of the true distribution of giant grenadier. There is a much greater proportion of male grenadier in the catch data compared to the longline survey. This could be partially explained by the diverse gear types in the fisheries that incidentally catch grenadier; however, a large proportion of the observed grenadier catch is from longline fisheries. The difference between the proportion of males in longline survey and fishery could also be attributed to seasonal variation in depth distribution. The longline survey takes place only in the summer, whereas fisheries take place nearly year round. More time is required to explore distribution differences in the fishery by season and we plan to examine this in the future.

The trawl survey had a greater proportion of males than the longline survey and the proportion of males increased with depth in all surveys. The sex proportions in the trawl survey were more similar to the fishery than the longline survey when all depths are considered; however, in the 1-500 depth stratum the trawl survey had a very low percentage of males (2-5%), whereas the majority of the fishery data was from 201-400 m and the percentage male was larger than 5% (12-16%).

In all data sources, including surveys and fisheries, the large majority of catch is females. Also, overall the proportion of male grenadier, by weight, increases with deeper depths. Taken together, this information indicates that our surveys and fisheries may not completely cover the range of grenadier distribution. However, it also indicates that a disproportionate harvest of females is occurring, and should continue to be monitored.

Although a portion of the male population may reside in depths deeper than surveys and fisheries, it is possible that there is not a 1:1 ratio of males to females. We have not aged males and, therefore, it is not known if the natural mortality rate is different between sexes for grenadiers. Given the sexual dimorphism in growth and differences in distribution by sex, it could be postulated that other life-history parameters, like natural mortality, may also vary by sex. For example, in some flatfish species there is sexual dimorphism in natural mortality, where males have a much higher rate than females (e.g., arrowtooth flounder, 0.2 for females and 0.35 for males). If this is true for grenadier, the sex ratio may not be 1:1. The number of females could be larger than the number of males. Even in a deep-water AFSC longline survey (down to 1,600 m in the WGOA), on average 24% were male by number. Also, because females were much larger at depths >1,000 m than depths <1,000 m, the weight ratio would likely be much less than 24%.

Table 1. Sum of observed giant grenadier catch in mt for males (M) and females (F) from 2003-2013.

Depth strata	Aleutian Islands			Bering Sea			Gulf of Alaska		
	AI	AI F	AI M	BS	BS F	BS M	GOA	GOA F	GOA M
1-100	31	27	5	441	383	58	69	59	10
101-200	802	701	106	1,280	1,113	166	1,891	1,607	284
201-300	10,183	8,855	1,328	5,300	4,610	692	8,849	7,522	1,327
301-400	6,338	5,512	828	8,457	7,355	1,104	8,726	7,415	1,309
401-500	265	230	34	788	684	104	671	570	101
501-600	4	4	0	8	6	0	20	16	1
601-700	0	0	0	2	2	0	4	4	0
701-800	0	0	0	0	0	0	3	3	0
801-900	0	0	0	0	0	0	1	1	0

Table 2. Percent of male giant grenadier in observed catch from 2003-2013 in numbers and weight. Weights were calculated from length frequencies by depth, sex, and area using sex specific growth curves from the AFSC trawl survey. The total sample size (n) for length frequencies is presented for each sex.

Depth (m)	BSAI		GOA	
	% male (numbers, weight)	n	% male (numbers, weight)	n
1-100		0	17%, 15%	6
101-200	25%, 22%	296	9%, 7%	690
201-300	19%, 16%	4,535	14%, 12%	6,623
301-400	17%, 14%	8,013	20%, 16%	11,986
401-500	23%, 17%	719	28%, 21%	1,603
501-600	20%, 13%	155	37%, 24%	123
601-700	11%, 6%	22	56%, 54%	18
701-800		0	20%, 12%	5

Table 3. Total estimated grenadier catch from 2003-2013 split by sex (mt). Observed lengths were converted to weights using area and sex specific growth curves and the percent male was calculated using these weights. The average proportion of males and females by weight in the catch was used to split catch.

Year	BS total	BS male	BS female	AI total	AI Male	AI Female	GOA total	GOA male	GOA female
2003	2,869	373	2,439	3,558	463	3,024	12,253	1,838	10,415
2004	2,223	289	1,890	1,251	163	1,063	11,989	1,798	10,191
2005	2,633	342	2,238	1,795	233	1,526	7,251	1,088	6,163
2006	2,067	269	1,757	2,195	285	1,866	8,429	1,264	7,165
2007	1,631	212	1,386	1,544	201	1,312	9,119	1,368	7,751
2008	2,820	367	2,397	2,525	328	2,146	11,333	1,700	9,633
2009	2,902	377	2,467	3,739	486	3,178	6,326	949	5,377
2010	2,799	364	2,379	3,553	462	3,020	5,419	813	4,606
2011	4,221	549	3,588	2,596	337	2,207	8,216	1,232	6,984
2012	2,276	296	1,935	4,383	570	3,726	7,206	1,081	6,125
2013	1,482	193	1,260	2,367	308	2,012	10,525	1,579	8,946
average	2,538	330	2,158	2,682	349	2,280	8,915	1,337	7,578

Table 4. Percent of fish that were male caught during the AFSC longline survey 2006-2013 in numbers and weight. Weights were calculated from length frequencies by depth, sex, and area using sex specific growth curves from the AFSC trawl survey. The total sample size (n) for length frequencies is presented for each sex.

Depth (m)	Aleutian Islands		Bering Sea		Gulf of Alaska	
	% male (numbers, weight)	n	% male (numbers, weight)	n	% male (numbers, weight)	n
101-200	0%, 0%	20	0%, 0%	9	0%, 0%	11
201-300	0%, 0%	312	0%, 0%	582	0.5%, 0.8%	2,098
301-400	0%, 0%	1,280	0%, 0%	1,559	1%, 0.5%	9,947
401-600	2%, 2%	2,912	0%, 0%	2,949	2%, 1%	19,527
601-800	6%, 5%	2,533	2%, 1%	3,038	5%, 3%	17,378
801-1000	20%, 15%	777	6%, 4%	1,015	9%, 6%	8,603
Total	5%, 4%	7,834	2%, 1%	9,255	4%, 3%	58,828

Table 5. Average AFCS longline survey Relative Population Weights split by sex and strata from 2006-2013.

Strata (m)	Aleutian Islands		Bering Sea		Gulf of Alaska	
	AI M	AI F	BS M	BS F	GOA M	GOA F
201-300	0	16,322	7	9,263	152	17,772
301-400	189	93,257	0	39,385	477	95,590
401-600	6,707	299,368	285	121,839	3,613	279,659
601-800	17,890	348,645	2,854	232,902	10,020	283,846
801-1000	37,164	218,898	6,692	148,634	17,131	271,927

Table 6. AFCS Gulf of Alaska trawl survey biomass estimates from recent years when the survey sampled down to 1,000 m (1999, 2005, 2007, 2009).

Depth strata (m)	Year	GOA M	GOA F	% male
1-500	1999	2,183	126,234	2%
1-500	2005	10,698	226,337	5%
1-500	2007	3,163	103,382	3%
1-500	2009	1,510	89,961	2%
501-700	1999	15,336	136,471	10%
501-700	2005	25,470	221,437	10%
501-700	2007	16,467	218,538	7%
501-700	2009	29,116	142,184	17%
701-1000	1999	28,466	81,219	26%
701-1000	2005	28,522	74,882	28%
701-1000	2007	46,574	99,862	32%
701-1000	2009	83,034	372,514	18%

Figure 1. Summed observed grenadier catch from 2003-2013, not total estimated catch, split by sex and depth strata.




Figure 2. Average AFCS longline survey giant grenadier Relative Population Weights from 2006-2013 split by sex and depth stratum.





Figure 3. Average GOA AFCS trawl survey giant grenadier biomass estimates in 1999, 2005, 2007, and 2009 split by sex and strata (recent years when the survey sampled down to 1,000 m).



Appendix B

Tabbe 1B-1.—Research catch (mt) of grenadier (giant, popeye, and pacific grenadier, but primarily giant grenadier) in AFSC trawl and longline (LL) surveys and the International Pacific Halibut Commission (IPHC) longline survey. Only numbers are available from the IPHC survey through 2009; 2010 and 2011 catch in weight is available. 0s indicate that there was catch but it is <1 mt.

Year	BSAI					GOA					Total
	IPHC #s	IPHC wt	AFSC Trawl	AFSC LL	Total BSAI	IPHC #s	IPH C wt	AFSC Trawl	AFSC LL	Total GOA	
1979			33		33			0		0	33
1980			85		85			1		1	86
1981			66		66			3		3	69
1982			124		124			0		0	125
1983			136		136			0		0	136
1984								59		59	59
1985			165		165			9		9	174
1986			90		90			0		0	90
1987			0		0			42		42	42
1988			30		30						30
1989											
1990								3	128	131	131
1991			10		10				113	113	123
1992									117	117	117
1993								6	135	141	141
1994			6		6				134	134	140
1995									191	191	191
1996				38	38			8	173	181	219
1997	1,184		9	78	87	258			169	169	256
1998	556			59	59	681		12	141	153	212
1999	165		0	57	57	660		47	157	204	261
2000	774		118	88	206	621			160	160	366
2001	1,313			43	43	287		11	161	173	215
2002	987		23	81	104	942			129	129	233
2003	1,792		91	50	141	1,344		27	151	178	320
2004	2,111		196	78	274	1,110			109	109	383
2005	1,404			71	71	1,266		49	120	169	240
2006	941		20	76	96	919			112	112	208
2007	1,224			77	77	849		44	166	209	286
2008	1,331		123	47	170	755			120	120	290
2009	2,710			86	86	785		39	154	193	279
2010	2,451	9	156	66	231	1,265	6		164	170	401
2011	1,808	7		75	82	751	2	20	124	145	227
2012		5	135	43	177		2		132	132	310
*2013		5		83	88		2		132	134	222
*2014				73	73				127	127	200

*GOA trawl survey data is not available through the AKRO for grenadier for 2013 (accessed through AKFIN, October 2014). Only AFSC longline survey data is available for 2014.

(This page intentionally left blank)