

3. Assessment of the sablefish stock in Alaska

by

Dana H. Hanselman, Chris R. Lunsford, and Cara J. Rodgveller

Executive Summary

Summary of changes in assessment inputs

Relative to last year's assessment, we made the following substantive changes in the current assessment.

Changes in the input data: New data included in the assessment model were relative abundance and length data from the 2015 longline survey, relative abundance and length data from the 2014 longline fishery, length data from the 2014 trawl fisheries, age data from the 2014 longline survey and 2014 fixed gear fishery, the 2015 Gulf of Alaska trawl survey abundance and length compositions, updated catch for 2014, and projected 2015- 2017 catches.

Changes in the assessment methodology: There are no model changes.

Summary of results

Quantity/Status	As estimated or specified <i>last</i> year for:		As estimated or recommended <i>this</i> year for:	
	2015	2016	2016*	2017*
M (natural mortality rate)	0.10	0.10	0.10	0.10
Tier	3b	3b	3b	3b
Projected total (age 2+) biomass (t)	219,997	227,042	204,796	214,552
Projected female spawning biomass (t)	91,183	88,345	86,471	81,986
$B_{100\%}$	262,269	262,269	257,018	257,018
$B_{40\%}$	104,908	104,908	102,807	102,807
$B_{35\%}$	91,794	91,794	89,956	89,956
F_{OFL}	0.098	0.091	0.093	0.086
$maxF_{ABC}$	0.082	0.078	0.078	0.073
F_{ABC}	0.082	0.078	0.078	0.073
OFL (t)	16,128	14,658	13,397	12,747
max ABC (t)	13,657	12,406	11,795	10,782
ABC (t)	13,657	12,406	11,795	10,782
Status	As determined <i>last</i> year for:		As determined <i>this</i> year for:	
	2013	2014	2014	2015
Overfishing	No	n/a	No	n/a
Overfished	n/a	No	n/a	No
Approaching overfished	n/a	No	n/a	No

* Projections are based on estimated catches of 9,781 t and 8,715 t used in place of maximum permissible ABC for 2016 and 2017. This was done in response to management requests for a more accurate two-year projection.

Assessment results

The longline survey abundance index decreased 21% from 2014 to 2015 following a 15% increase from 2013 to 2014 and is at the lowest point of the time series. The fishery abundance index increased 6% from 2013 to 2014 (the 2015 data are not available yet). The Gulf of Alaska trawl survey index was at its lowest point in 2013 but increased 12% in 2015. Spawning biomass is projected to decrease from 2016 to 2019, and then stabilize.

Sablefish are managed under Tier 3 of NPFMC harvest rules. Reference points are calculated using recruitments from 1977-2012. The updated point estimates of $B_{40\%}$, $F_{40\%}$, and $F_{35\%}$ from this assessment are 102,807 t (combined across the EBS, AI, and GOA), 0.094, and 0.112, respectively. Projected female spawning biomass (combined areas) for 2016 is 86,471 t (84% of $B_{40\%}$), placing sablefish in sub-tier “b” of Tier 3. The maximum permissible value of F_{ABC} under Tier 3b is 0.078, which translates into a 2016 ABC (combined areas) of 11,795 t. The OFL fishing mortality rate is 0.093 which translates into a 2016 OFL (combined areas) of 13,397 t. If the stock were in Tier 3a (above the $B_{40\%}$ reference point), the 2016 ABC would be 14,164 t. Model projections indicate that this stock is not subject to overfishing, overfished, nor approaching an overfished condition.

We recommend a 2016 ABC of 11,795 t. The maximum permissible ABC for 2016 based on Tier 3b of the harvest control rule, uses an adjusted $F_{40\%}$ which yields 11,795 t. The maximum permissible ABC for 2016 is 14% lower than the 2015 ABC of 13,657 t. The 2014 assessment projected a 10% decrease in ABC for 2016 from 2015. This slightly larger decrease is supported by a new low in the domestic longline survey index time series that offset the small increases in the fishery abundance index seen in 2014 and the Gulf of Alaska trawl survey index in 2015. The fishery abundance index has been trending down since 2007. The 2014 IPHC GOA sablefish index was not used in the model, but was similar and trending low in 2013 and 2014. The 2008 year class showed potential to be large in previous assessments based on patterns in the age and length compositions. However the estimate in this year’s assessment is only just above average because the recent large overall decrease in the longline survey and trawl indices have lowered the overall scale of the population. Spawning biomass is projected to decline through 2018, and then is expected to increase assuming average recruitment is achieved in the future. ABCs are projected to decrease in 2017 to 10,782 t and 10,869 t in 2018 (see Table 3.18).

Projected 2016 spawning biomass is 34% of unfished spawning biomass. Spawning biomass had increased from a low of 33% of unfished biomass in 2002 to 42% in 2008 and has now declined back to 34% of unfished biomass projected for 2016. The 1997 year class has been an important contributor to the population; however, it has been reduced and is predicted to comprise less than 6% of the 2016 spawning biomass. The last two above-average year classes, 2000 and 2008, each comprise 15% of the projected 2016 spawning biomass. The 2008 year class will be about 75% mature in 2016.

Apportionment

In December 1999, the Council apportioned the 2000 ABC and OFL based on a 5-year exponential weighting of the survey and fishery abundance indices. We have used the same algorithm to apportion the ABC and OFL since 2000. Following the standard apportionment scheme, we have observed that the objective to reduce variability in apportionment was not being achieved. Since 2007, the mean change in apportionment by area has increased annually (Figure 3.36A). While some of these changes may actually reflect interannual changes in regional abundance, they most likely reflect the high movement rates of the population and the high variability of our estimates of abundance in the Bering Sea and Aleutian Islands. For example, the apportionment for the Bering Sea has varied drastically since 2007, attributable to high variability in both survey abundance and fishery CPUE estimates in the Bering Sea (Figure 3.36B). These large annual changes in apportionment result in increased variability of ABCs by area, including areas other than the Bering Sea (Figure 3.36C). Because of the high variability in apportionment seen in recent years, we do not believe the standard method is meeting the goal of reducing the magnitude of interannual changes in the apportionment. Because of these reasons, we recommended fixing the apportionment at the proportions from the 2013 assessment, until the apportionment scheme is thoroughly reevaluated and reviewed. A Ph.D. student with the University of Alaska-Fairbanks began a project in 2013 with the objectives of re-examining the apportionment strategy and conducting a management strategy evaluation. A spatial sablefish model has been developed, but the management strategy evaluation is in early stages of development. Meanwhile, it seems imprudent to move to an interim apportionment or return to the former scheme until more satisfactory methods have been identified and evaluated. **Therefore, for 2016, we recommend continuing with the apportionment fixed at the proportions used in 2015.**

Area	2015 ABC	Standard apportionment for 2016 ABC	Recommended fixed apportionment for 2016 ABC*	Difference from 2015
Total	13,657	11,795	11,795	-13.6%
Bering Sea	1,333	1,816	1,151	-13.6%
Aleutians	1,802	1,627	1,557	-13.6%
Gulf of Alaska (subtotal)	10,522	8,352	9,087	-13.6%
Western	1,473	1,136	1,272	-13.6%
Central	4,658	3,451	4,023	-13.6%
W. Yakutat**	1,567	1,374	1,353	-13.6%
E. Yak. / Southeast**	2,823	2,391	2,438	-13.6%

* Fixed at the 2013 assessment apportionment proportions (Hanselman et al. 2012). ** Before 95:5 hook and line: trawl split shown below.

Adjusted for 95:5 hook-and-line: trawl split in EGOA	Year	W. Yakutat	E. Yakutat/Southeast
	2016	1,475 t	2,316 t
	2017	1,348 t	2,118 t

Plan team summaries

Area	Year	Biomass (4+)	OFL	ABC	TAC	Catch
GOA	2014	149,000	12,500	10,572	10,572	10,343
	2015	130,000	12,425	10,522	10,522	9,525
	2016	122,000	10,326	9,087		
	2017	123,000	9,825	8,307		
BS	2014	21,000	1,584	1,339	1,339	315
	2015	34,000	1,574	1,333	1,333	197
	2016	25,000	1,304	1,151		
	2017	26,000	1,241	1,052		
AI	2014	28,000	2,141	1,811	1,811	818
	2015	24,000	2,128	1,802	1,802	372
	2016	23,000	1,766	1,557		
	2017	23,000	1,681	1,423		

Year	2015				2016		2017	
Region	OFL	ABC	TAC	Catch*	OFL	ABC	OFL	ABC
BS	1,574	1,333	1,333	197	1,304	1,151	1,241	1,052
AI	2,128	1,802	1,802	372	1,766	1,557	1,681	1,423
GOA	12,425	10,522	10,522	9,525	10,326	9,087	9,825	8,307
WGOA	--	1,473	1,473	867	--	1,272	--	1,163
CGOA	--	4,658	4,658	4,176	--	4,023	--	3,678
**WYAK	--	1,708	1,708	1,794	--	1,475	--	1,348
**EY/SEO	--	2,682	2,682	2,688	--	2,316	--	2,118
Total	16,128	13,657	13,657	10,094	13,397	11,795	12,747	10,782

* As of October 29, 2015 Alaska Fisheries Information Network, (www.akfin.org). ** After 95:5 trawl split shown above.

Responses to SSC and Plan Team Comments on Assessments in General

“The SSC requests that stock assessment authors utilize the following model naming conventions in SAFE chapters:

Model 0: last years’ model with no new data,

Model 1: last years’ model with updated data, and

Model numbers higher than 1 are for proposed new models.”: SSC, December 2014

“For this year’s final assessments, the Teams recommend that each author of an age-structured assessment use one of the following model naming conventions (“TPA” represents the alternative described in the Team procedures document)...”: Joint Plan Team, September, 2015

“Of the options presented in the Joint Plan Teams minutes, the SSC agrees that that Option 4 has several advantages and recommends that this Option be advanced next year.”: SSC, October 2015

For this assessment, we will use the simplified convention suggested in the December SSC minutes and will investigate further detailed naming for the next assessment cycle in 2016.

The SSC also requests that stock assessment authors utilize the random effects model for area apportionment of ABCs”: SSC, December 2014

“The Teams recommend that the random effects survey smoothing model be used as a default for determining current survey biomass and apportionment among areas.”: Joint Plan Teams, September 2015

The sablefish model has used a 5 year exponential smoothing model of fishery and survey CPUE developed at the Council level that was based on the univariate Kalman filter model. This is similar to the random effects apportionment model, which smooths biomass by balancing process and measurement error. We will examine the random effects apportionment model in the future as different apportionment options are being examined for sablefish.

Responses to SSC and Plan Team Comments Specific to this Assessment

There were no recommendations specific to sablefish in 2014 or 2015.

Introduction

Distribution

Sablefish (*Anoplopoma fimbria*) inhabit the northeastern Pacific Ocean from northern Mexico to the Gulf of Alaska (GOA), westward to the Aleutian Islands (AI), and into the Bering Sea (BS) (Wolotira et al. 1993). Adult sablefish occur along the continental slope, shelf gullies, and in deep fjords, generally at depths greater than 200 m. Sablefish observed from a manned submersible were found on or within 1 m of the bottom (Krieger 1997). In contrast to the adult distribution, juvenile sablefish spend their first two to three years on the continental shelf of the GOA, and occasionally on the shelf of the southeast BS. The BS shelf is utilized significantly in some years and seldom used during other years (Shotwell et al. 2012).

Early life history

Spawning is pelagic at depths of 300-500 m near the edges of the continental slope (Mason et al. 1983, McFarlane and Nagata 1988), with eggs developing at depth and larvae developing near the surface as far offshore as 180 miles (Wing 1997). Along the Canadian coast (Mason et al. 1983) and off Southeast Alaska (Jennifer Stahl, February, 2010, ADF&G, pers. comm.) sablefish spawn from January-April with a peak in February. In a survey near Kodiak Island in December, 2011 that targeted sablefish preparing to spawn, spawning appeared to be imminent, but spent fish were not found. It is likely that they would spawn in January or February (Katy Echave, October 2012, AFSC, pers. comm.). Farther down the coast off of central California sablefish spawn earlier, from October-February (Hunter et al. 1989). An analysis of larval otoliths showed that spawning in the Gulf of Alaska may be a month later than southern sablefish (Sigler et al. 2001). Sablefish in spawning condition were also noted as far west as Kamchatka in November and December (Orlov and Biryukov 2005). Larval sablefish sampled by neuston net in the eastern Bering Sea fed primarily on copepod nauplii and adult copepods (Grover and Olla 1990). In gill nets set at night for several years on the AFSC longline survey, most young-of-the-year sablefish were caught in the central and eastern GOA (Sigler et al. 2001). Near the end of the first summer, pelagic juveniles less than 20 cm move inshore and spend the winter and following summer in inshore waters where they exhibit rapid growth, reaching 30-40 cm by the end of their second summer (Rutecki and Varosi 1997). Gao et al. (2004) studied stable isotopes in otoliths of juvenile sablefish from Oregon and Washington and found that as the fish increased in size they shifted from midwater prey to more benthic prey. In nearshore southeast Alaska, juvenile sablefish (20-45 cm) diets included fish such as Pacific herring and smelts and invertebrates such as krill, amphipods and polychaete worms (Coutré et al. 2015). In late summer, juvenile sablefish also consumed post-spawning pacific salmon carcass remnants in high volume revealing opportunistic scavenging (Coutré et al. 2015). After their second summer, they begin moving offshore to deeper water, typically reaching their adult habitat, the upper continental slope at 4 to 5 years. This corresponds to the age range when sablefish start becoming reproductively viable (Mason et al. 1983).

Movement

A movement model for Alaskan sablefish was developed for Alaskan sablefish by Heifetz and Fujioka (1991) based on 10 years of tagging data. The model has been updated by incorporating data from 1979-2009 in an AD Model Builder program, with time-varying reporting rates, and tag recovery data from ADF&G for State inside waters (Southern Southeast Inside and Northern Southeast Inside). In addition, the study estimated mortality rates from the tagging data (Hanselman et al. 2015). Annual movement probabilities were high, ranging from 10-88% depending on area of occupancy at each time step, and size group. Overall, movement probabilities were very different between areas of occupancy and moderately different between size groups. Estimated annual movement of small sablefish from the central Gulf of Alaska had the reverse pattern of a previous study, with 29% moving westward and 39% moving

eastward. Movement probabilities also varied annually with decreasing movement until the late 1990s and increasing movement until 2009. Year specific magnitude in movement probability of large fish was highly negatively correlated with female spawning biomass estimates from the federal stock assessment (i.e., when spawning biomass is high, they move less). Average mortality estimates from time at liberty were similar to the stock assessment.

Stock structure

Sablefish have traditionally been thought to form two populations based on differences in growth rate, size at maturity, and tagging studies (McDevitt 1990, Saunders et al. 1996, Kimura et al. 1998). The northern population inhabits Alaska and northern British Columbia waters and the southern population inhabits southern British Columbia, Washington, Oregon, and California waters, with mixing of the two populations occurring off southwest Vancouver Island and northwest Washington. Significant stock structure among the federal Alaska population is unlikely given extremely high movement rates throughout their lives (Hanselman et al. 2015, Heifetz and Fujioka 1991, Maloney and Heifetz 1997, Kimura et al. 1998).

Fishery

Early U.S. fishery, 1957 and earlier

Sablefish have been exploited since the end of the 19th century by U.S. and Canadian fishermen. The North American fishery on sablefish developed as a secondary activity of the halibut fishery of the United States and Canada. Initial fishing grounds were off Washington and British Columbia and then spread to Oregon, California, and Alaska during the 1920's. Until 1957, the sablefish fishery was exclusively a U.S. and Canadian fishery, ranging from off northern California northward to Kodiak Island in the GOA; catches were relatively small, averaging 1,666 t from 1930 to 1957, and generally limited to areas near fishing ports (Low et al. 1976).

Foreign fisheries, 1958 to 1987

Japanese longliners began operations in the eastern BS in 1958. The fishery expanded rapidly in this area and catches peaked at 25,989 t in 1962 (Table 3.1, Figures 3.1, 3.2). As the fishing grounds in the eastern Bering were preempted by expanding Japanese trawl fisheries, the Japanese longline fleet expanded to the AI region and the GOA. In the GOA, sablefish catches increased rapidly as the Japanese longline fishery expanded, peaking at 36,776 t overall in 1972. Catches in the AI region remained at low levels with Japan harvesting the largest portion of the sablefish catch. Most sablefish harvests were taken from the eastern Bering Sea until 1968, and then from the GOA until 1977. Heavy fishing by foreign vessels during the 1970's led to a substantial population decline and fishery regulations in Alaska, which sharply reduced catches. Catch in the late 1970's was restricted to about one-fifth of the peak catch in 1972, due to the passage of the Fishery Conservation and Management Act (FCMA).

Japanese trawlers caught sablefish mostly as bycatch in fisheries targeting other species. In the BS, the trawlers were mainly targeting rockfishes, Greenland turbot, and Pacific cod, and only a few vessels targeted sablefish. In the GOA, sablefish were mainly caught as bycatch in the directed Pacific Ocean perch fishery until 1972, when some vessels started targeting sablefish in 1972 (Sasaki 1985).

Other foreign nations besides Japan also caught sablefish. Substantial Soviet Union catches were reported from 1967-73 in the BS (McDevitt 1986). Substantial Korean catches were reported from 1974-1983 scattered throughout Alaska. Other countries reporting minor sablefish catches were Republic of Poland, Taiwan, Mexico, Bulgaria, Federal Republic of Germany, and Portugal. The Soviet gear was factory-type stern trawl and the Korean gears were longlines and pots (Low et al. 1976).

Recent U.S. fishery, 1977 to present

The U.S. longline fishery began expanding in 1982 in the GOA, and by 1988, the U.S. harvested all sablefish taken in Alaska, except minor joint venture catches. Following domestication of the fishery, the previously year-round season in the GOA began to shorten in 1984 from 12 months in 1983 to 10 days in 1994, warranting the label “derby” fishery.

In 1995, Individual Fishery Quotas (IFQ) were implemented for hook-and-line vessels along with an 8-month season. The IFQ Program is a catch share fishery that issued quota shares to individuals based on sablefish and halibut landings made from 1988-1990. Since the implementation of IFQ's, the number of longline vessels with sablefish IFQ harvests has experienced a substantial anticipated decline from 616 in 1995 to 362 in 2011 (NOAA 2012). This decrease was expected as shareholders have consolidated their holdings and fish them off fewer vessels to reduce costs (Fina 2011). The sablefish fishery has historically been a small boat fishery; the median vessel length in the 2011 fishery was 56ft. In recent years, approximately 30% of vessels eligible to fish in the IFQ fishery participate in both the halibut and sablefish fisheries and approximately 40% of vessels fish in more than one management area. The season dates have varied by several weeks since 1995, but the monthly pattern has been from March to November with the majority of landings occurring in May - June. The number of landings fluctuates with quota size, but in 2011 there were 1,726 landings recorded in the Alaska fishery (NOAA 2012).

Pot fishing in the IFQ fishery is not allowed in the GOA but is legal in the BSAI regions. In 2000, the pot fishery accounted for less than ten percent of the fixed gear sablefish catch in these areas but effort has increased substantially in response to killer whale depredation. Pots are longlined with approximately 40-135 pots per set. Since 2004, pot gear has accounted for over 50% of the BS fixed gear IFQ catch and up to 34% of the fixed gear catch in the AI.

Sablefish also are caught incidentally during directed trawl fisheries for other species groups such as rockfish and deepwater flatfish. Allocation of the TAC by gear group varies by management region and influences the amount of catch in each region (Table 3.1, Figures 3.1, 3.2). Five State of Alaska fisheries land sablefish outside the IFQ program; the major State fisheries occur in the Prince William Sound, Chatham Strait, and Clarence Strait and the minor fisheries in the northern GOA and AI. The minor state fisheries were established by the State of Alaska in 1995, the same time as the Federal Government established the IFQ fishery, primarily to provide open-access fisheries to fishermen who could not participate in the IFQ fishery.

IFQ management has increased fishery catch rates and decreased the harvest of immature fish (Sigler and Lunsford 2001). Catching efficiency (the average catch rate per hook for sablefish) increased 1.8 times with the change from an open-access to an IFQ fishery. The change to IFQ also decreased harvest and discard of immature fish which improved the chance that these fish will reproduce at least once. Thus, the stock can provide a greater yield under IFQ at the same target fishing rate because of the selection of older fish (Sigler and Lunsford 2001).

Longline gear in Alaska is fished on-bottom. Since the inception of the IFQ system, average set length in the directed fishery for sablefish has been near 9 km and average hook spacing near 1.2 m. The gear is baited by hand or by machine, with smaller boats generally baiting by hand and larger boats generally baiting by machine. Circle hooks are usually used, except for modified J-hooks on some boats with machine baiters. The gear usually is deployed from the vessel stern with the vessel traveling at 5-7 knots. Some vessels attach weights to the longline, especially on rough or steep bottom, so that the longline stays in place on bottom.

Management measures/units

A summary of historical catch and management measures pertinent to sablefish in Alaska are shown in Table 3.7. Influential management actions regarding sablefish include:

Management units

Sablefish are assessed as a single population in Federal waters off Alaska because of their high movement rates. Sablefish are managed by discrete regions to distribute exploitation throughout their wide geographical range. There are four management areas in the GOA: Western, Central, West Yakutat, and East Yakutat/Southeast Outside; and two management areas in the Bering Sea/Aleutian Islands (BSAI): the BS and the AI regions. Amendment 8 to the GOA Fishery Management Plan established the West and East Yakutat management areas for sablefish, effective 1980.

Quota allocation

Amendment 14 to the GOA Fishery Management Plan allocated the sablefish quota by gear type: 80% to fixed gear (including pots) and 20% to trawl in the Western and Central GOA, and 95% to fixed gear and 5% to trawl in the Eastern GOA, effective 1985. Amendment 15 to the BS/AI Fishery Management Plan, allocated the sablefish quota by gear type, 50% to fixed gear and 50% to trawl in the eastern BS, and 75% to fixed gear and 25% to trawl gear in the Aleutians, effective 1990.

IFQ management

Amendment 20 to the GOA Fishery Management Plan and 15 to the BS/AI Fishery Management Plan established IFQ management for sablefish beginning in 1995. These amendments also allocated 20% of the fixed gear allocation of sablefish to a CDQ reserve for the BS and AI.

Maximum retainable allowances

Maximum retainable allowances for sablefish as the “incidental catch species” were revised in the GOA by a regulatory amendment, effective April, 1997. The percentage depends on the basis species: 1% for pollock, Pacific cod, Atka mackerel, “other species”, and aggregated amount of non-groundfish species. Fisheries targeting deep flatfish, rex sole, flathead sole, shallow flatfish, Pacific ocean perch, northern rockfish, dusky rockfish, and demersal shelf rockfish in the Southeast Outside district, and thornyheads are allowed 7%. The MRA for arrowtooth flounder changed effective 2009 in the GOA, to 1% for sablefish as the basis species.

Allowable gear

Amendment 14 to the GOA Fishery Management Plan banned the use of pots for fishing for sablefish in the GOA, effective 18 November 1985, starting in the Eastern area in 1986, in the Central area in 1987, and in the Western area in 1989. An earlier regulatory amendment was approved in 1985 for 3 months (27 March - 25 June 1985) until Amendment 14 was effective. A later regulatory amendment in 1992 prohibited longline pot gear in the BS (57 FR 37906). The prohibition on sablefish longline pot gear use was removed for the BS, except from 1 to 30 June to prevent gear conflicts with trawlers during that month, effective 12 September 1996. Sablefish longline pot gear is allowed in the AI. In April of 2015 the NPFMC passed a motion to again allow for sablefish pot fishing in the GOA in response to increased sperm whale depredation. The final motion was passed and the new regulations are expected in early 2016. We will carefully monitor the development of this gear type in the Gulf of Alaska.

Catch

Annual catches in Alaska averaged about 1,700 t from 1930 to 1957 and exploitation rates remained low until Japanese vessels began fishing for sablefish in the BS in 1958 and the GOA in 1963. Catches rapidly escalated during the mid-1960s. Annual catches in Alaska reached peaks in 1962, 1972, and 1988 (Table 3.1, Figure 3.2). The 1972 catch was the all-time high, at 53,080 t, and the 1962 and 1988 catches were 50% and 72% of the 1972 catch. Evidence of declining stock abundance and passage of the MSFCMA led to significant fishery restrictions from 1978 to 1985, and total catches were reduced substantially.

Exceptional recruitment fueled increased abundance and increased catches during the late 1980's, which coincided with the domestic fishery expansion. Catches declined during the 1990's, increased in the early

2000s, and have since declined to near 12,000 t (Figure 3.1). TACs in the GOA are nearly fully utilized, while TACs in the BS and AI are rarely fully utilized.

Bycatch and discards

Sablefish discards by target fisheries are available for hook-and-line gear and other gear combined (Table 3.3). From 1994 to 2004 discards averaged 1,357 t for the GOA and BSAI combined (Hanselman et al. 2008). Since then, discards have been lower, averaging 608 t between 2007 and 2014. Discard rates are generally higher in the GOA than in the BSAI (Table 3.3).

Table 3.4 shows the average bycatch of Fishery Management Plans' (FMP) groundfish species in the sablefish target fishery from 2009-2015. The largest bycatch group is GOA thornyhead rockfish (575 t/year, 174 t discarded). Sharks and skates are also taken in substantial numbers and are mostly discarded.

Giant grenadiers, a non-target species that is soon entering both FMPs as an Ecosystem Component, make up the bulk of the nontarget species bycatch, with 2013 the highest in the last five years at 7,929 t (Table 3.5). Other nontarget taxa that have catches over one ton per year are corals, snails, sponges, sea stars, and miscellaneous fishes and crabs.

Prohibited species catches (PSC) in the targeted sablefish fisheries are dominated by halibut (334 t/year on average) and golden king crab (47,000 individuals/year on average) (Table 3.6). Crab catches are highly variable from year to year, probably as a result of relatively low observer sampling effort in sablefish fisheries.

Data

The following table summarizes the data used for this assessment:

Source	Data	Years
Fixed gear fisheries	Catch	1960-2015
Trawl fisheries	Catch	1960-2015
Japanese longline fishery	Catch-per-unit-effort (CPUE)	1964-1981
U.S. fixed gear fishery	CPUE, length	1990-2014
	Age	1999-2014
U.S. trawl fisheries	Length	1990,1991,1999, 2005-2014
Japan-U.S. cooperative longline survey	CPUE, length	1979-1994
	Age	1981, 1983, 1985, 1987, 1989, 1991, 1993
Domestic longline survey	CPUE, length	1990-2015
	Age	1996-2014
NMFS GOA trawl survey	Abundance index	1984, 1987, 1990, 1993, 1996, 1999, 2003, 2005, 2007, 2009, 2011, 2013, 2015
	Lengths	1984, 1987, 1990, 1993, 1996, 1999, 2003, 2005, 2007, 2009, 2011, 2013

Fishery

Length, catch, and effort data were historically collected from the Japanese and U.S. longline and trawl fisheries, and are now collected from U.S. longline, trawl, and pot fisheries (Table 3.8). The Japanese data were collected by fishermen trained by Japanese scientists (L. L. Low, August 25, 1999, AFSC, pers. comm.). The U.S. fishery length and age data were collected by at-sea and plant observers. No age data were collected from the fisheries until 1999 because of the difficulty of obtaining representative samples from the fishery and because only a small number of sablefish can be aged each year.

Catch

The catches used in this assessment (Table 3.1) include catches from minor State-managed fisheries in the northern GOA and in the AI region because fish caught in these State waters are reported using the area code of the adjacent Federal waters in the Alaska Regional Office catch reporting system (G. Tromble, July 12, 1999, Alaska Regional Office, pers. comm.), the source of the catch data used in this assessment. Minor State fisheries catches averaged 180 t from 1995-1998, about 1% of the average total catch. Most of the catch (80%) is from the AI region. The effect of including these State waters catches in the assessment is to overestimate biomass by about 1%, a negligible error considering statistical variation in other data used in this assessment. Catches from state areas that conduct their own assessments and set Guideline Harvest levels (e.g., Prince William Sound, Chatham Strait, and Clarence Strait), are not included in this assessment.

Some catches probably were not reported during the late 1980's (Kinoshita et al. 1995). Unreported catches could account for the Japan-U.S. cooperative longline survey index's sharp drop from 1989-90 (Table 3.8, Figures 3.3). We tried to estimate the amount of unreported catches by comparing reported catch to another measure of sablefish catch, sablefish imports to Japan, the primary buyer of sablefish. However the trends of reported catch and imports were similar, so we decided to change our approach for catch reporting in the 1999 assessment (Sigler et al. 1999). We assumed that non-reporting is due to at-sea discards, and apply discard estimates from 1994 to 1997 to inflate U.S. reported catches in all years prior to 1993 (2.9% for hook-and-line and 26.6% for trawl).

In response to Annual Catch Limit (ACL) requirements, assessments now document all removals including catch that are not associated with a directed fishery. Research catches of sablefish have been reported in previous stock assessments (Hanselman et al. 2009). Estimates of all removals not associated with a directed fishery including research catches are available and are presented in Appendix 3B. The sablefish research removals are small relative to the fishery catch, but substantial compared to the research removals for many other species. These research removals support a dedicated longline survey. Additional sources of significant removals are bottom trawl surveys and the International Pacific Halibut Commission's longline survey. Other removals are relatively minor for sablefish but the sport fishery catch has been increasing in recent years, but occurs primarily in State waters. Total removals from activities other than directed fishery have been between 239-359 t since 2006. These catches are not included in the stock assessment model. These removal estimates equate to approximately 2% of the recommended ABC and represent a relatively low risk to the sablefish stock.

Lengths

We use length compositions from the U.S. fixed gear (longline and pot) and U.S. trawl fisheries which are both measured by sex. The fixed gear fishery has large sample sizes and has annual data since 1990. The trawl fishery had low levels of observer sampling in much of the 1990s and early 2000s, and has a much smaller sample size than the fixed gear fishery. We only use years for the trawl fishery that have sample sizes of at least 300 per sex. The length compositions are weighted by catch in each FMP management area to obtain a representative estimate of catch-at-length.

Ages

We use age compositions from the U.S. fixed gear fishery since 1999. Sample sizes are similar to the longline survey with about 1,200 otoliths aged every year. The age compositions are weighted by the catch in each area to obtain a representative estimate of catch-at-age.

Longline fishery catch rate index

Fishery information is available from longline sets that target sablefish in the IFQ fishery. Records of catch and effort for these vessels are collected by observers and by vessel captains in voluntary and required logbooks. Fishery data from the Observer Program is available since 1990. Logbooks are required for vessels over 60 feet beginning in 1999. Since 2000, a longline fishery catch rate index has

been derived from observed sets and logbook data for use in the model and in apportionment. The mean CPUE is scaled to a relative population weight by the total area size in each area. In the years that logbook and observer CPUEs are available, the average of the two sources is computed by weighting with the inverse of the coefficient of variation.

Targeted sablefish longline sample sizes

For analysis of observed sablefish catch rates in the sablefish target fishery, we first have to determine the target of the set, because the target is not declared in the observer data set. To do this, we compare the catch of sablefish to other target species that are typically caught on longline gear: Greenland turbot, several rockfish species, Pacific halibut, and Pacific cod. Whichever target fishery has the greatest weight in the set is regarded as the target. Catch rates and sample sizes presented here only include sets where sablefish were determined to be the target. The total weight of all sets recorded by observers determined to be targeting sablefish represent on average 14% of the annual IFQ hook and line catch. In 2014 they comprised 12% of the hook and line catch (1,407 mt). On average, the percent of the IFQ catch observed is lowest in the EY/SE (5%), highest in WY and AI (~22%), and moderate in the BS, CGOA, and WGOA (10-14%). In 2014 20% of catch in the AI catch was observed, 4% in the BS, 9% in the WG, 15% in the CG, 18% in WY, and 10% in EY/SE. In 2014, coverage in EY/SE was higher than average and lower than average in the BS and WY areas. This may partially be due to observer restructuring, where more coverage was directed to smaller vessels, which are more common in EY/SE. Low longline fishery sample sizes in the BS are also likely a result of poor observer coverage for sablefish directed trips and an increase in pot fishing in the BS (Table 3.9). Because of confidentiality concerns, the catch rates with less than three vessels cannot be shown.

Killer whales impact sablefish catch rates in the BS, AI and WGOA and these sets are excluded from catch rate analyses in the observer data set. Whale data is not currently collected in logbooks. Since 2009, there has been an increase in killer whale depredation in the WGOA (average 6% from 2010-2013); however, this is only 7-22 sets per year. In the AI and BS, killer whale depredation has been variable, ranging from 0-12 sets per year in each area. In 2014 there was sperm whale depredation on 10 sets in the CGOA (1.7% of sets). Sperm whale depredation typically occurs in the CGOA, EY/SE, and WY. In 2014 7% of sets in the CG and EY were depredated and 20% in the WY. In 2014 there were a few sets with sperm whale depredation in the AI (6 sets) and in the WGOA (5 sets). The percent of sets affected by sperm whales varies greatly and determining if sperm whales are depredating can be subjective because whales do not take the great majority of the catch, like killer whales do. Therefore, measures of depredation in the fishery may not be accurate.

Logbook sample sizes are substantially higher than observer samples sizes, especially since 2004, and have continued to rise annually in many management areas (WGOA, WY, CGOA) (Table 3.9). Logbook participation increased sharply in 2004 in all areas primarily because the International Pacific Halibut Commission (IPHC) was used to collect, edit, and enter logbooks electronically. This increasing trend is likely due to the strong working relationship the IPHC has with fishermen, their diligence in collecting logbooks dockside, and because many vessels <60 feet are now participating in the program voluntarily. In 2014 68% of the logs collected that targeted sablefish were from vessels <60 ft. There is a higher proportion of the catch documented by logbooks than by observers; 50% of the hook and line catch was documented in logbooks, compared to 12% for observer data. Some data is included in both data sets if logbooks are required and an observer was onboard.

Longline catch rates

Sets where there was killer whale depredation are excluded for catch rate calculations in observer data, but whale depredation is not documented in logbooks and so no data are excluded. In general, catch rates are highest in the EY/SE and WY areas and are lowest in the BS and AI (Table 3.9, Figures 3.5 and 3.6). Recently, catch rate trends in the observer and logbook data have been similar in all areas except WY, where observer data shows a sharp decrease in 2014 (25%) and logbooks show only a 1% decrease. Catch

rates from logbooks were stable or slightly up in all areas while the sparser observer data were a little more variable.


Longline spatial and temporal patterns

Changes in spatial or temporal patterns of the fishery may cause fishery catch rates to be unrepresentative of abundance. For example, fishers sometimes target concentrations of fish, even as geographic distribution shrinks when abundance declines (Crecco and Overholtz 1990). This could lead to an incorrect interpretation of fishery catch rates, which could remain stable while the area occupied by the stock was diminishing (Rose and Kulka 1999).

We examined fishery longline data for seasonal and annual differences in effort and catch rate (CPUE, lbs/hook). Such changes may cause fishery catch rates to be unrepresentative of abundance. In the observed longline data since 2000, the majority of effort occurs in the spring, less in the summer, and least in the fall. Since 1998, catch rates are also highest in the spring, moderate in the summer, and variable in the fall (due to lower sample sizes in the fall). No temporal changes have emerged in the logbook or observer data.

Seasonal changes in fish size

From 2012-2014 there was an increase in the quantity of logbook data providing estimates of catch in weight and numbers. This enables us to examine the average fish weight by season and area. Data from 2012-2014 were combined to increase sample sizes. To further increase sample size, areas were aggregated into BS/AI, CG/WGOA, and WY/EY/SE (EGOA). Data were included unless there was missing weight or count information. There were very small differences between spring, summer, and fall in the west and central areas and larger differences in the EGOA (see figure below). In EGOA, weight in spring was 6.1 lbs, 7.1 lbs in summer, and 8.0 lbs in fall. Although fish size increases in the fall, catch rates and effort decreases.



Count of hook and line logbook sets used for calculations of average weight by area and season.

Area	Spring	Summer	Fall	Total
BS/AI	1,255	925	445	2,625
CG/WG	2,203	1,797	595	4,595
EGOA	1,419	365	166	1,950

Pot fishery catch rate analysis

Pot catch rates: Because pot data are sparser than longline data, and in some years is confidential due to fewer than 3 vessels participating, specific annual data are not presented. In addition, it is difficult to discern trends, since pot catch rates have wider confidence intervals than longline data due to smaller

sample sizes. Observed sets are determined to be targeting sablefish if sablefish comprise the greatest weight in the set. Overall, there are more vessels in both the logbook and observer data from the sablefish pot fishery in the BS than the AI. Since 2006, in the BS there have been from 3 to 9 vessels in logbook data and 5 to 8 vessels in observer data. In the AI, there have been from 1 to 5 vessels in logbooks and 1 to 4 in observer data.

In logbook data, since 2009 the number of pots, sets, and vessels has decreased. For example, in logbooks in 2014, only 276 sets were reported in the BS and 1,284 in 2009. From 2006-2014 the average catch rate in logbook data was 29 lbs/pot in the AI (number sets (n) = 1,271) and 18 lbs/pot in the BS (n = 3,237). The average catch rate in the observer data from 2006-2014 was 11 lbs/pot (n = 1,156) in the AI and 18 lbs/pot (n = 2,970) in the BS. The effort recorded by observers has also been decreasing since 2009 in the BS and 2011 in the AI. Pot effort is approximately equal throughout the fishing season, unlike hook and line fishing where effort is highest in the spring.

The composition of bycatch species caught in observed pots that retained sablefish in the BS and AI is comprised mostly of arrowtooth/Kamchatka flounder, Greenland turbot, Pacific halibut, giant grenadier, and snails. The estimated catch of golden king crab in the pot fishery was high in 2009 (Hanselman et al. 2010), but in 2014 it was 98% lower than 2009.

Surveys

A number of fishery independent surveys catch sablefish. The survey indices included in the model for this assessment are the AFSC longline survey and the AFSC GOA bottom trawl survey. For other surveys that occur in the same or adjacent geographical areas, but are not included as separate indices in the model, we provide trends and comparative analyses to the AFSC longline survey. Research catch removals including survey removals are documented in Appendix 3B.

AFSC Surveys

Longline survey

Overview: Catch, effort, age, length, weight, and maturity data are collected during sablefish longline surveys. These longline surveys likely provide an accurate index of sablefish abundance (Sigler 2000). Japan and the United States conducted a cooperative longline survey for sablefish in the GOA annually from 1978 to 1994, adding the AI region in 1980 and the eastern BS in 1982 (Sasaki 1985, Sigler and Fujioka 1988). Since 1987, the Alaska Fisheries Science Center has conducted annual longline surveys of the upper continental slope, referred to as domestic longline surveys, designed to continue the time series of the Japan-U.S. cooperative survey (Sigler and Zenger 1989). The domestic longline survey began annual sampling of the GOA in 1987, biennial sampling of the AI in 1996, and biennial sampling of the eastern BS in 1997 (Rutecki et al. 1997). The domestic survey also samples major gullies of the GOA in addition to sampling the upper continental slope. The order in which areas are surveyed was changed in 1998 to reduce interactions between survey sampling and short, intense fisheries. Before 1998, the order was AI and/or BS, Western Gulf, Central Gulf, Eastern Gulf. Starting in 1998, the Eastern Gulf area was surveyed before the Central Gulf area.

Specimen collections: Sablefish length data were randomly collected for all survey years. Otoliths were collected for age determination for most survey years. From 1979-1994 otolith collections were length-stratified; since 1994 otoliths have been collected randomly. Prior to 1996, otolith collections were aged but not consistently from year to year. Since 1996, a sample of otoliths collected during each survey has been aged in the years they were collected. Approximately one-half of the otoliths collected (~1,000) are aged annually. This sample size for age compositions should be large enough to get a precise age composition for the whole survey area, but may be too small to estimate the age composition in smaller areas by sex (P. Hulson, unpublished manuscript).

Standardization: Kimura and Zenger (1997) compared the performance of the two surveys from 1988 to

1994 in detail, including experiments comparing hook and gangion types used in the two surveys. The abundance index for both longline surveys decreased from 1988 to 1989, the cooperative survey decreased from 1989 to 1990, while the domestic survey increased (Table 3.9). Kimura and Zenger (1997) attributed the difference to the domestic longline survey not being standardized until 1990.

Survey Trends: Relative population abundance indices are computed annually using survey catch rates from stations sampled on the continental slope. Highest sablefish abundance indices occurred during the Japan-U.S. cooperative survey in the mid-1980's, in response to exceptional recruitment in the late 1970's (Figure 3.7). Relative population numbers declined through the 1990's in most areas during the domestic longline survey. Catches increased in the early 2000's but have trended down since 2006.

The 2013 and 2015 survey estimates of relative abundance in numbers (RPN) are the lowest points in the domestic time series despite modest increases in 2010, 2011 and 2014. The index remains below average because of recent weak recruitment.

Whale Depredation: Killer whale depredation of the survey's sablefish catches has been a problem in the BS since the beginning of the survey (Sasaki 1987). Killer whale depredation primarily occurs in the BS, AI, WGOA, and to a lesser extent in recent years in the CGOA (Table 3.11). Depredation is easily identified by reduced sablefish catch and the presence of lips or jaws and bent, straightened, or broken hooks. Since 1990, portions of the gear at stations affected by killer whale depredation during the domestic longline survey have been excluded from the analysis of catch rates, RPNs, and RPWs. The AI and the BS were added to the domestic longline survey in 1996 and this is when killer whale depredation increased. In 2009, 10 BS stations were depredated, which significantly impacted catch and biased the abundance index leading to using the 2007 BS RPN estimate to interpolate the 2009 and 2010 BS RPNs (Hanselman et al. 2009). In 2011, depredation levels in the BS were similar to previous years with catches at 7 of 16 stations affected. In 2013, a new high of 11 stations were depredated, although fewer skates were impacted and therefore removed from the analysis in comparison to what occurred in 2009. In the AI depredation was highest in 2012 and in 2014 was back to levels seen in 2008 and 2010.

In 2015 killer whale depredation was similar to recent years (Table 3.11). The number of stations in the BS was down to 9 from 11 stations in 2013. Although there has been some killer whale depredation in the CGOA in the past (1 - 2 stations), this year there was none.

Sperm whale depredation affects longline catches, but evidence of depredation is not accompanied by obvious decreases in sablefish catch or common occurrence of lips and jaws or bent and broken hooks. Data on sperm whale depredation have been collected since the 1998 longline survey (Table 3.11). Sperm whales are often observed from the survey vessel during haulback but do not appear to be depredating on the catch. Sperm whale depredation and presence is recorded during the longline survey at the station level, not the skate level like killer whales. Depredation is defined as sperm whales being present during haulback with the occurrence of damaged fish in the catch.

Sperm whale depredation is variable, but has generally been increasing since 1998 (Table 3.11). Whales are most common in the EGOA (WY and EY/SE), but are also seen in the CGOA. In 2015 there were sperm whales depredating at 19 stations (annual range 4-21) (Table 3.11). In 2015 in the CGOA sperm whales were present at 9 stations and depredating at 6, which is higher than in other years. Although sperm whales are sometimes observed in the WGOA, in 2015 there were no sightings.

Multiple studies have attempted to quantify sperm whale depredation rates. An early study using data collected by fisheries observers in Alaskan waters found no significant effect on the commercial fishery catch (Hill et al. 1999). Another study using data collected from commercial vessels in southeast Alaska, found a small, significant effect comparing longline fishery catches between sets with sperm whales present and sets with sperm whales absent (3% reduction, 95% CI of (0.4 – 5.5%), t-test, $p = 0.02$, Straley et al. 2005).

A general linear model fit to longline survey data from 1998-2004 found neither sperm whale presence (p

= 0.71) nor depredation rate ($p = 0.78$) increased significantly from 1998 to 2004. Catch rates were about 2% less at locations where depredation occurred, but the effect was not significant ($p = 0.34$). This analysis was updated through 2009 and now shows a significant effect of approximately four kilograms per hundred hooks in the Central and Eastern Gulf regions, which translates into approximately a 2% decrease in overall catch in those areas (J. Liddle, October, 2009, pers. comm.). A retrospective analysis of this data indicates the effect is not significant until the 2009 data are added, indicating the increasing depredation effect has combined with accumulating survey data to give increased power to detect this small reduction in CPUE.

Longline survey catch rates are not adjusted for sperm whale depredation because we do not know when measurable depredation began during the survey time series, because past studies of depredation on the longline survey showed no significant effect, and because sperm whale depredation is difficult to detect (Sigler et al. 2007). Because of recent increases in sperm whale presence and depredation at survey stations, as indicated by whale observations and significant results of recent studies, we evaluated a statistical adjustment to survey catch rates using a general linear modeling approach (Appendix 3C, Hanselman et al. 2010). This approach had promise but had issues with variance estimation and autocorrelation between samples. A new approach has been developed using a generalized linear mixed model that resolves these issues (see Appendix 3C in Hanselman et al. 2014).

Gully Stations: In addition to the continental slope stations sampled during the survey, twenty-seven stations are sampled in gullies at the rate of one to two stations per day. The sampled gullies are Shelikof Trough, Amatuli Gully, W-grounds, Yakutat Valley, Spencer Gully, Ommaney Trench, Dixon Entrance, and one station on the continental shelf off Baranof Island. The majority of these stations are located in deep gully entrances to the continental shelf in depths from 150-300 m in areas where the commercial fishery targets sablefish. No gullies are currently sampled in the Western GOA, AI, or BS.

Previous analyses have shown that on average gully stations catch fewer large fish and more small fish than adjacent slope stations (Rutecki et al. 1997, Zenger et al. 1994). Compared with the adjacent regions of the slope, sablefish catch rates for gully stations have been mixed with no significant trend (Zenger et al. 1994). Gully catches may indicate recruitment signals before slope areas because of their shallow depth, where younger, smaller sablefish typically inhabit. Catch rates from these stations have not been included in the historical abundance index calculations because preferred habitat of adult sablefish is on the slope.

These areas do support significant numbers of sablefish, however, and are important areas sampled by the survey. We compared the RPNs of gully stations to the RPNs of slope stations in the GOA to see if catches were comparable, or more importantly, if they portrayed different trends than the RPNs used in this assessment.

To compare trends, we computed Student's- t normalized residuals for all GOA gullies and slope stations and plotted them for the time series. If the indices were correlated, then the residuals would track one another over time (Figure 3.8). Overall, gully catches in the GOA from 1990-2014 are moderately correlated with slope catches ($r = 0.55$). There is no evidence of major differences in trends. In regards to gully catches being a recruitment indicator, the increase in the gully RPNs in 1999 and 2001-2002 may be in response to the above average 1997 and 2000 year classes. Both the 2001 and 2002 RPNs for the gully stations are higher than in 1999, which supports the current model estimate that the 2000 year class was larger than 1997. Both gully and slope trends were down in 2012 and 2013, consistent with the overall decrease in survey catch. However, the slope stations increased in 2014, while the gullies continued to decline. In 2015, the opposite pattern occurred, with the gullies showing a slight uptick while the slope stations declined again. In the future, we will continue to explore sablefish catch rates in gullies and explore their usefulness for indicating recruitment; they may also be useful for quantifying depredation, since sperm whales have rarely depredated on catches from gully stations.

Interactions between the fishery and survey are described in Appendix 3A.

Trawl surveys

Trawl surveys of the upper continental slope that adult sablefish inhabit have been conducted biennially or triennially since 1980 in the AI, and 1984 in the GOA, always to 500 m and occasionally to 700-1000 m. Trawl surveys of the BS slope were conducted biennially from 1979-1991 and redesigned and standardized for 2002, 2004, 2008, 2010, and 2012. Trawl surveys of the BS shelf are conducted annually but generally catch no sablefish. Trawl survey abundance indices were not used in the assessment model prior to 2007 in the sablefish assessment because they were not considered good indicators of the sablefish relative abundance. However, there is a long time series of data available and given the trawl survey's ability to sample smaller fish, it may be a better indicator of recruitment than the longline survey.

There is some difficulty with combining estimates from the BS and AI with the GOA estimates since they occur on alternating years. A method could be developed to combine these indices, but it leaves the problem of how to use the length data to predict recruitment since the data could give mixed signals on year class strength. At this time we are using only the GOA trawl survey biomass estimates (<500 m depth, Figure 3.4) and length data (<500 m depth) as a recruitment index for the whole population. The largest proportion of sablefish biomass is in the GOA so it should be indicative of the overall population. Biomass estimates used in the assessment for 1984-2013 are shown in Table 3.10. The GOA trawl survey index was at its lowest level of the time series in 2013, but increased 12% in 2015 from the 2013 estimate.

AI and BS Slope survey biomass estimates are not used in the assessment model but are tracked in Figure 3.9. Estimates in the two areas have decreased slowly since 2000.

Other surveys/areas not used in the assessment model

IPHC Longline Surveys

The IPHC conducts a longline survey each year to assess Pacific halibut. This survey differs from the AFSC longline survey in gear configuration and sampling design, but catches substantial numbers of sablefish. More information on this survey can be found in Soderlund et al. (2009). A major difference between the two surveys is that the IPHC survey samples the shelf consistently from ~ 10-500 meters, whereas the AFSC survey samples the slope and select gullies from 200-1000 meters. Because the majority of effort occurs on the shelf in shallower depths, the IPHC survey may catch smaller and younger sablefish than the AFSC survey; however, lengths of sablefish are not taken on the IPHC survey.

For comparison to the AFSC survey, IPHC relative population number's (RPN) were calculated using the same methods as the AFSC survey values, the only difference being the depth stratum increments. Area sizes used to calculate biomass in the RACE trawl surveys were utilized for IPHC RPN calculations.

We do not obtain IPHC survey estimates for the current year until the following year. We compared the IPHC and the AFSC RPNs for the GOA (Figure 3.10). The two series track well, but the IPHC survey RPN has more variability. This is likely because it surveys shallower water on the shelf where younger sablefish reside and are more patchily distributed. Since the abundance of younger sablefish will be more variable as year classes pass through, the survey more closely resembles the NMFS GOA trawl survey index described above which samples the same depths (Figure 3.10b).

While the two longline surveys have shown consistent patterns for most years, they diverged in 2010 and 2011, but the 2013 estimates both show the lowest point in the time series for each index (Figure 3.10). The IPHC estimate for the Gulf of Alaska for 2013 was a 21% decline from 2012. The uptick seen in 2014 in the AFSC survey was not apparent in the IPHC survey. We will continue to examine trends in each region and at each depth interval for evidence of recruiting year classes and for comparison to the AFSC longline survey. There is some effort in depths shallower than 200 meters on the AFSC longline survey, and we recently have computed RPNs for these depths for future comparisons with the IPHC RPNs.


Alaska Department of Fish and Game

The Alaska Department of Fish and Game conducts mark-recapture and a longline survey in Northern Southeast Alaska Inside (NSEI) waters. Sablefish in this area are treated as a separate population, but some migration into and out of Inside waters has been confirmed with tagging studies (Hanselman et al. 2015). This population seems to be stabilizing from previous steep declines. Their longline survey CPUE estimates (Figure 3.11a) and fishery CPUE estimates (Figure 3.11b) had been slowly increasing since 2000, confirming the lows in 1999/2000 estimated in our assessment. Like the AFSC longline survey, there was a sharp decline in the 2013 longline survey CPUE estimates for NSEI and a slight uptick in 2014.

Department of Fish and Oceans of Canada

In a 2011 Science Advisory Report, DFO reported : “*Stock reconstructions suggest that stock status is currently below B_{MSY} for all scenarios, with the stock currently positioned in the mid-Cautious to low-Healthy zones.*” Under these scenarios, recent harvest rates on adult sablefish potentially have been between 0.06 – 0.15¹.

The stratified random trap survey was up approximately 29% from 2012 to 2013 after a time series low in 2012 (see figure below) but has registered a new time series low in 2014. The estimated biomass trend in B.C. is similar to the trend in Alaska (see figure below)². The similarly low abundance south of Alaska concerns us, and points to the need to better understand the contribution to Alaska sablefish productivity from B.C. sablefish. Some potential ideas are to conduct an area-wide study of sablefish tag recoveries, and to attempt to model the population to include B.C. sablefish and U.S. West Coast sablefish.



¹ Science Advisory Report 2011/25: http://www.dfo-mpo.gc.ca/Csas-sccs/publications/sar-as/2011/2011_025-eng.pdf

² DFO. 2014. Performance of a revised management procedure for Sablefish in British Columbia. DFO Can. Sci. Advis. Sec. Sci. Resp. 2014 /025: http://www.dfo-mpo.gc.ca/csas-sccs/publications/scr-rs/2014/2014_025-eng.html

Overall abundance trends

Relative abundance has cycled through three valleys and two peaks near 1970 and 1985 (Table 3.10, Figures 3.3 and 3.4). The post-1970 decrease likely is due to heavy fishing. The 1985 peak likely is due to the exceptionally large late 1970's year classes. Since 1988, relative abundance has decreased substantially. Regionally, abundance decreased faster in the BS, AI, and western GOA and more slowly in the central and eastern GOA (Figure 3.7). The majority of the surveys show that sablefish were at their lowest levels in the early 2000s, with current abundance reaching these lows again in 2014 in the central and eastern GOA, and in 2015 in the western areas.

Analytic approach

Model Structure

The sablefish population is assessed with an age-structured model. The analysis presented here extends earlier age structured models developed by Kimura (1990) and Sigler (1999), which all stem from the work by Fournier and Archibald (1982). The current model configuration follows a more complex version of the GOA Pacific ocean perch model (Hanselman et al. 2005a); it includes split sexes and many more data sources to attempt to more realistically represent the underlying population dynamics of sablefish. The current configuration was accepted by the Groundfish Plan Team and NPFMC in 2010 ("Moonwater", Hanselman et al. 2010). The population dynamics and likelihood equations are described in Box 1. The analysis was completed using AD Model Builder software, a C++ based software for development and fitting of general nonlinear statistical models (Fournier et al. 2012).

Parameters Estimated Outside the Assessment Model

The following table lists the parameters estimated independently:

Parameter name	Value	Value	Source
Time period	1960-1995	1996-current	
Natural mortality	0.1	0.1	Johnson and Quinn (1988)
Female maturity-at-age	$m_a = 1/(1+e^{-0.84(a-6.60)})$		Sasaki (1985)
Length-at-age - females	$\bar{L}_a = 75.6(1-e^{-0.208(a+3.63)})$	$\bar{L}_a = 80.2(1-e^{-0.222(a+1.95)})$	Hanselman et al. (2007)
Length-at-age - males	$\bar{L}_a = 65.3(1-e^{-0.227(a+4.09)})$	$\bar{L}_a = 67.8(1-e^{-0.290(a+2.27)})$	Hanselman et al. (2007)
Weight-at-age - females	$\ln \hat{W}_a = \ln(5.47) + 3.02 \ln(1-e^{-0.238(a+1.39)})$		Hanselman et al. (2007)
Weight-at-age - males	$\ln \hat{W}_a = \ln(3.16) + 2.96 \ln(1-e^{-0.356(a+1.13)})$		Hanselman et al. (2007)
Ageing error matrix	From known-age tag releases, extrapolated for older ages		Heifetz et al. (1999)
Recruitment variability (σ_r)	1.2	1.2	Sigler et al. (2002)

Age and Size of Recruitment: Juvenile sablefish rear in nearshore and continental shelf waters, moving to the upper continental slope as adults. Fish first appear on the upper continental slope, where the longline survey and longline fishery occur, at age 2, and a fork length of about 45 cm. A higher proportion of young fish are susceptible to trawl gear compared to longline gear because trawl fisheries usually occur on the continental shelf and shelf break inhabited by younger fish, and catching small sablefish may be hindered by the large bait and hooks on longline gear.

Sablefish are difficult to age, especially those older than eight years (Kimura and Lyons 1991). To compensate, we use an ageing error matrix based on known-age otoliths (Heifetz et al. 1999; Hanselman

et al. 2012).

Growth and maturity: Sablefish grow rapidly in early life, growing 1.2 mm d^{-1} during their first spring and summer (Sigler et al. 2001). Within 100 days after first increment (first daily otolith mark for larvae) formation, they average 120 mm. Sablefish are currently estimated to reach average maximum lengths and weights of 68 cm and 3.2 kg for males and 80 cm and 5.5 kg for females (Echave et al. 2012).

New growth relationships were estimated in 2007 because many more age data were available (Hanselman et al. 2007); this analysis was accepted by the Plan Team in November 2007 and published in 2012 (Echave et al. 2012). We divided the data into two time periods based on the change in sampling design that occurred in 1995. It appears that sablefish maximum length and weight has increased slightly over time. New age-length conversion matrices were constructed using these curves with normal error fit to the standard deviations of the collected lengths at age (Figure 3.12). These new matrices provided for a superior fit to the data. Therefore, we use a bias-corrected and updated growth curve for the older data (1981-1993) and a new growth curve describing recent randomly collected data (1996-2004).

Fifty percent of females are mature at 65 cm, while 50 percent of males are mature at 57 cm (Sasaki 1985), corresponding to ages 6.6 for females and 5 for males (Table 3.12). Maturity parameters were estimated independently of the assessment model and then incorporated into the assessment model as fixed values. The maturity - length function is $m_l = 1 / (1 + e^{-0.40(L - 57)})$ for males and $m_l = 1 / (1 + e^{-0.40(L - 65)})$ for females. Maturity at age was computed using logistic equations fit to the length-maturity relationships shown in Sasaki (1985, Figure 23, GOA). Prior to the 2006 assessment, average male and female maturity was used to compute spawning biomass. Beginning with the 2006 assessment, female-only maturity has been used to compute spawning biomass. Female maturity-at-age from Sasaki (1985) is described by the logistic fit of $m_a = 1/(1+e^{-0.84(a-6.6)})$. In 2011, the AFSC conducted a winter cruise out of Kodiak to sample sablefish when they are preparing to spawn. Ovaries were examined histologically to determine maturity for a study of the age at maturity and fecundity. Skipped spawning was documented for the first time in sablefish. These winter samples provided a similar age at 50% maturity estimate (6.8 years) as the mean of visual observations taken during summer surveys from 1996-2012 (mean = 7.0 years) and the estimate currently used in the assessment (mean = 6.6 years), when skipped spawners were classified as mature. Skipped spawners were primarily found in gullies on the shelf and was positively correlated with age. A second survey will take place in December 2015 in the same areas that were sampled in 2011. Future analyses will aim to develop and evaluate methods to incorporate skipped spawning into maturity ogives and to better utilize the time series of visual maturity estimates.

Maximum age and natural mortality: Sablefish are long-lived; ages over 40 years are regularly recorded (Kimura et al. 1993). Reported maximum age for Alaska is 94 years (Kimura et al. 1998). Canadian researchers report age determinations up to 113 years¹. A natural mortality rate of $M=0.10$ has been assumed for previous sablefish assessments, compared to $M=0.112$ assumed by Funk and Bracken (1984). Johnson and Quinn (1988) used values of 0.10 and 0.20 in a catch-at-age analysis and found that estimated abundance trends agreed better with survey results when $M=0.10$ was used. Natural mortality has been modeled in a variety of ways in previous assessments. For sablefish assessments before 1999, natural mortality was assumed to equal 0.10. For assessments from 1999 to 2003, natural mortality was estimated rather than assumed to equal 0.10; the estimated value was about 0.10 but only with a precise prior imposed. For the 2004 assessment, a more detailed analysis of the posterior probability showed that natural mortality was not well-estimated by the available data (Sigler et al. 2004). Therefore in 2006, we returned to fixing the parameter at 0.10.

Variance and effective sample sizes: Several quantities were computed in order to compare the variance of the residuals to the assumed input variances. The standardized deviation of normalized residuals

¹Fisheries and Oceans Canada; <http://www.pac.dfo-mpo.gc.ca/fm-gp/commercial/ground-fond/sable-charbon/bio-eng.htm>

(SDNR) is closely related to the root mean squared error (RMSE) or effective sample size; values of SDNR of approximately 1 indicate that the model is fitting a data component as well as would be expected for a given specified input variance. The normalized residuals for a given year i of the abundance index was computed as

$$\delta_i = \frac{\ln(I_i) - \ln(\hat{I}_i)}{\sigma_i}$$

where σ_i is the input sampling log standard deviation of the estimated abundance index. For age or length composition data assumed to follow a multinomial distribution, the normalized residuals for age/length group a in year i were computed as

$$\delta_{i,a} = \frac{(y_{i,a} - \hat{y}_{i,a})}{\sqrt{\hat{y}_{i,a}(1 - \hat{y}_{i,a})/n_i}}$$

where y and \hat{y} are the observed and estimated proportion, respectively, and n is the input assumed sample size for the multinomial distribution. The effective sample size was also computed for the age and length compositions modeled with a multinomial distribution, and for a given year i was computed as

$$E_i = \frac{\sum_a \hat{y}_a * (1 - \hat{y}_a)}{\sum_a (\hat{y}_a - y_a)^2}.$$

An effective sample size that is nearly equal to the input sample size can be interpreted as having a model fit that is consistent with the input sample size.

For the 2010 recommended assessment model, we used average SDNR as a criterion to help reweight the age and length compositions. SDNR is a common metric used for goodness of fit in other fisheries, particularly in New Zealand (e.g. Langley and Maunder 2009) and has been recommended for use in fisheries models in Alaska during multiple CIE reviews, such as Atka mackerel and rockfish. We iteratively reweighted the model by setting an objective function penalty to reduce the deviations of average SDNR of a data component from one. Initially, we tried to fit all multinomial components this way, but due to tradeoffs in fit, it was found that the input sample sizes became too large and masked the influence of important data such as abundance indices. Given that we have age and length samples from nearly all years of the longline surveys, we chose to eliminate the attempt to fit the length data well enough to achieve an average SDNR of one, and reweighted all age components and only length components where no age data exists (e.g. domestic trawl fishery). The abundance index SDNRs were calculated, but no attempt was made to adjust their input variance because we have *a priori* knowledge about their sampling variances. This process was completed before the 2010 data were added into the assessment and endorsed by the Plan Teams and SSC in 2010. We continue to use these weightings. The table below shows the input CVs/sample sizes for the data sources and their associated output SDNR for the recommended model. This reweighting is intended to remain fixed for at least several years. The data weights in general continue to do well by these objectives (Table 3.13).

Parameters Estimated Inside the Assessment Model

Below is a summary of the parameters estimated within the recommended assessment model:

Parameter name	Symbol	Number of
Catchability	q	6
Log-mean-recruitment	μ_r	1
Spawners-per-recruit levels	F_{35}, F_{40}, F_{50}	3
Recruitment deviations	τ_y	83
Average fishing mortality	μ_f	2
Fishing mortality deviations	ϕ_v	112
Fishery selectivity	fs_a	8
Survey selectivity	ss_a	7
Total		222

Catchability is separately estimated for the Japanese longline fishery, the cooperative longline survey, the domestic longline survey, U.S. longline derby fishery, U.S. longline IFQ fishery, and the NMFS GOA trawl survey. Information is available to link these estimates of catchability. Kimura and Zenger (1997) analyzed the relationship between the cooperative and domestic longline surveys. For assessments through 2006, we used their results to create a prior distribution which linked catchability estimates for the two surveys. For 2007, we estimated new catchability prior distributions based on the ratio of the various abundance indices to a combined Alaskan trawl index. This resulted in similar mean estimates of catchability to those previously used, but allowed us to estimate a prior variance to be used in the model. This also facilitates linking the relative catchabilities between indices. These priors were used in the recommended model for 2008. This analysis was presented at the September 2007 Plan Team and is presented in its entirety in Hanselman et al. (2007). Lognormal prior distributions were used with the parameters shown below:

Index	U.S. LL Survey	Jap. LL Survey	Fisheries	GOA Trawl
Mean	7.857	4.693	4.967	0.692
CV	33%	24%	33%	30%

Recruitment is not estimated with a stock-recruit relationship, but is estimated with a level of average recruitment with deviations from average recruitment for the years 1933-2014.

Fishing mortality is estimated with two average fishing mortality parameters for the two fisheries (fixed gear and trawl) and deviations from the average for years 1960-2015 for each fishery.

Selectivity is represented using a function and is separately estimated by sex for the longline survey, fixed-gear fishery (pot and longlines combined), and the trawl survey. Selectivity for the longline surveys and fixed-gear fishery is restricted to be asymptotic by using the logistic function. Selectivity for the trawl fishery and trawl survey are dome-shaped (right descending limb) and estimated with a two-parameter gamma-function and a power function respectively (see Box 1 for equations). This right-descending limb is allowed because we do not expect that the trawl survey and fishery will catch older aged fish as frequently because they fish shallower than the fixed-gear fishery. Selectivity for the fixed-gear fishery is estimated separately for the “derby” fishery prior to 1995 and the IFQ fishery from 1995 thereafter. Fishers may choose where they fish in the IFQ fishery, compared to the crowded fishing grounds during the 1985-1994 “derby” fishery, when fishers reportedly often fished in less productive depths due to crowding (Sigler and Lunsford 2001). In choosing their ground, they presumably target bigger, older fish, and depths that produce the most abundant catches.

Bayesian analysis of reference points

Since the 1999 assessment, we have conducted a limited Bayesian analysis of assessment uncertainty. The

posterior distribution was computed based on 10 million MCMC simulations drawn from the posterior distribution. A burn-in of 1 million draws was removed from the beginning of the chain and then thinned to 4,000 parameter draws to remove serial correlation between successive draws. This was determined to be sufficient through simple chain plots, and comparing the means and standard deviations of the first half of the chain with the second half.

In previous assessments, we estimated the posterior probability that projected abundance will fall below the decision analysis thresholds based on Mace and Sissenwine (1993). However, in the North Pacific Fishery Management Council setting we have thresholds that are defined in the Council harvest rules. These are when the spawning biomass falls below $B_{40\%}$, $B_{35\%}$, and when the spawning biomass falls below $\frac{1}{2}$ MSY or $B_{17.5\%}$ which calls for a rebuilding plan under the Magnuson-Stevens Act. For the previous analysis based on Mace and Sissenwine (1993), see Hanselman et al. 2005b. To examine the posterior probability, we project spawning biomass into the future with recruitments varied as random draws from a lognormal distribution with the mean and standard deviation of 1979-2013 age-2 recruitments. The fishing mortality used is the current yield ratio described in the *Catch specification* section multiplied by maxABC for each year.

Box 1 Model Description

Y	Year, $y=1, 2, \dots, T$
T	Terminal year of the model
A	Model age class, $a = a_0, a_0+1, \dots, a_+$
a_0	Age at recruitment to the model
a_+	Plus-group age class (oldest age considered plus all older ages)
L	Length class
Ω	Number of length bins (for length composition data)
G	Gear-type (g = longline surveys, longline fisheries, or trawl fisheries)
X	Index for likelihood component
$w_{a,s}$	Average weight at age a and sex s
φ_a	Proportion of females mature at age a
μ_r	Average log-recruitment
μ_f	Average log-fishing mortality
$\phi_{y,g}$	Annual fishing mortality deviation
τ_y	Annual recruitment deviation $\sim \ln(0, \sigma_r)$
σ_r	Recruitment standard deviation
$N_{y,a,s}$	Numbers of fish at age a in year y of sex s
M	Natural mortality
$F_{y,a,g}$	Fishing mortality for year y , age class a and gear g
$Z_{y,a}$	Total mortality for year y and age class a ($= \sum_g F_{y,a,g} + M$)
R_y	Recruitment in year y
B_y	Spawning biomass in year y
$s_{a,s}^g$	Selectivity at age a for gear type g and sex s
$A_{50\%}, d_{50\%}$	Age at 50% selection for ascending limb, age at 50% deselection for descending limb
δ	Slope/shape parameters for different logistic curves
\mathbf{A}	Ageing-error matrix dimensioned $a_+ \times a_+$
\mathbf{A}_s^l	Age to length conversion matrix by sex s dimensioned $a_+ \times \Omega$
q_g	Abundance index catchability coefficient by gear
λ_x	Statistical weight (penalty) for component x
I_y, \hat{I}_y	Observed and predicted survey index in year y
$P_{y,l,s}^g, \hat{P}_{y,l,s}^g$	Observed and predicted proportion at length l for gear g in year y and sex s
$P_{y,a,s}^g, \hat{P}_{y,a,s}^g$	Observed and predicted proportion at observed age a for gear g in year y and sex s
ψ_y^g	Sample size assumed for gear g in year y (for multinomial likelihood)
n_g	Number of years that age (or length) composition is available for gear g
$q_{\mu,g}, \sigma_{q,g}$	Prior mean, standard deviation for catchability coefficient for gear g
M_{μ}, σ_M	Prior mean, standard deviation for natural mortality
$\sigma_{r,\mu}, \sigma_{\sigma_r}$	Prior mean, standard deviation for recruitment variability

Equations describing state dynamics	Model Description (continued)
$N_{1,a} = \begin{cases} R_1, & a = a_0 \\ e^{(\mu_r + \tau_{a_0 - a + 1})} e^{-(a - a_0)M}, & a_0 < a < a_+ \\ e^{(\mu_r)} e^{-(a - a_0)M} (1 - e^{-M})^{-1}, & a = a_+ \end{cases}$	Initial year recruitment and numbers at ages.
$N_{y,a} = \begin{cases} R_y, & a = a_0 \\ N_{y-1,a-1} e^{-Z_{y-1,a-1}}, & a_0 < a < a_+ \\ N_{y-1,a-1} e^{-Z_{y-1,a-1}} + N_{y-1,a} e^{-Z_{y-1,a}}, & a = a_+ \end{cases}$	Subsequent years recruitment and numbers at ages
$R_y = e^{(\mu_r + \tau_y)}$	Recruitment
Selectivity equations	
$s_{a,s}^g = \left(1 + e^{(-\delta_{g,s}^g (a - a_{50\%g,s}))}\right)^{-1}$	Logistic selectivity
$s_{a,s}^g = \frac{a^{\delta_{g,s}^g}}{\max(s_{a,s}^g)}$	Inverse power family
$s_{a,s}^g = \left(\frac{a}{a_{\max}}\right)^{a_{\max,g,s} / p} e^{(a_{\max,g,s} - a) / p}$	Reparameterized gamma distribution
$p = 0.5 \left[\sqrt{a_{\max,g,s}^2 + 4\delta_{g,s}^2} - a_{\max,g,s} \right]$	
$s_{a,s}^g = (1 - \varphi_s^g)^{-1} \left(\frac{(1 - \varphi_s^g)}{\varphi_s^g} \right)^{\varphi_s^g} \frac{e^{(\delta_{g,s}^g \varphi_s^g (a_{50\%g,s} - a))}}{(1 + e^{(\delta_{g,s}^g (a_{50\%g,s} - a))})}$	Exponential-logistic selectivity
Observation equations	
$\hat{C}_{y,g} = \sum_1^g \sum_1^s w_{a,s} N_{y,a,g,s} F_{y,a,g,s} \left(1 - e^{-Z_{y,a,g,s}}\right) Z_{y,a,g,s}^{-1}$	Catch biomass in year y
$\hat{I}_{y,g} = q^g \sum_{a_0}^{a_+} \sum_1^s N_{y,a,s} \frac{s_{a,s}^g}{\max(s_{a,s}^g)} w_{a,s}$	Survey biomass index (weight)
$\hat{I}_{y,g} = q^g \sum_{a_0}^{a_+} \sum_1^s N_{y,a,s} \frac{s_{a,s}^g}{\max(s_{a,s}^g)}$	Survey abundance index (numbers)
$\hat{P}_{y,a,s}^g = N_{y,a,s} s_{a,s}^g \left(\sum_{a_0}^{a_+} N_{y,a,s} s_{a,s}^g \right)^{-1} \mathbf{A}_s$	Vector of fishery or survey predicted proportions at age
$\hat{P}_{y,a,s}^g = N_{y,s} s_s^g \left(\sum_{a_0}^{a_+} N_{y,a,s} s_{a,s}^g \right)^{-1} \mathbf{A}'_s$	Vector of fishery or survey predicted proportions at length

Posterior distribution components

$$L_C = \lambda_c \sum_1^g \sum_y \left(\ln C_{g,y} - \ln \hat{C}_{g,y} \right)^2 / (2\sigma_C^2)$$

$$L_I = \lambda_I \sum_1^g \sum_y \left(\ln I_{g,y} - \ln \hat{I}_{g,y} \right)^2 / (2\sigma_I^2)$$

$$L_{age} = \lambda_{age} \sum_{i=1}^{n_g} -\psi_y^g \sum_{a_0}^{a_+} (P_{i,a}^g + \nu) \ln(\hat{P}_{i,a}^g + \nu)$$

$$L_{length} = \lambda_{length} \sum_1^s \sum_{i=1}^{n_g} -\psi_y^g \sum_{l=1}^{\Omega} (P_{i,l,s}^g + \nu) \ln(\hat{P}_{i,l,s}^g + \nu)$$

$$L_q = \left(\ln \hat{q}^g - \ln q_{\mu}^g \right)^2 / 2\sigma_q^2$$

$$L_M = \left(\ln \hat{M} - \ln M_{\mu} \right)^2 / 2\sigma_M^2$$

$$L_{\sigma_r} = \left(\ln \hat{\sigma}_r - \ln \sigma_{r,\mu} \right)^2 / 2\sigma_{\sigma_r}^2$$

$$L_{\tau} = 0.1 \sum_{y=1}^T \frac{\tau_y^2}{2\hat{\sigma}_r^2} + n \ln \hat{\sigma}_r$$

$$L_f = \lambda_f \sum_1^g \sum_{y=1}^T \phi_{y,g}^2$$

$$L_{Total} = \sum_x L_x$$

Model Description (continued)

Catch likelihood

Survey biomass index likelihood

Age composition likelihood

Length composition likelihood

(ψ_y^g = sample size, n_g = number of years of data for gear g , i = year of data availability, ν is a constant set at 0.001)

Prior on survey catchability coefficient for gear g

Prior for natural mortality

Prior distribution for σ_r

Prior on recruitment deviations

Regularity penalty on fishing mortality

Total objective function value

Results

Model Evaluation

For this assessment, we present last year's model updated for 2015 with no model changes. A comparison of the model likelihood components and key parameter estimates from 2014 are compared with the 2015 updated model.

Box 2: Model comparison of the 2014 (M0) and 2015 (M1) models by contribution to the objective function (negative log-likelihood values) and key parameters.

Model	<u>M0</u> 2014	<u>M1</u> 2015
Likelihood Components (Data)		
Catch	7	6
Domestic LL survey RPN	47	49
Japanese LL survey RPN	18	18
Domestic LL fishery RPW	10	10
Japanese LL fishery RPW	13	13
NMFS GOA trawl survey	19	22
Domestic LL survey ages	180	192
Domestic LL fishery ages	238	264
Domestic LL survey lengths	59	64
Japanese LL survey ages	144	144
Japanese LL survey lengths	46	46
NMFS trawl survey lengths	286	314
Domestic LL fishery lengths	207	211
Domestic trawl fishery lengths	194	204
Data likelihood	1469	1559
Total objective function value	1489	1579
Key parameters		
Number of parameters	219	222
$B_{next\ year}$ (Female spawning (kt) biomass for next year)	92	86
$B_{40\%}$ (Female spawning biomass (kt))	105	103
B_{1960} (Female spawning biomass (kt))	161	174
$B_{0\%}$ (Female spawning biomass (kt))	262	257
$SPR\%$ current	35.1%	33.6%
$F_{40\%}$	0.094	0.094
$F_{40\%}$ (Tier 3b adjusted)	0.082	0.078
$ABC(kt)$	13.7	11.8
$q_{Domestic\ LL\ survey}$	7.6	7.6
$q_{Japanese\ LL\ survey}$	6.2	6.2
$q_{Domestic\ LL\ fishery}$	4.0	4.0
$q_{Trawl\ Survey}$	1.3	1.3
$a_{50\%}$ (domestic LL survey selectivity)	3.8	3.8
$a_{50\%}$ (LL fishery selectivity)	3.9	3.9
μ_r (average recruitment)	16.7	16.3
σ_r (recruitment variability)	1.20	1.20

The two models are identical in all aspects except for inclusion of new data. Our usual criteria for choosing a superior model are: (1) the best overall fit to the data (in terms of negative log-likelihood), (2) biologically reasonable patterns of estimated recruitment, catchabilities, and selectivities, (3) a good visual fit to length and age compositions, and (4) parsimony.

Because the models presented have different amounts of data and different data weightings, it is not reasonable to compare their negative log likelihoods so we cannot compare them by the first criterion above. In general we can only evaluate the 2015 model based on changes in results from 2014 and it is unlikely we would reject the model that included the most recent data. The model generally produces good visual fits to the data, and biologically reasonable patterns of recruitment, abundance, and selectivities. An exception to the generally good fits to the data is the fit to the recent fishery age compositions, which fit the plus group poorly (see further discussion in *Goodness of fit* below). The 2015 update shows a slight decrease in spawning and total biomass from previous projections. Therefore the 2015 (M1) model is utilizing the new information effectively, and we use it to recommend 2016 ABC and OFL.

Time Series Results

Definitions

Spawning biomass is the biomass estimate of mature females. Total biomass is the estimate of all sablefish age-two and greater. Recruitment is measured as the number of age-two sablefish. Fishing mortality is fully-selected F, meaning the mortality at the age the fishery has fully selected the fish.

Abundance trends

Sablefish abundance increased during the mid-1960's (Table 3.15, Figure 3.13) due to strong year classes in the early 1960's. Abundance subsequently dropped during the 1970's due to heavy fishing and relatively low recruitment; catches peaked at 53,080 t in 1972. The population recovered due to a series of strong year classes from the late 1970's (Figure 3.14, Table 3.14) and also recovered at different rates in different areas (Table 3.15); spawning abundance peaked again in 1987. The population then decreased because these strong year classes expired. The model suggested an increasing trend in spawning biomass since the all-time low in 2002, which changed directions again in 2008 (Figure 3.13). The low 2012-2013 longline survey RPN values changed what was a stable trend in 2011 to a downward trajectory in 2015.

Projected 2016 spawning biomass is 34% of unfished spawning biomass. Spawning biomass has increased from a low of 33% of unfished biomass in 2002 to 42% in 2008 and has now declined back to 34% of unfished biomass projected for 2016. The 1997 year class has been an important contributor to the population; however, it has been reduced and is predicted to comprise less than 6% of the 2016 spawning biomass. The last two above-average year classes, 2000 and 2008, classes each comprise 15% of the projected 2016 spawning biomass. The 2008 year class will be about 75% mature in 2016. Figure 3.15 shows the relative contribution of each year class to next year's spawning biomass.

Recruitment trends

Annual estimated recruitment varies widely (Figure 3.14b). The two recent strong year classes in 1997 and 2000 are evident in all data sources. After 2000, few strong year classes are apparent, but the 2008 year class is currently estimated to be the largest since 2000. Few small fish were caught in the 2005 through 2009 trawl surveys, but the 2008 year class appeared in the 2011 trawl survey length composition. Larger one year olds may be showing up in the 2015 trawl survey length composition in the 41-43 cm bins (Figures 3.16, 3.17). The 2010 and 2011 longline survey age compositions show the 2008 year class appearing relatively strong in all three areas for lightly selected 2 and 3 year old fish (Figures 3.18-3.20). The 2014 survey age composition is dominated by 2007-2008 year classes which make up more than 30% of the composition. Large year classes often appear in the western areas first and then in subsequent years in the Central and Eastern GOA. While this was true for the 1997 and 2000 year classes,

the 2008 year class is appearing in all areas at approximately the same magnitude at the same time (Figure 3.18).

Average recruitment during 1979-2015 was 16.3 million 2-year-old sablefish per year, which is slightly less than average recruitment during 1958-2015. Estimates of recruitment strength during the 1960s are less certain because they depend on age data from the 1980s with older aged fish that are subject to more ageing error. In addition the size of the early recruitments is based on an abundance index during the 1960s based only on the Japanese fishery catch rate, which may be a weak measure of abundance. The 2008 year class is being estimated at just above average in this year's model. Because of the very low survey abundance indices in 2012, 2013, and 2015, the 2008 year class thus far is only just above average. If the 2008 year class is actually strong, the estimate may increase if the survey abundance estimates become stronger in future years.

Juvenile sablefish are pelagic and at least part of the population inhabits shallow near-shore areas for their first one to two years of life (Rutecki and Varosi 1997). In most years, juveniles have been found only in a few places such as Saint John Baptist Bay near Sitka, Alaska. Widespread, abundant age-1 juveniles likely indicate a strong year class. Abundant age-1 juveniles were reported for the 1960 (J. Fujioka & H. Zenger, 1995, NOAA, pers. comm.), 1977 (Bracken 1983), 1980, 1984, and 1998 year classes in southeast Alaska, the 1997 and 1998 year classes in Prince William Sound (W. Bechtol, 2004, ADFG, pers. comm.), the 1998 year class near Kodiak Island (D. Jackson, 2004, ADFG, pers. comm.), and the 2008 year class in Uganik Bay on Kodiak Island (P. Rigby, June, 2009, NOAA, pers. comm.). Numerous reports of young of the year being caught in 2014 have been received including large catches in NOAA surface trawl surveys in the EGOA in the summer (W. Fournier, August, 2014, NOAA, pers. comm.) and in Alaska Department of Fish and Game surveys in Prince William Sound (M. Byerly, 2014, ADFG, pers. comm.). Additionally, salmon fishermen in the EGOA reported large quantities of YOY sablefish in the stomachs of troll caught coho salmon in 2014 and 2015. The Gulf of Alaska NMFS bottom trawl survey caught a substantial number of one year old sablefish in 2015, particularly in the Western GOA. These size fish are not large enough or old enough to enter the model estimates. Surface trawl surveys in the Gulf of Alaska also reported finding YOY sablefish in Pacific pomfret stomachs in the summer of 2015 (C. Denenbaum, September 2015, NOAA, pers. comm.). Charter fishermen in the CGOA also reported frequent catches of one year old sablefish in 2015 while targeting coho salmon (K. Echave, September 2015, NOAA, pers. comm.).

Sablefish recruitment varies greatly from year to year (Figure 3.14b), but shows some relationship to environmental conditions. Sablefish recruitment success is related to winter current direction and water temperature; above average recruitment is more common for years with northerly drift or above average sea surface temperature (Sigler et al. 2001). Sablefish recruitment success is also coincidental with recruitment success of other groundfish species. Strong year classes were synchronous for many northeast Pacific groundfish stocks for the 1961, 1970, 1977, and 1984 year classes (Hollowed and Wooster 1992). For sablefish in Alaska, the 1960-1961 and 1977 year classes also were strong. Some of the largest year classes of sablefish occurred when abundance was near the historic low, the 1977-1978 and 1980-1981 year classes (Figures 3.14, 3.21). These strong year classes followed the 1976/1977 North Pacific regime shift. The 1977 year class was associated with the Pacific Decadal Oscillation (PDO) phase change and the 1977 and 1981 year classes were associated with warm water and unusually strong northeast Pacific pressure index (Hollowed and Wooster 1992). Larger than average year classes were produced again in 1997-2000, when the population was low indicating that recruitment is only weakly related to spawning biomass. Some species such as walleye pollock and sablefish may exhibit increased production at the beginning of a new environmental regime, when bottom up forcing prevails and high turnover species compete for dominance, which later shifts to top down forcing once dominance is established (Bailey 2000, Hunt et al. 2002). The large year classes of sablefish indicate that the population, though low, still was able to take advantage of favorable environmental conditions and produce large year classes. Shotwell et al. (2014) used a two-stage model selection process to examine relevant environmental variables that affect recruitment and included them directly into the assessment model. The best model

suggested that colder than average wintertime sea surface temperatures in the central North Pacific represent oceanic conditions that create positive recruitment events for sablefish in their early life history.

Goodness of fit

The model generally fit the data well. Abundance indices generally track through the middle of the confidence intervals of the estimates (Figures 3.3, 3.4), with the exception of the trawl survey, where predictions are typically lower in the early years and higher in later years. This index is given less weight than the other indices based on higher sampling error so it does not fit as well. Like the trawl survey index, the fishery CPUE does not fit as well as the longline survey, because the index has a higher variance. All age compositions were predicted well, except for not quite reaching the magnitude of the 1997 and 2000 year classes in several years (Figures 3.19, 3.21, 3.24). The model is not fitting the 2008 year class well in 2014 because of its weak presence in the 2013 age composition. The length frequencies from the fixed gear fishery are predicted well in most years, but the model appears to not fit the smallest fish that appear in 2011 (Figure 3.22, 3.23). The fits to the trawl survey and trawl fishery length compositions were generally mediocre, because of the small sample sizes relative to the longline survey and fishery length compositions (Figures 3.16, 3.17., 3.25). The model fit the domestic longline survey lengths poorly in the 1990s, then fit well until 2011 and 2012 where the smallest and largest fish were not fit well (Figures 3.26, 3.27). By 2014, the 2008 year class has grown large enough (in length) to be included in the main groups in the length compositions. The 2013 fixed gear fishery age composition is fit poorly, particularly in the plus group. This was due to an exceptionally high proportion of the catch caught in the AI being older than 30 years old. Examination of the origin of these older fish showed that this shift in fishery age composition was caused by a westward shift of the observed fishery into grounds that are not surveyed by the longline survey where there is an apparent abundance of older fish that are unknown to the model. This problem is similar, but lessened in the 2014 age composition. We will explore methods to consider these shifts in future spatial assessment models.

Selectivities

We assume that selectivity is asymptotic for the longline survey and fisheries and dome-shaped (or descending right limb) for the trawl survey and trawl fishery (Figure 3.28). The age-of-50% selection is 3.8 years for females in the longline survey and 3.9 years in the IFQ longline fishery. Females are selected at an older age in the IFQ fishery than in the derby fishery (Figure 3.28). Males were selected at an older age than females in both the derby and IFQ fisheries, likely because they are smaller at the same age. Selection of younger fish during short open-access seasons likely was due to crowding of the fishing grounds, so that some fishers were pushed to fish shallower water that young fish inhabit (Sigler and Lunsford 2001). Relative to the longline survey, younger fish are more vulnerable and older fish are less vulnerable to the trawl fishery because trawling often occurs on the continental shelf in shallower waters (< 300 m) where young sablefish reside. The trawl fishery selectivities are similar for males and females (Figure 3.28). The trawl survey selectivity curves differ between males and females, where males stay selected by the trawl survey longer (Figure 3.28). These trawl survey patterns are consistent with the idea that sablefish move out on the shelf at 2 years of age and then gradually become less available to the trawl fishery and survey as they move offshore into deeper waters.

Fishing mortality and management path

Fishing mortality was estimated to be high in the 1970s, relatively low in the early 1980s and then increased and held relatively steady in the 1990s and 2000s (Figure 3.29). Goodman et al. (2002) suggested that stock assessment authors use a “management path” graph as a way to evaluate management and assessment performance over time. In this “management path” we plot estimated fishing mortality relative to the (current) limit value and the estimated spawning biomass relative to limit spawning biomass ($B_{35\%}$). Figure 3.30 shows that recent management has generally constrained fishing

mortality below the limit rate, and until recently kept the stock above the $B_{35\%}$ limit. Projected 2015 and 2016 and 2017 spawning biomass is slightly below $B_{35\%}$.

Uncertainty

We compared a selection of parameter estimates from the Markov-Chain Monte Carlo (MCMC) simulations with the maximum-likelihood estimates, and compared each method's associated level of uncertainty (Table 3.16). Mean and median catchability estimates were nearly identical. The estimate of $F_{40\%}$ was lower by maximum likelihood and shows some skewness as indicated by the difference between the MCMC mean and median values. MCMC standard deviations were generally slightly higher in all cases which shows that there is more uncertainty captured through MCMC.

Retrospective analysis

Retrospective analysis is the examination of the consistency among successive estimates of the same parameters obtained as new data are added to a model. Retrospective analysis has been applied most commonly to age-structured assessments. Retrospective biases can arise for many reasons, ranging from bias in the data (e.g., catch misreporting, non-random sampling) to different types of model misspecification such as wrong values of natural mortality, or temporal trends in values set to be invariant. Classical retrospective analysis involves starting from some time period earlier in the model and successively adding data and testing if there is a consistent bias in the outputs (NRC 1998).

For this assessment, we show the retrospective trend in spawning biomass and total biomass for ten previous assessment years (2005-2014) compared to estimates from the current preferred model. This analysis is simply removing all new data that have been added for each consecutive year to the preferred model. Each year of the assessment generally adds one year of longline fishery lengths, trawl fishery lengths, longline survey lengths, longline and fishery ages (from one year prior), fishery abundance index, and longline survey index. Every other year, a trawl survey estimate and corresponding length composition are added.

In the first four years of the retrospective plot we see that estimates of spawning biomass were consistently lower for the last few years in the next assessment year (Figure 3.31a). In recent years, the retrospective plot of spawning biomass shows only small changes from year to year (e.g., Table 3.17). One common measure of the retrospective bias is Mohn's revised rho which indicates the size and direction of the bias. The revised Mohn's rho of 0.023 is very low (a small positive retrospective bias) relative to most assessments at the AFSC (Hanselman et al. 2013). The retrospective patterns are well within the posterior uncertainty of each assessment (Figure 3.31b). Recruitment estimates appear to have little trend over time with the exception of the 2002 year class which increased from a very low value to near average (Figure 3.31c). Only the 2008 year class started near average indicating low presence of 2 year olds in most of the recent data.

Examining retrospective trends can show potential biases in the model, but may not identify what their source is. Other times a retrospective trend is merely a matter of the model having too much inertia in the age-structure and other historic data to respond to the most recent data. This retrospective pattern likely to be considered mild, but at issue is the "one-way" pattern in the early part of the retrospective time series. It is difficult to isolate the cause of this pattern but several possibilities exist. For example, hypotheses could include environmental changes in catchability, time-varying natural mortality, or changes in selectivity of the fishery or survey. One other issue is that fishery abundance and lengths, and all age compositions are added into the assessment with a one year lag to the current assessment. This estimate of rho is down from 0.089 in 2013 and about the same as in 2014 (0.019), which we attribute to two factors: 1) 2003 was dropped out of the retrospective window which had a relatively large change from the terminal year; and 2) The updated catch data that was used in 2014 added a significant amount of catch in the early part of the retrospective window, which increased the estimate of spawning biomass at the

recent low point. We will monitor and explore these patterns in the future.

The 2010 Joint Plan Team requested that we examine what the current model configuration would have recommended for ABCs going back in time to see how much model and author changes has affected management advice. We examined this in the 2011 SAFE and concluded that despite many model changes, including growth updates and a split-gender model, the management advice would have been similar (Hanselman et al. 2011).

Harvest Recommendations

Reference fishing mortality rate

Sablefish are managed under Tier 3 of NPFMC harvest rules. Reference points are calculated using recruitments from 1977-2012. The updated point estimates of $B_{40\%}$, $F_{40\%}$, and $F_{35\%}$ from this assessment are 102,807 t (combined across the EBS, AI, and GOA), 0.094, and 0.112, respectively. Projected female spawning biomass (combined areas) for 2016 is 86,471 t (84% of $B_{40\%}$), placing sablefish in sub-tier “b” of Tier 3. The maximum permissible value of F_{ABC} under Tier 3b is 0.078, which translates into a 2016 ABC (combined areas) of 11,795 t. The OFL fishing mortality rate is 0.093 which translates into a 2016 OFL (combined areas) of 13,397 t. If the stock were in Tier 3a (above the $B_{40\%}$ reference point), the 2016 ABC would be 14,164 t. Model projections indicate that this stock is not subject to overfishing, overfished, nor approaching an overfished condition.

Population projections

A standard set of projections is required for each stock managed under Tiers 1, 2, or 3 of Amendment 56. This set of projections encompasses seven harvest scenarios designed to satisfy the requirements of Amendment 56, the National Environmental Policy Act, and the Magnuson-Stevens Fishery Conservation and Management Act (MSFCMA).

For each scenario, the projections begin with the vector of 2015 numbers at age as estimated in the assessment. This vector is then projected forward to the beginning of 2016 using the schedules of natural mortality and selectivity described in the assessment and the best available estimate of total (year-end) catch for 2015. In each subsequent year, the fishing mortality rate is prescribed on the basis of the spawning biomass in that year and the respective harvest scenario. In each year, recruitment is drawn from an inverse Gaussian distribution whose parameters consist of maximum likelihood estimates determined from recruitments estimated in the assessment. Spawning biomass is computed in each year based on the time of peak spawning and the maturity and weight schedules described in the assessment. Total catch after 2015 is assumed to equal the catch associated with the respective harvest scenario in all years. This projection scheme is run 1,000 times to obtain distributions of possible future stock sizes, fishing mortality rates, and catches.

Five of the seven standard scenarios will be used in an Environmental Assessment prepared in conjunction with the final SAFE. These five scenarios, which are designed to provide a range of harvest alternatives that are likely to bracket the final TAC for 2016, are as follow (“ $max F_{ABC}$ ” refers to the maximum permissible value of F_{ABC} under Amendment 56):

Scenario 1: In all future years, F is set equal to $max F_{ABC}$. (Rationale: Historically, TAC has been constrained by ABC, so this scenario provides a likely upper limit on future TACs.)

Scenario 2: In 2016 and 2017, F is set equal to a constant fraction of $max F_{ABC}$, where this fraction is equal to the ratio of the realized catches in 2012-2014 to the TAC for each of those years. For the remainder of the future years, maximum permissible ABC is used. (Rationale: In many fisheries the ABC is routinely not fully utilized, so assuming an average ratio of F will yield more realistic projections.)

Scenario 3: In all future years, F is set equal to 50% of $max F_{ABC}$. (Rationale: This scenario

provides a likely lower bound on F_{ABC} that still allows future harvest rates to be adjusted downward when stocks fall below reference levels.)

Scenario 4: In all future years, F is set equal to the 2010-2014 average F . (Rationale: For some stocks, TAC can be well below ABC, and recent average F may provide a better indicator of F_{TAC} than F_{ABC} .)

Scenario 5: In all future years, F is set equal to zero. (Rationale: In extreme cases, TAC may be set at a level close to zero.)

Two other scenarios are needed to satisfy the MSFCMA's requirement to determine whether a stock is currently in an overfished condition or is approaching an overfished condition. These two scenarios are as follows (for Tier 3 stocks, the MSY level is defined as $B_{35\%}$):

Scenario 6: In all future years, F is set equal to F_{OFL} . (Rationale: This scenario determines whether a stock is overfished. If the stock is expected to be 1) above its MSY level in 2015 or 2) above $\frac{1}{2}$ of its MSY level in 2015 and above its MSY level in 2025 under this scenario, then the stock is not overfished.)

Scenario 7: In 2016 and 2017, F is set equal to $\max F_{ABC}$, and in all subsequent years F is set equal to F_{OFL} . (Rationale: This scenario determines whether a stock is approaching an overfished condition. If the stock is 1) above its MSY level in 2017 or 2) above $\frac{1}{2}$ of its MSY level in 2017 and expected to be above its MSY level in 2027 under this scenario, then the stock is not approaching an overfished condition.)

Spawning biomass, fishing mortality, and yield are tabulated for the seven standard projection scenarios (Table 3.18). The difference for this assessment for projections is in Scenario 2 (Author's F); we use pre-specified catches to increase accuracy of short-term projections in fisheries (such as sablefish) where the catch is usually less than the ABC. This was suggested to help management with setting more accurate preliminary ABCs and OFLs for 2016 and 2017. The methodology for determining these pre-specified catches is described below in ***Specified catch estimation***.

Status determination

In addition to the seven standard harvest scenarios, Amendments 48/48 to the BSAI and GOA Groundfish Fishery Management Plans require projections of the likely OFL two years into the future. While Scenario 6 gives the best estimate of OFL for 2016, it does not provide the best estimate of OFL for 2017, because the mean 2016 catch under Scenario 6 is predicated on the 2016 catch being equal to the 2016 OFL, whereas the actual 2016 catch will likely be less than the 2016 OFL. A better approach is to estimate catches that are more likely to occur as described below under *Specified Catch Estimation*. The executive summary contains the appropriate one- and two-year ahead projections for both ABC and OFL.

Under the MSFCMA, the Secretary of Commerce is required to report on the status of each U.S. fishery with respect to overfishing. This report involves the answers to three questions: 1) Is the stock being subjected to overfishing? 2) Is the stock currently overfished? 3) Is the stock approaching an overfished condition?

Is the stock being subjected to overfishing? The official catch estimate for the most recent complete year (2014) is 11,582 t. This is less than the 2014 OFL of 16,160 t. Therefore, the stock is not being subjected to overfishing.

Harvest Scenarios #6 and #7 (Table 3.18) are intended to permit determination of the status of a stock with respect to its minimum stock size threshold (MSST). Any stock that is below its MSST is defined to be *overfished*. Any stock that is expected to fall below its MSST in the next two years is defined to be

approaching an overfished condition. Harvest Scenarios #6 and #7 are used in these determinations as follows:

Is the stock currently overfished? This depends on the stock's estimated spawning biomass in 2015:

- a. If spawning biomass for 2015 is estimated to be below $\frac{1}{2} B_{35\%}$, the stock is below its MSST.
- b. If spawning biomass for 2015 is estimated to be above $B_{35\%}$, the stock is above its MSST.
- c. If spawning biomass for 2015 is estimated to be above $\frac{1}{2} B_{35\%}$ but below $B_{35\%}$, the stock's status relative to MSST is determined by referring to harvest Scenario #6 (Table 3.18). If the mean spawning biomass for 2025 is below $B_{35\%}$, the stock is below its MSST. Otherwise, the stock is above its MSST.

Is the stock approaching an overfished condition? This is determined by referring to harvest Scenario #7 (Table 3.18):

- a. If the mean spawning biomass for 2017 is below $\frac{1}{2} B_{35\%}$, the stock is approaching an overfished condition.
- b. If the mean spawning biomass for 2017 is above $B_{35\%}$, the stock is not approaching an overfished condition.
- c. If the mean spawning biomass for 2017 is above $\frac{1}{2} B_{35\%}$ but below $B_{35\%}$, the determination depends on the mean spawning biomass for 2027. If the mean spawning biomass for 2027 is below $B_{35\%}$, the stock is approaching an overfished condition. Otherwise, the stock is not approaching an overfished condition.

Based on the above criteria and the results of the seven scenarios in Table 3.18, the stock is not overfished and is not approaching an overfished condition.

Specified catch estimation

In response to GOA Plan Team minutes in 2010, we have established a consistent methodology for estimating current-year and future year catches in order to provide more accurate two-year projections of ABC and OFL to management. We explained the methods and gave examples in the 2011 SAFE (Hanselman et al. 2011). Going forward, for current year catch, we are applying an expansion factor to the official catch on or near October 1 by the 3-year average of catch taken between October 1 and December 31 in the last three complete catch years (e.g. 2012-2014 for this year).

For catch projections into the next two years, we are using the ratio of the last three official catches to the last three TACs multiplied against the future two years' ABCs (if TAC is normally the same as ABC). This method results in slightly higher ABCs in each of the future two years of the projection, based on both the lower catch in the first year out, and on the amount of catch taken before spawning in the projection two years out.

Bayesian analysis

The model estimates of projected spawning biomass fall near the center of the posterior distribution of spawning biomass. Most of the probability lies between 80,000 and 100,000 t (Figure 3.32). The probability changes smoothly and exhibits a relatively normal distribution. The posterior distribution clearly indicates the stock is below $B_{40\%}$.

Scatter plots of selected pairs of model parameters were produced to evaluate the shape of the posterior distribution (Figure 3.33). The plots indicate that the parameters are reasonably well defined by the data. As expected, catchabilities, $F_{40\%}$, and ending spawning biomass were confounded. The catchability of the longline survey is most confounded with ending spawning biomass because it has the most influence in the model in recent abundance predictions.

We estimated the posterior probability that projected abundance will fall, or stay below thresholds of 17.5% (MSST), and 35% (MSY), and 40% (B_{target}) of the unfished spawning biomass based on the posterior probability estimates. Abundance was projected for 14 years. For management, it is important to know the risk of falling under these thresholds. The probability that spawning biomass falls below key biological reference points was estimated based on the posterior probability distribution for spawning biomass. The probability that next year's spawning biomass was below $B_{35\%}$ was 0.89. During the next three years, the probability of being below $B_{17.5\%}$ is near zero, the probability of being below $B_{35\%}$ is 0.98, and the probability of staying below $B_{40\%}$ is near 100% (Figure 3.34).

Alternative Projection

We also use an alternative projection that considers uncertainty from the whole model by running projections within the model. This projection propagates uncertainty throughout the entire assessment procedure and is based on 10,000,000 MCMC (burnt-in and thinned) using the standard Tier 3 harvest rules. The projection shows wide credible intervals on future spawning biomass (Figure 3.35). The $B_{35\%}$ and $B_{40\%}$ reference points are based on the 1979-2013 recruitments, and this projection predicts that the mean and median spawning biomass will stay below $B_{35\%}$ until after 2020, and then return to $B_{40\%}$ if average recruitment is attained. This projection is run with the same ratio for catch as described in Alternative 2 above, except for all future years instead of the next two.

Acceptable biological catch

We recommend a 2016 ABC of 11,795 t. The maximum permissible ABC for 2016 based on Tier 3b of the harvest control rule, uses an adjusted $F_{40\%}$ which yields 11,795 t. The maximum permissible ABC for 2016 is 14% lower than the 2015 ABC of 13,657 t. The 2014 assessment projected a 10% decrease in ABC for 2016 from 2015. This slightly larger decrease is supported by a new low in the domestic longline survey index time series that offset the small increases in the fishery abundance index seen in 2014 and the Gulf of Alaska trawl survey index in 2015. The fishery abundance index has been trending down since 2007. The 2014 IPHC GOA sablefish index was not used in the model, but was similar and trending low in 2013 and 2014. The 2008 year class showed potential to be large in previous assessments based on patterns in the age and length compositions. However the estimate in this year's assessment is only just above average because the recent large overall decrease in the longline survey and trawl indices have lowered the overall scale of the population. Spawning biomass is projected to decline through 2018, and then is expected to increase assuming average recruitment is achieved in the future. ABCs are projected to decrease in 2017 to 10,782 t and 10,869 t in 2018 (see Table 3.18).

Area allocation of harvests

The combined ABC has been apportioned to regions using weighted moving average methods since 1993; these methods reduce the magnitude of inter-annual changes in the apportionment. Weighted moving average methods are robust to uncertainties about movement rates and measurement error of the biomass distribution, while adapting to current information about the biomass distribution. The 1993 TAC was apportioned using a 5 year running average with emphasis doubled for the current year survey abundance index in weight (relative population weight or RPW). Since 1995, the ABC was apportioned using an exponential weighting of regional RPWs. Exponential weighting is implied under certain conditions by the Kalman filter. The exponential factor is the measurement error variance divided by the prediction error variance (Meinhold and Singpurwalla 1983). Prediction error variance depends on the variances of the previous year's estimate, the process error, and the measurement error. When the ratio of measurement error variance to process error variance is r , the exponential factor is equal to

$1 - 2/(\sqrt{4r+1} + 1)$ (Thompson 2004). For sablefish we do not estimate these values, but instead set the

exponential factor at $\frac{1}{2}$, so that, except for the first year, the weight of each year's value is $\frac{1}{2}$ the weight of the following year. The weights are year index 5: 0.0625; 4: 0.0625; 3: 0.1250; 2: 0.2500; 1: 0.5000. A $(\frac{1}{2})^x$ weighting scheme, where x is the year index, reduced annual fluctuations in regional ABC, while keeping regional fishing rates from exceeding overfishing levels in a stochastic migratory model (J. Heifetz, 1999, NOAA, pers. comm.). Because mixing rates for sablefish are sufficiently high and fishing rates sufficiently low, moderate variations of biomass-based apportionment would not significantly change overall sablefish yield unless there are strong differences in recruitment, growth, and survival by area (Heifetz et al. 1997).

Previously, the Council approved apportionments of the ABC based on survey data alone. Starting with the 2000 ABC, the Council approved an apportionment based on survey and fishery data. The fishery and survey information were combined to apportion ABC using the following method: The RPWs based on the fishery data were weighted with the same exponential weights used to weight the survey data (year index 5: 0.0625; 4: 0.0625; 3: 0.1250; 2: 0.2500; 1: 0.5000). The fishery and survey data were combined by computing a weighted average of the survey and fishery estimates, with the weight inversely proportional to the variability of each data source. The variance for the fishery data has typically been twice that of the survey data, so the survey data was weighted twice as much as the fishery data. Below are area-specific apportionments following the traditional apportionment scheme, which we are **not recommending for 2016:**

Apportionments are based on survey and fishery information	2014 ABC Percent	2015 Survey RPW	2014 Fishery RPW	2016 ABC Percent	2015 ABC	2016 ABC	Change
Total					13,657	11,795	-14%
Bering Sea	10%	15%	16%	10%	1,333	1,816	36%
Aleutians	13%	13%	15%	13%	1,802	1,627	-10%
Gulf of Alaska	77%	72%	69%	77%	10,522	8,352	-21%
Western	14%	14%	13%	14%	1,473	1,136	-23%
Central	44%	43%	36%	44%	4,658	3,451	-26%
W. Yakutat*	15%	16%	17%	15%	1,567	1,374	-12%
E. Yakutat / Southeast*	27%	27%	34%	27%	2,823	2,391	-15%

Following the standard apportionment scheme, we have observed that the objective to reduce variability in apportionment was not being achieved. Since 2007, the mean change in apportionment by area has increased annually (Figure 3.36A). While some of these changes may actually reflect interannual changes in regional abundance, they most likely reflect the high movement rates of the population and the high variability of our estimates of abundance in the Bering Sea and Aleutian Islands. For example, the apportionment for the Bering Sea has varied drastically since 2007, attributable to high variability in both survey abundance and fishery CPUE estimates in the Bering Sea (Figure 3.36B). These large annual changes in apportionment result in increased variability of ABCs by area, including areas other than the Bering Sea (Figure 3.36C). Because of the high variability in apportionment seen in recent years, we do not believe the standard method is meeting the goal of reducing the magnitude of interannual changes in the apportionment. Because of these reasons, we recommended fixing the apportionment at the proportions from the 2013 assessment, until the apportionment scheme is thoroughly reevaluated and reviewed. A Ph.D. student with the University of Alaska-Fairbanks began a project in 2013 with the objectives of re-examining the apportionment strategy and conducting a management strategy evaluation. A spatial sablefish model has been developed, but the management strategy evaluation is in early stages of development. Meanwhile, it seems imprudent to move to an interim apportionment or return to the former scheme until more satisfactory methods have been identified and evaluated. **Therefore, for 2016, we recommend continuing with the apportionment fixed at the proportions used in 2015.**

Area	2015 ABC	Standard apportionment for 2016 ABC	Recommended fixed apportionment for 2016 ABC*	Difference from 2015
Total	13,657	11,795	11,795	-13.6%
Bering Sea	1,333	1,816	1,151	-13.6%
Aleutians	1,802	1,627	1,557	-13.6%
Gulf of Alaska (subtotal)	10,522	8,352	9,087	-13.6%
Western	1,473	1,136	1,272	-13.6%
Central	4,658	3,451	4,023	-13.6%
W. Yakutat**	1,567	1,374	1,353	-13.6%
E. Yak. / Southeast**	2,823	2,391	2,438	-13.6%

* Fixed at the 2013 assessment apportionment proportions (Hanselman et al. 2012). ** Before 95:5 hook and line: trawl split shown below.

Adjusted for 95:5 hook-and-line: trawl split in EGOA	Year	W. Yakutat	E. Yakutat/Southeast
	2016	1,475 t	2,316 t
	2017	1,348 t	2,118 t

Overfishing level (OFL)

Applying an adjusted $F_{35\%}$ as prescribed for OFL in Tier 3b, results in a value of 13,397 t for the combined stock. The OFL is apportioned by region, Bering Sea (1,304 t), AI (1,766 t), and GOA (10,326 t), by the same method as the ABC apportionment.

Ecosystem considerations

Ecosystem considerations for the Alaska sablefish fishery are summarized in Table 3.19.

Ecosystem effects on the stock

Prey population trends

Young-of-the-year sablefish prey mostly on euphausiids (Sigler et al. 2001) and copepods (Grover and Olla 1990), while juvenile and adult sablefish are opportunistic feeders. Larval sablefish abundance has been linked to copepod abundance and young-of-the-year abundance may be similarly affected by euphausiid abundance because of their apparent dependence on a single species (McFarlane and Beamish 1992). The dependence of larval and young-of-the-year sablefish on a single prey species may be the cause of the observed wide variation in annual sablefish recruitment. No time series is available for copepod and euphausiid abundance, so predictions of sablefish abundance based on this predator-prey relationship are not possible.

Juvenile and adult sablefish feed opportunistically, so diets differ throughout their range. In general, sablefish < 60 cm consume more euphausiids, shrimp, and cephalopods, while sablefish > 60 cm consume more fish (Yang and Nelson 2000). In the GOA, fish constituted 3/4 of the stomach content weight of adult sablefish with the remainder being invertebrates (Yang and Nelson 2000). Of the fish found in the diets of adult sablefish, pollock were the most abundant item while eulachon, capelin, Pacific herring, Pacific cod, Pacific sand lance, and flatfish also were found. Squid were the most important invertebrate and euphausiids and jellyfish were also present. In southeast Alaska, juvenile sablefish also consume juvenile salmon at least during the summer months (Sturdevant et al. 2009). Off the coast of Oregon and California, fish made up 76 percent of the diet (Laidig et al. 1997), while euphausiids dominated the diet off the southwest coast of Vancouver Island (Tanasichuk 1997). Off Vancouver Island, herring and other fish were increasingly important as sablefish size increased; however, the most important prey item was euphausiids. It is unlikely that juvenile and adult sablefish are affected by availability and abundance of

individual prey species because they are opportunistic feeders. The only likely way prey could affect growth or survival of juvenile and adult sablefish is by overall changes in ecosystem productivity.

Predators/Competitors: The main juvenile sablefish predators are adult coho and chinook salmon, which prey on young-of-the-year sablefish during their pelagic stage. Sablefish were the fourth most commonly reported prey species in the salmon troll logbook program from 1977 to 1984 (Wing 1985), however the effect of salmon predation on sablefish survival is unknown. The only other fish species reported to prey on sablefish in the GOA is Pacific halibut; however, sablefish comprised less than 1% of their stomach contents (M. Yang, October 14, 1999, NOAA, pers. comm.). Although juvenile sablefish may not be a prominent prey item because of their relatively low and sporadic abundance compared to other prey items, they share residence on the continental shelf with potential predators such as arrowtooth flounder, halibut, Pacific cod, bigmouth sculpin, big skate, and Bering skate, which are the main piscivorous groundfishes in the GOA (Yang et al. 2006). It seems possible that predation of sablefish by other fish is significant to the success of sablefish recruitment even though they are not a common prey item.

Sperm whales are likely a major predator of adult sablefish. Fish are an important part of sperm whale diet in some parts of the world, including the northeastern Pacific Ocean (Kawakami 1980). Fish have appeared in the diets of sperm whales in the eastern AI and GOA. Although fish species were not identified in sperm whale diets in Alaska, sablefish were found in 8.3% of sperm whale stomachs off of California (Kawakami 1980).

Sablefish distribution is typically thought to be on the upper continental slope in deeper waters than most groundfish. However, during the first two to three years of their life sablefish inhabit the continental shelf. Length samples from the NMFS bottom trawl survey suggest that the geographic range of juvenile sablefish on the shelf varies dramatically from year to year. In particular, juveniles utilize the Bering Sea shelf extensively in some years, while not at all in others (Shotwell et al. 2014). Juvenile sablefish (< 60 cm FL) prey items overlap with the diet of small arrowtooth flounder. On the continental shelf of the GOA, both species consumed euphausiids and shrimp predominantly; these prey are prominent in the diet of many other groundfish species as well. This diet overlap may cause competition for resources between small sablefish and other groundfish species.

Changes in the physical environment: Mass water movements and temperature changes appear related to recruitment success. Above-average recruitment was somewhat more likely with northerly winter currents and much less likely for years when the drift was southerly. Recruitment was above average in 61% of the years when temperature was above average, but was above average in only 25% of the years when temperature was below average. Growth rate of young-of-the-year sablefish is higher in years when recruitment is above average (Sigler et al. 2001). Shotwell et al. (2014) showed that colder than average wintertime sea surface temperatures in the central North Pacific may represent oceanic conditions that create positive recruitment events for sablefish in their early life history.

Anthropogenic changes in the physical environment: The Essential Fish Habitat Environmental Impact Statement (EFH EIS) (NMFS 2005) concluded that the effects of commercial fishing on the habitat of sablefish is minimal or temporary in the current fishery management regime primarily based on the criterion that sablefish are currently above Minimum Stock Size Threshold (MSST).

Juvenile sablefish are partly dependent on benthic prey (18% of diet by weight) and the availability of benthic prey may be adversely affected by fishing. Little is known about effects of fishing on benthic habitat or the habitat requirements for growth to maturity. Although sablefish do not appear to be directly dependent on physical structure, reduction of living structure is predicted in much of the area where juvenile sablefish reside and this may indirectly reduce juvenile survivorship by reducing prey availability or by altering the abilities of competing species to feed and avoid predation.

Fishery effects on the ecosystem

Fishery-specific contribution to bycatch of prohibited species, forage species, HAPC biota, marine mammals and birds, and other sensitive non-target species: The sablefish fishery catches significant portions of the shark and thornyhead rockfish total catch (Table 3.4). The sablefish fishery catches the majority of grenadier total catch; the annual amount is variable (Table 3.5). The trend in seabird catch is variable, but is substantially low compared to the 1990s, presumably due to widespread use of measures to reduce seabird catch. Prohibited species catches (PSC) in the targeted sablefish fisheries are dominated by halibut (334 t/year) and golden king crab (47,000 individuals/year). Halibut catches were average in 2015, while golden king crab catches have dropped precipitously from almost 200,000 individuals in 2011 to 3,193 t in 2014 and 11,270 t in 2015 (Table 3.6).

The shift from an open-access to an IFQ fishery has increased catching efficiency which has reduced the number of hooks deployed (Sigler and Lunsford 2001). Although the effects of longline gear on bottom habitat are poorly known, the reduced number of hooks deployed during the IFQ fishery must reduce the effects on benthic habitat. The IFQ fishery likely has also reduced discards of other species because of the slower pace of the fishery and the incentive to maximize value from the catch.

Fishery-specific concentration of target catch in space and time relative to predator needs in space and time (if known) and relative to spawning components: The sablefish fishery largely is dispersed in space and time. The longline fishery lasts 8-1/2 months. The quota is apportioned among six regions of Alaska.

Fishery-specific effects on amount of large size target fish: The longline fishery catches mostly medium and large-size fish which are typically mature. Length frequencies from the pot fishery in the BSAI are very similar to the longline fishery. The trawl fishery, which on average accounts for about 10% of the total catch, often catches slightly smaller fish. The trawl fishery typically occurs on the continental shelf where juvenile sablefish sometimes occur. Catching these fish as juveniles reduces the yield available from each recruit.

Fishery-specific contribution to discards and offal production: Discards of sablefish in the longline fishery are small, typically less than 5% of total catch (Table 3.3). The catch of sablefish in the longline fishery typically consists of a high proportion of sablefish, 90% or more. However, at times grenadiers may be a significant catch and they are almost always discarded.

Fishery-specific effects on age-at-maturity and fecundity of the target species: The shift from an open-access to an IFQ fishery has decreased harvest of immature fish and improved the chance that individual fish will reproduce at least once (Sigler and Lunsford 2001).

Fishery-specific effects on EFH non-living substrate: The primary fishery for sablefish is with longline gear. While it is possible that longlines could move small boulders it is unlikely fishing would persist where this would often occur. Relative to trawl gear, a significant effect of longlines on bedrock, cobbles, or sand is unlikely.

Data gaps and research priorities

There is little information on early life history of sablefish and recruitment processes. A better understanding of juvenile distribution, habitat utilization, and species interactions would improve understanding of the processes that determine the productivity of the stock. Better estimation of recruitment and year class strength would improve assessment and management of the sablefish population.

The Essential Fish Habitat 5-year Update will begin in 2016 and new habitat suitability models will be distributed on all life history stages for FMP species where data were available. Following evaluation of this information for sablefish, we plan to include the new EFH model results and maps in future assessments. A newly revamped species-specific ecosystem consideration (SEC) protocol is also planned

to be introduced over the next several years. The SECs will likely replace the Ecosystem Consideration section of the single-species assessment reports and include updated species profiles, climate vulnerability analyses, and stock/habitat prioritization information. The intention of these SECs is to improve the process of integrating ecosystem information into the stock assessments and facilitate ecosystem based fishery management.

Future sablefish research is going to focus on several directions:

- 1) Evaluating different apportionment strategies for the ABC.
- 2) Refine survey abundance index model for inclusion in future assessment model that accounts for whale depredation and potentially includes gully abundance data and other covariates.
- 3) Refine fishery abundance index to utilize a core fleet, and identify covariates that affect catch rates.
- 4) Improve knowledge of sperm whale and killer whale depredation in the fishery and begin to quantify depredation effects on fishery catch rates.
- 5) Continue to explore the use of environmental data to aid in determining recruitment.
- 6) An integrated GOA Ecosystem project funded by the North Pacific Research Board is underway and is looking at recruitment processes of major groundfish including sablefish. We hope to work closely with this project to help understand sablefish recruitment dynamics.
- 7) Develop a species-specific report card and enhanced ecosystem considerations section based on the the results of the GOA project described above.
- 8) We are developing a spatially explicit research assessment model that includes movement, which will help in examining smaller-scale population dynamics while retaining a single stock hypothesis Alaska-wide sablefish model. This is to include a management strategy evaluation of apportionment strategies.

Literature Cited

- Bailey, K. M. 2000. Shifting control of recruitment of walleye pollock *Theragra chalcogramma* after a major climatic and ecosystem change. *Marine Ecology Progress Series*, 198, 215–224.
- Bracken, B. 1983. Sablefish migration in the Gulf of Alaska based on tag recoveries. *Proceedings of the International Sablefish Symposium*. Alaska Sea Grant Report 83-8.
- Coutré, K. M., A.H. Beaudreau, and P.W. Malecha. 2015. Temporal Variation in Diet Composition and Use of Pulsed Resource Subsidies by Juvenile Sablefish. *Transactions of the American Fisheries Society*, 144(4), 807-819.
- Crecco, V. and W. J. Overholtz. 1990. Causes of density-dependent catchability for Georges Bank haddock *Melanogrammus aeglefinus*. *Can. J. Fish. Aquat. Sci.* 47: 385-394.
- Echave, K. B., D. H. Hanselman, M. D. Adkison, M. F. Sigler. 2012. Inter-decadal changes in sablefish, *Anoplopoma fimbria*, growth in the northeast Pacific Ocean. *Fish. Bull.* 210:361-374.
- Fina, M. 2011. Evolution of Catch Share Management: Lessons from Catch Share Management in the North Pacific. *Fisheries*. 36(4):164-177.
- Fournier, D. and C. P. Archibald. 1982. A general theory for analyzing catch at age data. *Can. J. Fish. Aquat. Sci.* 39: 1195-1207.
- Fournier, D.A., H.J. Skaug, J. Ancheta, J. Ianelli, A. Magnusson, M.N. Maunder, A. Nielsen, and J. Sibert. 2012. AD Model Builder: using automatic differentiation for statistical inference of highly parameterized complex nonlinear models. *Optim. Methods Softw.* 27, 233-249.
- Funk, F. and B. E. Bracken. 1984. Status of the Gulf of Alaska sablefish (*Anoplopoma fimbria*) resource in 1983. *Alaska Dept. Fish Game., Info. Leaflet*. 235, 55 p.
- Gao, Y., S.H. Jøner, R.A. Svec, and K.L. Weinberg. 2004. Stable isotopic comparison in otoliths of juvenile sablefish (*Anoplopoma fimbria*) from waters off the Washington and Oregon coast. *Fisheries Research*, 68(1), 351-360.
- Goodman, D., M. Mangel, G. Parkes, T.J. Quinn II, V. Restrepo, T. Smith, and K. Stokes. 2002. Scientific Review of the Harvest Strategy Currently Used in the BSAI and GOA Groundfish Fishery Management Plans. Draft report. North Pacific Fishery Management Council, 605 W 4th Ave, Suite 306 Anchorage, AK 99501.
- Grover, J. J. and B. L. Olla. 1990. Food habits of larval sablefish *Anoplopoma fimbria* from the Bering Sea. *Fish. Bull.* 88: 811-814.
- Hanselman, D.H., J. Heifetz, K.B. Echave, and S.C. Dressel. 2015. Move it or lose it: Movement and mortality of sablefish tagged in Alaska. *Canadian Journal of Fish and Aquatic Sciences*.
<http://www.nrcresearchpress.com/doi/abs/10.1139/cjfas-2014-0251>
- Hanselman, D.H., C. Lunsford, and C. Rodgveller. 2014. Assessment of the sablefish stock in Alaska. *In* Stock assessment and fishery evaluation report for the groundfish resources of the GOA and BS/AI. North Pacific Fishery Management Council, 605 W 4th Ave, Suite 306 Anchorage, AK 99501.
- Hanselman, D.H., W. Clark, and M. Sigler. 2013. Retrospective investigations group report.
http://www.afsc.noaa.gov/REFM/stocks/Plan_Team/2013/Sept/Retrospectives_2013_final3.pdf
- Hanselman, D.H., W. Clark, J. Heifetz, and D. Anderl. 2012. Statistical distribution of age readings of known-age sablefish (*Anoplopoma fimbria*). *Fish. Res.* 131: 1-8.
- Hanselman, D.H., C. Lunsford, and C. Rodgveller. 2012. Assessment of the sablefish stock in Alaska. *In* Stock assessment and fishery evaluation report for the groundfish resources of the GOA and BS/AI. North Pacific Fishery Management Council, 605 W 4th Ave, Suite 306 Anchorage, AK 99501. pp. 323-432.

- Hanselman, D.H., C. Lunsford, and C. Rodgveller. 2011. Assessment of the sablefish stock in Alaska. *In* Stock assessment and fishery evaluation report for the groundfish resources of the GOA and BS/AI. North Pacific Fishery Management Council, 605 W 4th Ave, Suite 306 Anchorage, AK 99501. pp. 307-412.
- Hanselman, D. H., C.R. Lunsford, and C. Rodgveller. 2010. Assessment of the sablefish stock in Alaska. *In* Stock assessment and fishery evaluation report for the groundfish resources of the GOA and BS/AI as projected for 2011. North Pacific Fishery Management Council, 605 W 4th Ave, Suite 306 Anchorage, AK 99501.
- Hanselman, D. H., J. Fujioka, C. Lunsford, and C. Rodgveller. 2009. Assessment of the sablefish stock in Alaska. *In* Stock assessment and fishery evaluation report for the groundfish resources of the GOA and BS/AI as projected for 2010. North Pacific Fishery Management Council, 605 W 4th Ave, Suite 306 Anchorage, AK 99501.
- Hanselman, D. H., C. Lunsford, J. Fujioka, and C. Rodgveller. 2008. Alaskan Sablefish. *In* Stock assessment and fishery evaluation report for the groundfish resources of the GOA and BS/AI as projected for 2008. North Pacific Fishery Management Council, 605 W 4th Ave, Suite 306 Anchorage, AK 99501.
- Hanselman, D. H., C. Lunsford, J. Fujioka, and C. Rodgveller. 2007. Alaskan Sablefish. *In* Stock assessment and fishery evaluation report for the groundfish resources of the GOA and BS/AI as projected for 2008. North Pacific Fishery Management Council, 605 W 4th Ave, Suite 306 Anchorage, AK 99501.
- Hanselman, D.H., J. Heifetz, J.T. Fujioka, and J.N. Ianelli. 2005a. Gulf of Alaska Pacific ocean perch. *In* Stock assessment and fishery evaluation report for the groundfish fisheries of the Gulf of Alaska. pp. 525-578. North Pacific Fishery Management Council, 605 W 4th Avenue, Suite 306, Anchorage, AK 99510.
- Hanselman, D.H., C.R. Lunsford, M.F. Sigler, and J.T. Fujioka. 2005b. Alaska sablefish assessment for 2006. *In* Stock assessment and fishery evaluation report for the groundfish fisheries of the Gulf of Alaska. North Pacific Fishery Management Council, 605 W 4th Avenue, Suite 306, Anchorage, AK 99510.
- Heifetz, J., D. Anderl, N.E. Maloney, and T.L. Rutecki. 1999. Age validation and analysis of ageing error from marked and recaptured sablefish, *Anoplopoma fimbria*. *Fish. Bull.* 97: 256-263
- Heifetz, J. and J. T. Fujioka. 1991. Movement dynamics of tagged sablefish in the northeastern Pacific Ocean. *Fish. Res.*, 11: 355-374.
- Heifetz, J., J. T. Fujioka, and T. J. Quinn II. 1997. Geographic apportionment of sablefish, *Anoplopoma fimbria*, harvest in the northeastern Pacific Ocean. *In* M. Saunders and M. Wilkins (eds.). Proceedings of the International Symposium on the Biology and Management of Sablefish. pp 229-238. NOAA Tech. Rep. 130.
- Hill, P. S., J. L. Laake, and E. Mitchell. 1999. Results of a pilot program to document interactions between sperm whales and longline vessels in Alaska waters. NOAA Tech. Memo. NMFS-AFSC-108. 42 p.
- Hollowed, A.B. and W.S. Wooster. 1992. Variability of winter ocean conditions and strong year classes of Northeast Pacific groundfish. *ICES Mar. Sci. Symp.* 195, 433-444.
- Hunt, G. L., P. Stabeno, G. Walters, E. Sinclair, R.D. Brodeur, J.M. Napp and N. Bond. 2002. Climate change and control of the southeastern Bering Sea pelagic ecosystem. *Deep-Sea Res.* 49: 5821-5853.
- Hunter, J. R., B. J. Macewicz, and C. A. Kimbrell. 1989. Fecundity and other aspects of reproduction in sablefish, *Anoplopoma fimbria*, in central California waters. *CalCOFI Rep.* 30: 61-72.
- Johnson, S. L., and T. J. Quinn II. 1988. Catch-Age Analysis with Auxiliary Information of sablefish in

- the Gulf of Alaska. Contract report to National Marine Fisheries Service, Auke Bay, Alaska. 79 pp. Center for Fisheries and Ocean Sciences, University of Alaska, Juneau, Alaska.
- Kawakami, T. 1980. A review of sperm whale food. *Sci. Rep. Whales Res. Inst.* 32: 199-218.
- Kimura, D. K. 1990. Approaches to age-structured separable sequential population analysis. *Can. J. Fish. Aquat. Sci.* 47: 2364-2374.
- Kimura, D. K. and J. J. Lyons. 1991. Between-reader bias and variability in the age-determination process. *Fish. Bull.* 89: 53-60.
- Kimura, D. K., A. M. Shimada, and S. A. Lowe. 1993. Estimating von Bertalanffy growth parameters of sablefish, *Anoplopoma fimbria*, and Pacific cod *Gadus macrocephalus* using tag-recapture data. *Fish. Bull.* 91: 271-280.
- Kimura, D. K., A. M. Shimada, and F. R. Shaw. 1998. Stock structure and movement of tagged sablefish, *Anoplopoma fimbria*, in offshore northeast Pacific waters and the effects of El Niño-Southern Oscillation on migration and growth. *Fish. Bull.* 96: 462-481.
- Kimura, D. K., and H. H. Zenger. 1997. Standardizing sablefish (*Anoplopoma fimbria*) longline survey abundance indices by modeling the log-ratio of paired comparative fishing cpues. *ICES J. Mar. Sci.* 54:48-59.
- Kinoshita, R. K., A. Greig, and J. M. Terry. 1995. Economic status of the groundfish fisheries off Alaska, 1995. Available North Pacific Fishery Management Council, 605 W 4th Avenue, Suite 306, Anchorage, Alaska 99510.
- Krieger, K. J. 1997. Sablefish, *Anoplopoma fimbria*, observed from a manned submersible. *In* M. Saunders and M. Wilkins (eds.). *Proceedings of the International Symposium on the Biology and Management of Sablefish.* pp 115-121. NOAA Tech. Rep. 130.
- Laidig, T. E., P. B. Adams, and W. M. Samiere. 1997. Feeding habits of sablefish, *Anoplopoma fimbria*, off the coast of Oregon and California. *In* M. Saunders and M. Wilkins (eds.). *Proceedings of the International Symposium on the Biology and Management of Sablefish.* pp 65-80. NOAA Tech. Rep. 130.
- Langley, A.D. and M. Maunder. 2009. Stock assessment of TRE 7. New Zealand Fisheries Assessment Report 2009/49. 42p.
- Low, L. L., G. K. Tanonaka, and H. H. Shippen. 1976. Sablefish of the Northeastern Pacific Ocean and Bering Sea. Northwest Fisheries Science Center Processed Report. 115 p.
- Mace, P. M. and M. P. Sissenwine. 1993. How much spawning per recruit is enough? *In* S. J. Smith, J. J. Hunt, and D. Rivard [ed.] *Risk evaluation and biological reference points for fisheries management.* *Can. Spec. Publ. Fish. Aquat. Sci.* 120: 101-118.
- Maloney, N. E. and J. Heifetz. 1997. Movements of tagged sablefish, *Anoplopoma fimbria*, released in the eastern Gulf of Alaska. *In* M. Saunders and M. Wilkins (eds.). *Proceedings of the International Symposium on the Biology and Management of Sablefish.* pp 115-121. NOAA Tech. Rep. 130.
- Mason, J. C. R. J. Beamish, and G. A. McFarlane. 1983. Sexual maturity, fecundity, spawning, and early life history of sablefish (*Anoplopoma fimbria*) off the Pacific coast of Canada. *Can. J. Fish. Aquat. Sci.* 40: 2126-2134.
- McDevitt, S. A. 1986. A summary of sablefish catches in the Northeast Pacific Ocean, 1956-84. NOAA Tech. Memo. NMFS F/NWC-101. 34 p.
- McDevitt, S. A. 1990. Growth analysis of sablefish from mark-recapture data from the northeast Pacific. M.S. University of Washington. 87 p.
- McFarlane, G.A. and R.J. Beamish. 1992. Climatic influence linking copepod production with strong year-class in sablefish, *Anoplopoma fimbria*. *Can. J. Fish. Aquat. Sci.* 49: 743– 753.
- McFarlane, G. A. and W. D. Nagata. 1988. Overview of sablefish mariculture and its potential for

- industry. Alaska Sea Grant Report 88-4. PP. 105-120. University of Alaska Fairbanks, Fairbanks, Alaska 99775.
- Meinhold, R. J. and N. D. Singpurwalla, 1983. Understanding the Kalman Filter. *The American Statistician*, May 1983, Vol 37, No. 2, pp. 123-127.
- NMFS (National Marine Fisheries Service). 2005. Final Environmental Impact Statement for Essential Fish Habitat Identification and Conservation in Alaska.
<http://www.fakr.noaa.gov/habitat/seis/efheis.htm>.
- NOAA (National Oceanic and Atmospheric Administration). 2012. NOAA's National Marine Fisheries Service (NMFS), Alaska Region Restricted Access Management (RAM) Pacific Halibut - Sablefish IFQ Report, Fishing Year 2011. alaskafisheries.noaa.gov/ram/ifqreports.htm. National Marine Fisheries Service, Juneau, Alaska.
- NRC (National Research Council). 1998. Improving fish stock assessments. National Academy Press, Washington, DC, 177 pp
- Orlov, A. M., and I. A. Biryukov. 2005. First report of sablefish in spawning condition off the coast of Kamchatka and the Kuril Islands. *ICES J. Mar. Sci.* 62: 1016-1020.
- Rose, G. A. and D. W. Kulka. 1999. Hyperaggregation of fish and fisheries: how catch-per-unit-effort increased as the northern cod (*Gadus morhua*) declined. *Can. J. Fish. Aquat. Sci.* 56 (Suppl. 1): 118-127.
- Rutecki, T. L., M. F. Sigler and H. H. Zenger Jr. 1997. Data report: National Marine Fisheries Service longline surveys, 1991-97.
- Rutecki, T.L. and E.R. Varosi. 1997. Distribution, age, and growth of juvenile sablefish, *Anoplopoma fimbria*, in Southeast Alaska. In M. Saunders and M. Wilkins (eds.). Proceedings of the International Symposium on the Biology and Management of Sablefish. pp 45-54. NOAA Tech. Rep. 130.
- Sasaki, T. 1973. Data on the Japanese blackcod fisheries in the North Pacific--I. Development and history of the Japanese blackcod fisheries through 1972. Unpubl. Rep., 22 p. Far Seas Fish. Res. Lab., Japan Fish Agency, 7-1, Orido 5 chome, Shimizu 424, Japan.
- Sasaki, T. 1985. Studies on the sablefish resources in the North Pacific Ocean. Bulletin 22, (1-108), Far Seas Fishery Laboratory. Shimizu, 424, Japan.
- Sasaki, T. 1987. Stock assessment of sablefish in the eastern Bering Sea, Aleutian Islands region, and the Gulf of Alaska in 1987. Unpubl. Rep., 33 p. Far Seas Fish. Res. Lab., Japan Fish Agency, 7-1, Orido 5 chome, Shimizu 424, Japan.
- Saunders, M. W., B. M. Leaman, V. Haist, R. Hilborn, and G. A. McFarlane. 1996. Sablefish stock assessment for 1996 and recommended yield options for 1997. Unpublished report available Department of Fisheries and Oceans, Biological Sciences Branch, Pacific Biological Station, Nanaimo, British Columbia, V9R 5K6.
- Shotwell, S.K., D.H. Hanselman, and I.M. Belkin. 2014. Toward biophysical synergy: Investigating advection along the Polar Front to identify factors influencing Alaska sablefish recruitment. *Deep-Sea Res. II*, <http://dx.doi.org/10.1016/j.dsr2.2012.08.024>.
- Sigler, M. F. and J. T. Fujioka. 1988. Evaluation of variability in sablefish, *Anoplopoma fimbria*, abundance indices in the Gulf of Alaska using the bootstrap method. *Fish. Bull.* 86: 445-452.
- Sigler, M.F., C.R. Lunsford, and J.T. Fujioka. 2004. Alaska sablefish assessment for 2005. In Stock assessment and fishery evaluation report for the groundfish fisheries of the Gulf of Alaska. North Pacific Fishery Management Council, 605 W 4th Avenue, Suite 306, Anchorage, AK 99510.
- Sigler, M.F., C.R. Lunsford, J.T. Fujioka, and S.A. Lowe. 2002. Alaska sablefish assessment for 2003. In Stock assessment and fishery evaluation report for the groundfish fisheries of the Bering Sea and Aleutian Islands. pp. 449-514. North Pacific Fishery Management Council, 605 W 4th Avenue,

Suite 306, Anchorage, AK 99510.

- Sigler, M. F. and C. R. Lunsford. 2001. Effects of individual quotas on catching efficiency and spawning potential in the Alaska sablefish fishery. *Can. J. Fish. Aquat. Sci.* 58: 1300-1312.
- Sigler, M.F., C.R. Lunsford, J.M. Straley, and J.B. Liddle. 2007. Sperm whale depredation of sablefish longline gear in the northeast Pacific Ocean. *Mar. Mammal Sci.* doi:10.1111/j.1748-7692.2007.00149.
- Sigler, M. F. 2000. Abundance estimation and capture of sablefish, *Anoplopoma fimbria*, by longline gear. *Can. J. Fish. Aquat. Sci.* 57: 1270-1283.
- Sigler, M.F., J.T. Fujioka, and S.A. Lowe. 1999. Alaska sablefish assessment for 2000. *In* Stock assessment and fishery evaluation report for the groundfish fisheries of the Gulf of Alaska. http://www.afsc.noaa.gov/refm/docs/historic_assess/sable99.pdf. North Pacific Fishery Management Council, 605 W 4th Avenue, Suite 306, Anchorage, AK 99510.
- Sigler, M. F., T. L. Rutecki, D. L. Courtney, J. F. Karinen, and M.-S. Yang. 2001. Young-of-the-year sablefish abundance, growth, and diet. *Alaska Fish. Res. Bull.* 8(1): 57-70.
- Sigler, M. F. and H. H. Zenger. 1989. Assessment of Gulf of Alaska sablefish and other groundfish based on the domestic longline survey, 1987. NOAA Tech. Memo. NMFS F/NWC-169. 54 p.
- Soderlund, E., E. Anderson, C. L. Dykstra, T. Geernaert, and A. M. Ranta. 2009. 2008 standardized stock assessment survey. *Int. Pac. Comm. Report of Assessment and Research Activities 2008* 469-496.
- Straley, J., T. O'Connell, S. Mesnick, L. Behnken, and J. Liddle. 2005. Sperm Whale and Longline Fisheries Interactions in the Gulf of Alaska. North Pacific Research Board R0309 Final Report, 15 p.
- Sturdevant, M. V., M. F. Sigler, and J. A. Orsi. 2009. Sablefish predation on juvenile Pacific salmon in the coastal marine waters off southeast Alaska in 1999. *Trans. Amer. Fish. Soc.* 138: 675-691.
- Tanasichuk, R. W. 1997. Diet of sablefish, *Anoplopoma fimbria*, from the southwest coast of Vancouver Island. *In* M. Saunders and M. Wilkins (eds.). *Proceedings of the International Symposium on the Biology and Management of Sablefish.* pp 93-98. NOAA Tech. Rep. 130.
- Thompson, G. G. 2004. Estimation of Pacific Cod Biomass Distributions Based on Alternative Weightings of Trawl Survey Estimates. Appendix to Nov. 2004 Pacific Cod Stock Assessment and Fishery Evaluation report.
- Wing, B. L. 1985. Salmon stomach contents from the Alaska Troll Logbook Program, 1977-84. NOAA Tech. Memo. NMFS F/NWC-91. 41 p.
- Wing, B. L. 1997. Distribution of sablefish, *Anoplopoma fimbria*, larvae in the Eastern Gulf of Alaska. *In* M. Saunders and M. Wilkins (eds.). *Proceedings of the International Symposium on the Biology and Management of Sablefish.* pp 13-26. NOAA Tech. Rep. 130.
- Wolotira, R. J. J., T. M. Sample, S. F. Noel, and C. R. Iten. 1993. Geographic and bathymetric distributions for many commercially important fishes and shellfishes off the west coast of North America, based on research survey and commercial catch data, 1912-1984. NOAA Tech. Memo. NMFS-AFSC-6. 184 pp.
- Yang, M-S. and M. W. Nelson. 2000. Food habits of the commercially important groundfishes in the Gulf of Alaska in 1990, 1993, and 1996. NOAA Tech. Memo. NMFS-AFSC-112. 174 p.
- Yang, M-S. K. Dodd, R. Hibpshman, and A. Whitehouse. 2006. Food habits of groundfishes in the Gulf of Alaska in 1999 and 2001. NOAA Tech. Memo. NMFS-AFSC-164. 199 p.
- Zenger, H. H., Jr., M. F. Sigler, and E. R. Varosi. 1994. Assessment of Gulf of Alaska sablefish and other groundfish species based on the 1988 National Marine Fisheries Service longline survey. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-AFSC-39, 55 p.

Tables

Table 3.1. Alaska sablefish catch (t). The values include landed catch and discard estimates. Discards were estimated for U.S. fisheries before 1993 by multiplying reported catch by 2.9% for fixed gear and 26.9% for trawl gear (1994-1997 averages) because discard estimates were unavailable. Eastern includes West Yakutat and East Yakutat / Southeast. 2015 catches are as of 10/29/2015 (www.akfin.org).

Year	Grand total	BY AREA								BY GEAR	
		Bering Sea	Aleutians	Western	Central	Eastern	West Yakutat	East Yak/SEO	Unknown	Fixed	Trawl
1960	3,054	1,861	0	0	0	1,193			0	3,054	0
1961	16,078	15,627	0	0	0	451			0	16,078	0
1962	26,379	25,989	0	0	0	390			0	26,379	0
1963	16,901	13,706	664	266	1,324	941			0	10,557	6,344
1964	7,273	3,545	1,541	92	955	1,140			0	3,316	3,957
1965	8,733	4,838	1,249	764	1,449	433			0	925	7,808
1966	15,583	9,505	1,341	1,093	2,632	1,012			0	3,760	11,823
1967	19,196	11,698	1,652	523	1,955	3,368			0	3,852	15,344
1968	30,940	14,374	1,673	297	1,658	12,938			0	11,182	19,758
1969	36,831	16,009	1,673	836	4,214	14,099			0	15,439	21,392
1970	37,858	11,737	1,248	1,566	6,703	16,604			0	22,729	15,129
1971	43,468	15,106	2,936	2,047	6,996	16,382			0	22,905	20,563
1972	53,080	12,758	3,531	3,857	11,599	21,320			15	28,538	24,542
1973	36,926	5,957	2,902	3,962	9,629	14,439			37	23,211	13,715
1974	34,545	4,258	2,477	4,207	7,590	16,006			7	25,466	9,079
1975	29,979	2,766	1,747	4,240	6,566	14,659			1	23,333	6,646
1976	31,684	2,923	1,659	4,837	6,479	15,782			4	25,397	6,287
1977	21,404	2,718	1,897	2,968	4,270	9,543			8	18,859	2,545
1978	10,394	1,193	821	1,419	3,090	3,870			1	9,158	1,236
1979	11,814	1,376	782	999	3,189	5,391			76	10,350	1,463
1980	10,444	2,205	275	1,450	3,027	3,461			26	8,396	2,048
1981	12,604	2,605	533	1,595	3,425	4,425			22	10,994	1,610
1982	12,048	3,238	964	1,489	2,885	3,457			15	10,204	1,844
1983	11,715	2,712	684	1,496	2,970	3,818			35	10,155	1,560
1984	14,109	3,336	1,061	1,326	3,463	4,618			305	10,292	3,817
1985	14,465	2,454	1,551	2,152	4,209	4,098			0	13,007	1,457
1986	28,892	4,184	3,285	4,067	9,105	8,175			75	21,576	7,316
1987	35,163	4,904	4,112	4,141	11,505	10,500			2	27,595	7,568
1988	38,406	4,006	3,616	3,789	14,505	12,473			18	29,282	9,124
1989	34,829	1,516	3,704	4,533	13,224	11,852			0	27,509	7,320
1990	32,115	2,606	2,412	2,251	13,786	11,030			30	26,598	5,518
1991	27,073	1,318	2,168	1,821	11,662	10,014			89	23,124	3,950
1992	24,932	586	1,497	2,401	11,135	9,171			142	21,614	3,318
1993	25,417	669	2,078	740	11,955	9,976	4,620	5,356	0	22,912	2,506
1994	23,577	694	1,725	539	9,376	11,243	4,493	6,750	0	20,639	2,938
1995	20,692	930	1,119	1,747	7,673	9,223	3,872	5,352	0	18,079	2,613
1996	17,275	648	764	1,542	6,773	7,548	2,893	4,655	0	15,088	2,187
1997	14,607	552	781	1,374	6,234	5,666	1,930	3,735	0	12,975	1,632
1998	13,867	563	535	1,432	5,915	5,422	1,956	3,467	0	12,380	1,487
1999	13,585	675	681	1,488	5,874	4,867	1,709	3,159	0	11,601	1,985
2000	15,565	742	1,049	1,582	6,173	6,020	2,066	3,953	0	13,546	2,019
2001	14,064	864	1,074	1,588	5,518	5,021	1,737	3,284	0	12,281	1,783
2002	14,748	1,144	1,119	1,865	6,180	4,441	1,550	2,891	0	12,505	2,243
2003	16,411	1,012	1,118	2,118	6,993	5,170	1,822	3,347	0	14,351	2,060
2004	17,518	1,041	955	2,170	7,310	6,041	2,241	3,801	0	15,861	1,656
2005	16,580	1,070	1,481	1,929	6,701	5,399	1,824	3,575	0	15,024	1,556
2006	15,551	1,079	1,151	2,151	5,921	5,251	1,889	3,362	0	14,305	1,246
2007	15,957	1,182	1,168	2,101	6,003	5,502	2,074	3,429	0	14,721	1,235
2008	14,674	1,141	901	1,679	5,543	5,410	2,056	3,354	0	13,552	1,122
2009	13,128	916	1,100	1,423	5,005	4,684	1,831	2,853	0	12,071	1,057
2010	11,980	755	1,094	1,354	4,508	4,269	1,578	2,690	0	10,976	1,004
2011	12,971	705	1,024	1,402	4,919	4,921	1,896	3,024	0	11,792	1,179
2012	13,868	743	1,205	1,353	5,329	5,238	2,033	3,205	0	12,767	1,102
2013	13,642	634	1,062	1,385	5,207	5,354	2,106	3,247	0	12,604	1,038
2014	11,582	315	818	1,201	4,755	4,493	1,671	2,822	0	10,557	1,025
2015	10,094	197	372	867	4,176	4,482	1,794	2,688	0	9,699	902

Table 3.2. Catch (t) in the Aleutian Islands and the Bering Sea by gear type from 1991-2015. Both CDQ and non-CDQ catches are included. Catches in 1991-1999 are averages. Catch as of October 29, 2015 (www.akfin.org).

Aleutian Islands				
<u>Year</u>	<u>Pot</u>	<u>Trawl</u>	<u>Longline</u>	<u>Total</u>
1991-1999	6	73	1,210	1,289
2000	103	33	913	1,049
2001	111	39	925	1,074
2002	105	39	975	1,119
2003	316	42	760	1,118
2004	384	32	539	955
2005	688	115	679	1,481
2006	461	60	629	1,151
2007	632	40	496	1,169
2008	177	76	646	899
2009	78	75	947	1,100
2010	59	74	963	1,097
2011	141	47	837	1,024
2012	77	148	979	1,205
2013	87	58	917	1,062
2014	160	26	632	818
2015	12	15	345	372
Bering Sea				
1991-1999	5	189	539	733
2000	40	284	418	742
2001	106	353	405	864
2002	382	295	467	1,144
2003	363	231	417	1,012
2004	435	293	313	1,041
2005	595	273	202	1,070
2006	621	84	373	1,078
2007	879	92	211	1,182
2008	754	183	204	1,141
2009	557	93	266	916
2010	452	30	274	755
2011	405	44	256	705
2012	431	93	218	743
2013	352	133	149	634
2014	164	34	116	315
2015	100	17	81	197

Table 3.3. Discarded catches of sablefish (amount [t], percent of total catch, total catch [t]) by gear (H&L=hook & line, Other = Pot, trawl, and jig, combined for confidentiality) by FMP area for 2007-2014. Source: NMFS Alaska Regional Office via AKFIN, October 23, 2015.

Year	Gear	BSAI			GOA			Combined		
		Discard	%Discard	Catch	Discard	%Discard	Catch	Discard	%Discard	Catch
2007	Total	70	3.00%	2,350	556	4.09%	13,608	627	3.93%	15,958
	H&L	16	2.25%	707	256	2.07%	12,379	272	2.08%	13,086
	Other	55	3.32%	1,643	300	24.43%	1,229	355	12.35%	2,872
2008	Total	100	4.90%	2,040	750	5.99%	12,512	850	5.84%	14,552
	H&L	93	10.99%	850	669	5.74%	11,639	762	6.10%	12,490
	Other	7	0.55%	1,190	81	9.27%	872	87	4.24%	2,062
2009	Total	25	1.23%	2,016	736	6.66%	11,046	761	5.82%	13,062
	H&L	17	1.39%	1,213	655	6.45%	10,157	672	5.91%	11,370
	Other	8	0.98%	803	81	9.10%	889	89	5.25%	1,692
2010	Total	43	2.31%	1,852	419	4.13%	10,131	462	3.85%	11,983
	H&L	36	2.89%	1,237	371	4.02%	9,231	407	3.89%	10,468
	Other	7	1.15%	615	47	5.27%	900	54	3.60%	1,515
2011	Total	25	1.47%	1,730	575	5.11%	11,239	600	4.63%	12,969
	H&L	18	1.63%	1,093	396	3.90%	10,145	413	3.68%	11,237
	Other	8	1.20%	637	179	16.36%	1,095	187	10.79%	1,732
2012	Total	25	1.28%	1,948	318	2.67%	11,921	343	2.47%	13,868
	H&L	13	1.10%	1,197	253	2.29%	11,060	266	2.17%	12,257
	Other	12	1.56%	750	65	7.52%	861	76	4.75%	1,611
2013	Total	30	1.79%	1,696	637	5.34%	11,947	668	4.90%	13,643
	H&L	27	2.51%	1,066	590	5.31%	11,101	617	5.07%	12,167
	Other	4	0.59%	630	48	5.62%	846	51	3.47%	1,476
2014	Total	38	3.37%	1,132	516	4.94%	10,450	554	4.79%	11,582
	H&L	37	4.94%	748	438	4.62%	9,483	475	4.64%	10,231
	Other	1	0.33%	385	78	8.10%	967	80	5.89%	1,351
2007-2014 Mean	Total	45	2.42%	1,845	563	4.85%	11,607	608	4.52%	13,452
	H&L	32	3.16%	1,014	453	4.26%	10,649	486	4.16%	11,663
	Other	13	1.51%	832	110	11.48%	957	122	6.84%	1,789

Table 3.4. Bycatch (t) of FMP Groundfish species in the targeted sablefish fishery averaged from 2010-2014. Other = Pot and trawl combined because of confidentiality. Source: AKFIN, October 23, 2015.

Species	Hook and Line			Other Gear			All Gear		
	Discard	Retained	Total	Discard	Retained	Total	Discard	Retained	Total
GOA Thornyhead Rockfish	170	377	546	4	25	29	174	402	575
GOA Shark	380	0	380	0	0	0	380	0	380
GOA Skate	301	10	312	2	0	2	304	10	314
GOA Shortraker Rockfish	141	94	235	10	8	18	151	102	253
GOA Arrowtooth Flounder	160	18	178	50	1	51	210	19	229
GOA Roughey Rockfish	69	83	152	1	3	4	70	86	156
BSAI Other Rockfish	41	81	122	2	1	2	42	82	124
GOA Pacific Cod	45	36	81	0	2	2	45	39	83
BSAI Skate	77	1	78	0	0	0	77	1	78
BSAI Arrowtooth Flounder	29	10	40	37	1	37	66	11	77
BSAI Greenland Turbot	22	39	60	5	0	6	27	39	66
GOA Other Rockfish	30	9	39	0	0	0	30	9	40
GOA Deep Water Flatfish	9	0	9	16	5	21	25	5	30
BSAI Shortraker Rockfish	8	7	15	0	0	0	8	7	15
GOA Pacific Ocean Perch	1	0	1	1	11	12	2	11	13
BSAI Other Flatfish	11	0	11	1	0	1	12	0	12
GOA Sculpin	11	0	11	0	0	0	11	0	11
GOA Rex Sole	0	0	0	8	3	11	8	3	11
GOA Demersal Shelf Rockfish	1	9	10	0	0	0	1	9	10
Total	1,505	775	2,280	137	60	197	1,642	835	2,477

Table 3.5. Bycatch of nontarget species and HAPC biota in the targeted sablefish fishery. Source: NMFS AKRO Blend/Catch Accounting System via AKFIN, October 23, 2015. Birds are in numbers.

Group Name	<u>Estimated Catch (t)</u>					
	<u>2009</u>	<u>2010</u>	<u>2011</u>	<u>2012</u>	<u>2013</u>	<u>2014</u>
Benthic urochordata	0.01	0.13	0.13	1.08	0.00	0.00
Birds	519	524	1,868	227	665	539
Bivalves	0.04	0.04	0.05	0.01	0.00	0.00
Brittle star unidentified	0.44	0.12	0.44	4.52	0.10	0.60
Corals Bryozoans	2.20	3.35	5.54	7.55	12.75	0.91
Dark Rockfish	0.14	0.00	0.00	0.03	0.06	0.03
Eelpouts	1.83	1.37	0.58	0.63	1.11	0.74
Giant Grenadier	5,979	4,770	6,943	7,009	7,929	4,577
Greenlings	0.07	0.00	0.02	0.00	0.00	0.00
Grenadier	1,133	858	842	1,017	1,466	854
Hermit crab unidentified	0.10	0.17	0.22	0.08	0.09	0.16
Large sculpin	30.56	19.04	3.88	5.13	19.63	5.53
Invertebrate unidentified	1.52	2.40	2.02	6.81	0.18	0.11
Misc crabs	3.29	1.81	1.14	0.31	0.51	0.50
Misc crustaceans	2.34	0.00	0.00	0.00	0.00	0.15
Misc fish	5.02	6.20	8.43	10.11	27.75	28.09
Scypho jellies	0.08	0.10	0.68	0.00	0.00	5.51
Sea anemone unidentified	2.25	1.40	3.29	0.99	0.88	2.99
Sea pens whips	0.52	0.32	1.58	0.25	0.27	1.92
Sea star	2.95	4.11	3.45	2.99	18.38	10.74
Snails	10.79	11.02	20.08	12.07	8.80	3.65
Sponge unidentified	2.17	1.12	2.09	0.94	3.30	1.60
Urchins, dollars, cucumbers	1.64	0.55	0.26	0.78	0.72	0.79

Table 3.6. Prohibited Species Catch (PSC) estimates reported in tons for halibut and numbers of animals for crab and salmon, by year, and fisheries management plan (BSAI or GOA) for the sablefish fishery. Other = Pot and trawl combined because of confidentiality. Source: NMFS AKRO Blend/Catch Accounting System PSCNQ via AKFIN, October 29, 2015.

BSAI								
Hook and Line	Year	Bairdi	Chinook	Golden KC	Halibut	Other salmon	Opilio	Red KC
	2011	0	0	527	101	0	18	18
	2012	0	0	420	82	0	0	7
	2013	0	15	465	66	8	0	0
	2014	0	0	471	38	0	0	40
	2015	0	9	181	23	0	0	159
	Mean	0	5	413	62	2	4	45
Other	2011	808	0	198,724	13	0	249	294
	2012	0	0	16,754	10	0	119	0
	2013	222	0	788	18	0	314	0
	2014	0	0	3,193	6	0	1,679	0
	2015	0	0	11,270	1	0	26	0
	Mean	206	0	46,146	10	0	477	59
BSAI Mean		206	5	46,558	72	2	481	104
GOA								
Hook and Line	2011	0	0	120	212	304	0	0
	2012	0	0	23	293	248	0	0
	2013	78	4	93	273	519	0	24
	2014	6	39	39	250	284	0	0
	2015	165	25	24	256	375	0	29
	Mean	50	14	60	257	346	0	11
Other	2011	0	0	132	6	0	0	0
	2012	0	0	9	5	0	0	0
	2013	0	0	0	12	12	0	0
	2014	0	0	18	2	0	0	0
	2015	20	0	0	3	0	0	0
	Mean	4	0	32	5	2	0	0
GOA Mean		54	14	92	262	348	0	11

Table 3.7. Summary of management measures with time series of catch, ABC, OFL, and TAC.

Year	Catch(t)	OFL	ABC	TAC	Management measure
1980	10,444			18,000	Amendment 8 to the Gulf of Alaska Fishery Management Plan established the West and East Yakutat management areas for sablefish.
1981	12,604			19,349	
1982	12,048			17,300	
1983	11,715			14,480	
1984	14,109			14,820	
1985	14,465			13,480	Amendment 14 of the GOA FMP allocated sablefish quota by gear type: 80% to fixed gear and 20% to trawl gear in WGOA and CGOA and 95% fixed to 5% trawl in the EGOA.
1986	28,892			21,450	Pot fishing banned in Eastern GOA.
1987	35,163			27,700	Pot fishing banned in Central GOA.
1988	38,406			36,400	
1989	34,829			32,200	Pot fishing banned in Western GOA.
1990	32,115			33,200	Amendment 15 of the BSAI FMP allocated sablefish quota by gear type: 50% to fixed gear in and 50% to trawl in the EBS, and 75% fixed to 25% trawl in the Aleutian Islands.
1991	27,073			28,800	
1992	24,932			25,200	Pot fishing banned in Bering Sea (57 FR 37906).
1993	25,417			25,000	
1994	23,577			28,840	
1995	20,692			25,300	Amendment 20 to the Gulf of Alaska Fishery Management Plan and 15 to the Bering Sea/Aleutian Islands Fishery Management Plan established IFQ management for sablefish beginning in 1995. These amendments also allocated 20% of the fixed gear allocation of sablefish to a CDQ reserve for the Bering Sea and Aleutian Islands.
1996	17,275			19,380	Pot fishing ban repealed in Bering Sea except from June 1-30.
1997	14,607	27,900	19,600	17,200	Maximum retainable allowances for sablefish were revised in the Gulf of Alaska. The percentage depends on the basis species.
1998	13,867	26,500	16,800	16,800	
1999	13,585	24,700	15,900	15,900	
2000	15,565	21,400	17,300	17,300	
2001	14,064	20,700	16,900	16,900	
2002	14,748	26,100	17,300	17,300	
2003	16,411	28,900	18,400	20,900	
2004	17,518	30,800	23,000	23,000	
2005	16,580	25,400	21,000	21,000	
2006	15,551	25,300	21,000	21,000	
2007	15,957	23,750	20,100	20,100	
2008	14,674	21,310	18,030	18,030	Pot fishing ban repealed in Bering Sea for June 1-30 (74 FR 28733).
2009	13,128	19,000	16,080	16,080	
2010	11,980	21,400	15,230	15,230	
2011	12,971	20,700	16,040	16,040	
2012	13,868	20,400	17,240	17,240	
2013	13,642	19,180	16,230	16,230	
2014	11,582	16,160	13,722	13,722	
2015	10,094	16,128	13,657	13,657	

Table 3.8. Sample sizes for aged fish and length data collected from Alaska sablefish. Japanese fishery data from Sasaki (1985), U.S. fishery data from the observer databases, and longline survey data from longline survey databases. Trawl survey data from AKFIN. All fish were sexed before measurement, except for the Japanese fishery data.

Year	LENGTH						AGE			
	U.S. NMFS trawl survey (GOA)	Japanese fishery Trawl Longline		U.S. fishery Trawl Longline		Cooperative longline survey	Domestic longline survey	Cooperative longline survey	Domestic longline survey	U.S. longline fishery
1963			30,562							
1964		3,337	11,377							
1965		6,267	9,631							
1966		27,459	13,802							
1967		31,868	12,700							
1968		17,727								
1969		3,843								
1970		3,456								
1971		5,848	19,653							
1972		1,560	8,217							
1973		1,678	16,332							
1974			3,330							
1975										
1976			7,704							
1977			1,079							
1978			9,985							
1979			1,292			19,349				
1980			1,944			40,949				
1981						34,699		1,146		
1982						65,092				
1983						66,517		889		
1984	12,964					100,029				
1985						125,129		1,294		
1986						128,718				
1987	9,610					102,639		1,057		
1988						114,239				
1989						115,067		655		
1990	4,969			1,229	32,936	78,794	101,530			
1991				721	28,182	69,653	95,364	902		
1992				0	20,929	79,210	104,786			
1993	7,168			468	21,943	80,596	94,699	1,178		
1994				89	11,914	74,153	70,431			
1995				87	17,735		80,826			
1996	4,615			239	14,416		72,247		1,176	
1997				0	20,330		82,783		1,214	
1998				35	8,932		57,773		1,191	
1999	4,281			1,268	28,070		79,451		1,186	1,141
2000				472	32,208		62,513		1,236	1,152
2001				473	30,315		83,726		1,214	1,003
2002				526	33,719		75,937		1,136	1,059
2003	5,003			503	36,077		77,678		1,128	1,185
2004				694	31,199		82,767		1,185	1,145
2005	4,901			2,306	36,213		74,433		1,074	1,164
2006				721	32,497		78,625		1,178	1,154
2007	3,773			860	29,854		73,480		1,174	1,115
2008				2,018	23,414		71,661		1,184	1,164
2009	3,934			1,837	24,674		67,978		1,197	1,126
2010				1,634	24,530		75,010		1,176	1,159
2011	2,114			1,877	22,659		87,498		1,199	1,190
2012				2,533	22,203		63,116		1,186	1,165
2013	1,249			2,674	16,093		51,586		1,190	1,157
2014				2,210	19,524		52,290		1,183	1,126
2015	3,277						52,110			

Table 3.9. Average catch rate (pounds/hook) for fishery data by year and region. SE = standard error, CV = coefficient of variation. C = confidential due to less than three vessels or sets. These data are still used in the combined index.

Observer Fishery Data											
Aleutian Islands-Observer						Bering Sea-Observer					
Year	CPUE	SE	CV	Sets	Vessels	Year	CPUE	SE	CV	Sets	Vessels
1990	0.53	0.05	0.10	193	8	1990	0.72	0.11	0.15	42	8
1991	0.50	0.03	0.07	246	8	1991	0.28	0.06	0.20	30	7
1992	0.40	0.06	0.15	131	8	1992	0.25	0.11	0.43	7	4
1993	0.28	0.04	0.14	308	12	1993	0.09	0.03	0.36	4	3
1994	0.29	0.05	0.18	138	13	1994	C	C	C	2	2
1995	0.30	0.04	0.14	208	14	1995	0.41	0.07	0.17	38	10
1996	0.23	0.03	0.12	204	17	1996	0.63	0.19	0.30	35	15
1997	0.35	0.07	0.20	117	9	1997	C	C	C	0	0
1998	0.29	0.05	0.17	75	12	1998	0.17	0.03	0.18	28	9
1999	0.38	0.07	0.17	305	14	1999	0.29	0.09	0.32	27	10
2000	0.29	0.03	0.11	313	15	2000	0.28	0.09	0.31	21	10
2001	0.26	0.04	0.15	162	9	2001	0.31	0.02	0.07	18	10
2002	0.32	0.03	0.11	245	10	2002	0.10	0.02	0.22	8	4
2003	0.26	0.04	0.17	170	10	2003	C	C	C	8	2
2004	0.21	0.04	0.21	138	7	2004	0.17	0.05	0.31	9	4
2005	0.15	0.05	0.34	23	6	2005	0.23	0.02	0.16	9	6
2006	0.23	0.04	0.16	205	11	2006	0.17	0.05	0.21	68	15
2007	0.35	0.10	0.29	198	7	2007	0.28	0.05	0.18	34	8
2008	0.37	0.04	0.10	247	6	2008	0.38	0.22	0.58	12	5
2009	0.29	0.05	0.22	335	10	2009	0.14	0.04	0.21	24	5
2010	0.27	0.04	0.14	459	12	2010	0.17	0.03	0.19	42	8
2011	0.25	0.05	0.19	401	9	2011	0.10	0.01	0.13	12	4
2012	0.25	0.10	0.15	363	8	2012	C	C	C	6	1
2013	0.28	0.06	0.22	613	7	2013	0.21	0.10	0.46	27	5
2014	0.24	0.04	0.18	487	6	2014	0.25	0.12	0.48	8	3

Table 3.9 (cont.)

Western Gulf-Observer						Central Gulf-Observer					
Year	CPUE	SE	CV	Sets	Vessels	Year	CPUE	SE	CV	Sets	Vessels
1990	0.64	0.14	0.22	178	7	1990	0.54	0.04	0.07	653	32
1991	0.44	0.06	0.13	193	16	1991	0.62	0.06	0.09	303	24
1992	0.38	0.05	0.14	260	12	1992	0.59	0.05	0.09	335	19
1993	0.35	0.03	0.09	106	12	1993	0.60	0.04	0.07	647	32
1994	0.32	0.03	0.10	52	5	1994	0.65	0.06	0.09	238	15
1995	0.51	0.04	0.09	432	22	1995	0.90	0.07	0.08	457	41
1996	0.57	0.05	0.10	269	20	1996	1.04	0.07	0.07	441	45
1997	0.50	0.05	0.10	349	20	1997	1.07	0.08	0.08	377	41
1998	0.50	0.03	0.07	351	18	1998	0.90	0.06	0.06	345	32
1999	0.53	0.07	0.12	244	14	1999	0.87	0.08	0.10	269	28
2000	0.49	0.06	0.13	185	12	2000	0.93	0.05	0.06	319	30
2001	0.50	0.05	0.10	273	16	2001	0.70	0.04	0.06	347	31
2002	0.51	0.05	0.09	348	15	2002	0.84	0.07	0.08	374	29
2003	0.45	0.04	0.10	387	16	2003	0.99	0.07	0.07	363	34
2004	0.47	0.08	0.17	162	10	2004	1.08	0.10	0.09	327	29
2005	0.58	0.07	0.13	447	13	2005	0.89	0.06	0.07	518	32
2006	0.42	0.04	0.13	306	15	2006	0.82	0.06	0.08	361	33
2007	0.37	0.04	0.11	255	12	2007	0.93	0.06	0.07	289	30
2008	0.46	0.07	0.16	255	11	2008	0.84	0.07	0.08	207	27
2009	0.44	0.09	0.21	208	11	2009	0.77	0.06	0.07	320	33
2010	0.42	0.06	0.14	198	10	2010	0.80	0.05	0.07	286	31
2011	0.54	0.12	0.22	196	12	2011	0.85	0.08	0.10	213	28
2012	0.38	0.04	0.11	147	13	2012	0.74	0.07	0.09	298	27
2013	0.34	0.02	0.06	325	18	2013	0.51	0.05	0.10	419	34
2014	0.41	0.06	0.15	190	16	2014	0.56	0.03	0.05	585	57

Table 3.9 (cont.)

West Yakutat-Observer						East Yakutat/SE-Observer					
Year	CPUE	SE	CV	Sets	Vessels	Year	CPUE	SE	CV	Sets	Vessels
1990	0.95	0.24	0.25	75	9	1990	C	C	C	0	0
1991	0.65	0.07	0.10	164	12	1991	C	C	C	17	2
1992	0.64	0.18	0.27	98	6	1992	C	C	C	20	1
1993	0.71	0.07	0.10	241	12	1993	C	C	C	26	2
1994	0.65	0.17	0.27	81	8	1994	C	C	C	5	1
1995	1.02	0.10	0.10	158	21	1995	1.45	0.20	0.14	101	19
1996	0.97	0.07	0.07	223	28	1996	1.20	0.11	0.09	137	24
1997	1.16	0.11	0.09	126	20	1997	1.10	0.14	0.13	84	17
1998	1.21	0.10	0.08	145	23	1998	1.27	0.12	0.10	140	25
1999	1.20	0.15	0.13	110	19	1999	0.94	0.12	0.13	85	11
2000	1.28	0.10	0.08	193	32	2000	0.84	0.13	0.16	81	14
2001	1.03	0.07	0.07	184	26	2001	0.84	0.08	0.09	110	14
2002	1.32	0.13	0.10	155	23	2002	1.20	0.23	0.19	121	14
2003	1.36	0.10	0.07	216	27	2003	1.29	0.13	0.10	113	19
2004	1.23	0.09	0.08	210	24	2004	1.08	0.10	0.09	135	17
2005	1.32	0.09	0.07	352	24	2005	1.18	0.13	0.11	181	16
2006	0.96	0.10	0.10	257	30	2006	0.93	0.11	0.11	104	18
2007	1.02	0.11	0.11	208	24	2007	0.92	0.15	0.17	85	16
2008	1.40	0.12	0.08	173	23	2008	1.06	0.13	0.12	103	17
2009	1.34	0.12	0.09	148	23	2009	0.98	0.12	0.12	94	13
2010	1.11	0.09	0.08	136	22	2010	0.97	0.17	0.17	76	12
2011	1.18	0.09	0.07	186	24	2011	0.98	0.09	0.10	196	16
2012	0.97	0.09	0.10	255	24	2012	0.93	0.11	0.12	104	15
2013	1.11	0.15	0.13	109	20	2013	0.91	0.12	0.14	165	22
2014	0.83	0.07	0.09	149	22	2014	0.88	0.08	0.09	207	33

Table 3.9 (cont.)

Aleutian Islands-Logbook						Bering Sea-Logbook					
Year	CPUE	SE	CV	Sets	Vessels	Year	CPUE	SE	CV	Sets	Vessels
1999	0.29	0.04	0.15	167	15	1999	0.56	0.08	0.14	291	43
2000	0.24	0.05	0.21	265	16	2000	0.21	0.05	0.22	169	23
2001	0.38	0.16	0.41	36	5	2001	0.35	0.11	0.33	61	8
2002	0.48	0.19	0.39	33	5	2002	C	C	C	5	2
2003	0.36	0.11	0.30	139	10	2003	0.24	0.13	0.53	25	6
2004	0.45	0.11	0.25	102	7	2004	0.38	0.09	0.24	202	8
2005	0.46	0.15	0.33	109	8	2005	0.36	0.07	0.19	86	10
2006	0.51	0.16	0.31	61	5	2006	0.38	0.07	0.18	106	9
2007	0.38	0.22	0.58	61	3	2007	0.37	0.08	0.21	147	8
2008	0.30	0.03	0.12	119	4	2008	0.52	0.20	0.39	94	7
2009	0.23	0.07	0.06	204	7	2009	0.25	0.04	0.14	325	18
2010	0.25	0.05	0.20	497	9	2010	0.30	0.08	0.27	766	12
2011	0.23	0.07	0.30	609	12	2011	0.22	0.03	0.13	500	24
2012	0.26	0.03	0.14	893	12	2012	0.30	0.04	0.15	721	21
2013	0.26	0.06	0.22	457	7	2013	0.20	0.04	0.18	460	15
2014	0.25	0.07	0.27	272	5	2014	0.34	0.05	0.15	436	15

Western Gulf-Logbook						Central Gulf-Logbook					
Year	CPUE	SE	CV	Sets	Vessels	Year	CPUE	SE	CV	Sets	Vessels
1999	0.64	0.06	0.09	245	27	1999	0.80	0.05	0.06	817	60
2000	0.60	0.05	0.09	301	32	2000	0.79	0.04	0.05	746	64
2001	0.47	0.05	0.10	109	24	2001	0.74	0.06	0.08	395	52
2002	0.60	0.08	0.13	78	14	2002	0.83	0.06	0.07	276	41
2003	0.39	0.04	0.11	202	24	2003	0.87	0.07	0.08	399	45
2004	0.65	0.06	0.09	766	26	2004	1.08	0.05	0.05	1676	80
2005	0.78	0.08	0.11	571	33	2005	0.98	0.07	0.07	1154	63
2006	0.69	0.08	0.11	1067	38	2006	0.87	0.04	0.05	1358	80
2007	0.59	0.06	0.10	891	31	2007	0.83	0.04	0.05	1190	69
2008	0.71	0.06	0.08	516	29	2008	0.88	0.05	0.06	1039	68
2009	0.53	0.06	0.11	824	33	2009	0.95	0.08	0.08	1081	73
2010	0.48	0.04	0.08	1297	46	2010	0.66	0.03	0.05	1171	80
2011	0.50	0.05	0.10	1148	46	2011	0.80	0.06	0.07	1065	71
2012	0.50	0.04	0.08	1142	37	2012	0.79	0.06	0.07	1599	82
2013	0.35	0.03	0.07	1476	32	2013	0.48	0.03	0.07	2102	73
2014	0.39	0.03	0.08	1008	28	2014	0.52	0.04	0.08	2051	72

Table 3.9 (cont.)

West Yakutat-Logbook						East Yakutat/SE-Logbook					
Year	CPUE	SE	CV	Sets	Vessels	Year	CPUE	SE	CV	Sets	Vessels
1999	1.08	0.08	0.08	233	36	1999	0.91	0.08	0.08	183	22
2000	1.04	0.06	0.06	270	42	2000	0.98	0.08	0.08	190	26
2001	0.89	0.09	0.11	203	29	2001	0.98	0.09	0.09	109	21
2002	0.99	0.07	0.07	148	28	2002	0.83	0.06	0.07	108	22
2003	1.26	0.10	0.08	104	23	2003	1.13	0.10	0.09	117	22
2004	1.27	0.06	0.05	527	54	2004	1.19	0.05	0.04	427	55
2005	1.13	0.05	0.04	1158	70	2005	1.15	0.05	0.05	446	77
2006	0.97	0.05	0.06	1306	84	2006	1.06	0.04	0.04	860	107
2007	0.97	0.05	0.05	1322	89	2007	1.13	0.04	0.04	972	122
2008	0.97	0.05	0.05	1118	74	2008	1.08	0.05	0.05	686	97
2009	1.23	0.07	0.06	1077	81	2009	1.12	0.05	0.05	620	87
2010	0.98	0.05	0.05	1077	85	2010	1.04	0.05	0.05	744	99
2011	0.95	0.07	0.07	1377	75	2011	1.01	0.04	0.04	877	112
2012	0.89	0.06	0.06	1634	86	2012	1.00	0.05	0.05	972	102
2013	0.74	0.06	0.07	1953	79	2013	0.86	0.05	0.06	865	88
2014	0.73	0.04	0.06	1591	74	2014	0.88	0.05	0.05	797	83

Table 3.10. Sablefish abundance index values (1,000's) for Alaska (200-1,000 m) including deep gully habitat, from the Japan-U.S. Cooperative Longline Survey, Domestic Longline Survey, and Japanese and U.S. longline fisheries. Relative population number equals CPUE in numbers weighted by respective strata areas. Relative population weight equals CPUE measured in weight multiplied by strata areas. Indices were extrapolated for survey areas not sampled every year, including Aleutian Islands 1979, 1995, 1997, 1999, 2001, 2003, 2005, and 2007, 2009, 2011, 2013, and 2015, and Bering Sea 1979-1981, 1995, 1996, 1998, 2000, 2002, 2004, 2006, 2008, 2009, 2010, 2012, and 2014. NMFS trawl survey biomass estimates (kilotons) are from the Gulf of Alaska at depths <500 m.

Year	RELATIVE POPULATION NUMBER		RELATIVE POPULATION WEIGHT/BIOMASS				
	Coop. longline survey	Dom. longline survey	Jap. longline fishery	Coop. longline survey	Dom. longline survey	U.S. fishery	NMFS Trawl survey
1964			1,452				
1965			1,806				
1966			2,462				
1967			2,855				
1968			2,336				
1969			2,443				
1970			2,912				
1971			2,401				
1972			2,247				
1973			2,318				
1974			2,295				
1975			1,953				
1976			1,780				
1977			1,511				
1978			942				
1979	413		809	1,075			
1980	388		1,040	968			
1981	460		1,343	1,153			
1982	613			1,572			
1983	621			1,595			
1984	685			1,822			294
1985	903			2,569			
1986	838			2,456			
1987	667			2,068			271
1988	707			2,088			
1989	661			2,178			
1990	450	649		1,454	2,141	1,201	214
1991	386	593		1,321	2,071	1,066	
1992	402	511		1,390	1,758	908	
1993	395	563		1,318	1,894	904	250
1994	366	489		1,288	1,882	822	
1995		501			1,803	1,243	
1996		520			2,017	1,201	145
1997		491			1,764	1,341	
1998		477			1,662	1,130	
1999		520			1,740	1,316	104
2000		462			1,597	1,139	
2001		535			1,798	1,111	238
2002		561			1,916	1,152	
2003		532			1,759	1,218	189
2004		544			1,738	1,357	
2005		533			1,695	1,304	179
2006		580			1,848	1,206	
2007		500			1,584	1,268	111
2008		472			1,550	1,361	
2009		491			1,580	1,152	107
2010		542			1,778	1,054	
2011		556			1,683	1,048	84
2012		438			1,280	1,023	
2013		416			1,276	893	60
2014		479			1,432	949	
2015		378			1,169		67

Table 3.11. Count of stations where sperm (S) or killer whale (K) depredation occurred in the six sablefish management areas. The number of stations sampled that are used for RPN calculations are in parentheses. Areas not surveyed in a given year are left blank. If there were no whale depredation data taken, it is denoted with an “n/a”. Killer whale depredation did not always occur on all skates of gear, and only those skates with depredation were cut from calculations of RPNs and RPWs.

Year	BS (16)		AI (14)		WG (10)		CG (16)		WY (8)		EY/SE (17)	
	S	K	S	K	S	K	S	K	S	K	S	K
1996			n/a	1	n/a	0	n/a	0	n/a	0	n/a	0
1997	n/a	2			n/a	0	n/a	0	n/a	0	n/a	0
1998			0	1	0	0	0	0	4	0		0
1999	0	7			0	0	3	0	6	0	4	0
2000			0	1	0	1	0	0	4	0	2	0
2001	0	5			0	0	3	0	2	0	2	0
2002			0	1	0	4	3	0	4	0	2	0
2003	0	7			0	3	2	0	1	0	2	0
2004			0	0	0	4	3	0	4	0	6	0
2005	0	2			0	4	0	0	2	0	8	0
2006			0	1	0	3	2	1	4	0	2	0
2007	0	7			0	5	1	1	5	0	6	0
2008			0	3	0	2	2	0	8	0	9	0
2009	0	10			0	2	5	1	3	0	2	0
2010			0	3	0	1	2	1	2	0	6	0
2011	0	7			0	5	1	1	4	0	9	0
2012			1	5	1	5	2	0	4	0	3	0
2013	0	11			0	2	2	2	3	0	7	0
2014			1	3	0	4	4	0	6	0	4	0
2015	0	9			0	5	6	0	6	0	7	0

Table 3.12. Sablefish fork length (cm), weight (kg), and proportion mature by age and sex (weight-at age modeled from 1996-2004 age-length data from the AFSC longline survey).

<u>Age</u>	<u>Fork length (cm)</u>		<u>Weight (kg)</u>		<u>Fraction mature</u>	
	<u>Male</u>	<u>Female</u>	<u>Male</u>	<u>Female</u>	<u>Male</u>	<u>Female</u>
2	48.1	46.8	1.0	0.9	0.059	0.006
3	53.1	53.4	1.5	1.5	0.165	0.024
4	56.8	58.8	1.9	2.1	0.343	0.077
5	59.5	63.0	2.2	2.6	0.543	0.198
6	61.6	66.4	2.5	3.1	0.704	0.394
7	63.2	69.2	2.7	3.5	0.811	0.604
8	64.3	71.4	2.8	3.9	0.876	0.765
9	65.2	73.1	2.9	4.2	0.915	0.865
10	65.8	74.5	3.0	4.4	0.939	0.921
11	66.3	75.7	3.0	4.6	0.954	0.952
12	66.7	76.6	3.1	4.8	0.964	0.969
13	67.0	77.3	3.1	4.9	0.971	0.979
14	67.2	77.9	3.1	5.1	0.976	0.986
15	67.3	78.3	3.1	5.1	0.979	0.99
16	67.4	78.7	3.1	5.2	0.982	0.992
17	67.5	79.0	3.1	5.3	0.984	0.994
18	67.6	79.3	3.2	5.3	0.985	0.995
19	67.6	79.4	3.2	5.3	0.986	0.996
20	67.7	79.6	3.2	5.4	0.987	0.997
21	67.7	79.7	3.2	5.4	0.988	0.997
22	67.7	79.8	3.2	5.4	0.988	0.998
23	67.7	79.9	3.2	5.4	0.989	0.998
24	67.7	80.0	3.2	5.4	0.989	0.998
25	67.7	80.0	3.2	5.4	0.989	0.998
26	67.8	80.1	3.2	5.4	0.999	0.998
27	67.8	80.1	3.2	5.4	0.999	0.999
28	67.8	80.1	3.2	5.4	0.999	0.999
29	67.8	80.1	3.2	5.5	0.999	0.999
30	67.8	80.2	3.2	5.5	0.999	0.999
31+	67.8	80.2	3.2	5.5	1.000	1.000

Table 3.13. Input and output sample sizes and standard deviation of normalized residuals (SDNR) for data sources in the sablefish assessment model.

Multinomial Compositions	Input N/CV	SDNR	Effective N
Domestic LL Fishery Ages	200	1.14	167
Domestic LL Fishery Lengths	120	0.89	389
Trawl Fishery Lengths	50	0.84	94
LL Survey Ages	160	0.86	192
NMFS Trawl Survey Lengths	140	0.97	147
Domestic LL Survey Lengths	20	0.30	225
Japanese/Coop LL Survey Lengths	20	0.32	197
<hr/>			
Lognormal abundance indices			
Domestic RPN	5%	3.83	
Japanese/Coop RPN	5%	3.00	
Domestic Fishery RPW	10%	0.87	
Foreign Fishery RPW	10%	1.30	
NMFS Trawl Survey	10-20%	1.91	

Table 3.14. Sablefish recruits, total biomass (2+), and spawning biomass plus lower and upper lower 95% credible intervals (2.5%, 97.5%) from MCMC. Recruits are in millions, and biomass is in kt.

Year	Recruits (Age 2)			Total Biomass			Spawning Biomass		
	Mean	2.5%	97.5%	Mean	2.5%	97.5%	Mean	2.5%	97.5%
1977	1.6	0	12	294	275	319	130	118	149
1978	2.4	0	11	265	247	288	119	108	135
1979	83.3	65	105	322	303	350	114	104	129
1980	27.7	3	49	356	336	381	109	100	122
1981	8.6	0	31	373	352	398	107	99	120
1982	48.2	28	75	418	398	449	111	103	122
1983	22.6	1	41	446	425	471	123	115	134
1984	42.6	33	58	488	469	515	139	131	151
1985	0.5	0	4	491	472	518	154	146	167
1986	24.0	12	35	502	484	527	169	160	182
1987	18.7	11	29	491	473	515	175	166	189
1988	4.4	0	13	458	442	479	174	165	188
1989	4.5	0	11	415	399	433	167	159	181
1990	5.5	3	10	372	359	389	158	150	171
1991	28.8	24	34	355	342	370	147	139	160
1992	0.3	0	2	325	313	340	136	128	148
1993	25.4	21	30	318	306	334	125	118	137
1994	3.4	0	8	296	284	311	114	107	125
1995	6.4	2	10	275	264	290	106	99	116
1996	7.2	5	10	257	246	270	101	95	110
1997	19.2	16	23	253	241	267	98	92	106
1998	1.2	0	4	238	227	252	95	89	103
1999	30.9	26	36	249	237	263	91	86	99
2000	20.1	14	28	259	246	275	88	83	95
2001	10.0	1	17	259	245	274	85	80	92
2002	44.4	38	54	290	276	308	84	79	91
2003	6.1	1	11	295	279	312	87	81	94
2004	14.8	11	19	299	283	317	90	85	97
2005	6.3	3	10	291	276	309	95	89	102
2006	11.0	7	15	285	270	302	101	95	108
2007	8.1	5	12	275	260	293	106	99	114
2008	10.4	7	14	266	251	282	107	100	115
2009	9.1	6	12	256	242	273	106	99	114
2010	19.8	16	25	259	244	275	104	97	112
2011	3.8	0	7	251	236	267	101	95	109
2012	8.8	5	13	243	228	259	98	92	106
2013	0.3	0	1	226	212	242	95	88	102
2014	2.8	6	26	208	193	223	92	86	100
2015	13.3	5	23	202	185	216	90	80	94
2016	17.2	5	25	189	141	238	86	73	89
2017	17.2	5	36	194	154	233	82	64	87

Table 3.15. Regional estimates of sablefish total biomass (Age 2+). Partitioning was done using RPWs from Japanese LL survey from 1979-1989 and domestic LL survey from 1990-2015 using a 2 year moving average. For 1960-1978, a prospective 4:6:9 - year average of forward proportions was used.

Year	Bering Sea	Aleutian Islands	Western GOA	Central GOA	West Yakutat	EYakutat/Southeast	Alaska
1960	99	118	51	149	46	71	535
1961	102	121	53	153	48	73	549
1962	114	136	59	171	54	82	616
1963	115	137	59	172	54	82	620
1964	114	137	59	172	54	82	618
1965	117	140	61	176	55	84	633
1966	127	152	66	191	60	91	687
1967	128	153	66	192	60	92	691
1968	126	151	66	190	59	91	684
1969	120	143	62	181	56	86	649
1970	110	132	57	166	52	79	597
1971	99	119	52	150	47	72	538
1972	91	109	47	137	43	66	492
1973	82	98	43	124	39	59	445
1974	74	89	39	112	35	53	401
1975	66	80	35	100	31	48	360
1976	62	73	32	93	29	44	333
1977	54	65	28	82	26	39	294
1978	49	60	26	72	23	36	265
1979	61	66	30	96	28	42	322
1980	64	85	34	95	31	47	356
1981	67	93	39	83	35	57	373
1982	76	87	54	101	40	60	418
1983	80	93	69	113	37	54	446
1984	92	113	78	117	35	54	488
1985	101	112	71	122	36	49	491
1986	108	106	68	125	43	53	502
1987	80	107	65	131	49	60	491
1988	48	93	61	147	47	61	458
1989	56	81	48	133	43	54	415
1990	57	61	40	114	43	57	372
1991	39	41	38	112	47	78	355
1992	23	37	25	103	51	86	325
1993	15	35	29	105	54	80	318
1994	18	34	32	97	45	69	296
1995	26	31	28	90	39	62	275
1996	24	26	28	93	33	52	257
1997	24	23	26	98	31	50	253
1998	21	30	27	84	27	49	238
1999	20	41	29	83	27	50	249
2000	20	42	34	86	26	49	259
2001	28	41	41	81	22	45	259
2002	40	44	43	94	24	45	290
2003	40	45	41	100	26	43	295
2004	40	46	37	106	28	43	299
2005	42	44	38	94	26	47	291
2006	45	40	40	86	26	49	285
2007	48	35	29	85	29	49	275
2008	51	34	26	83	26	46	266
2009	49	33	30	80	23	41	256
2010	51	29	27	76	29	48	259
2011	33	25	25	89	32	46	251
2012	13	31	28	97	27	46	243
2013	30	31	23	76	21	46	226
2014	44	26	22	59	18	39	208
2015	35	27	22	58	22	38	202

Table 3.16. Key parameter estimates and their uncertainty and Bayesian credible intervals (BCI). Recruitment is in millions.

Parameter	μ (MLE)	μ (MCMC)	Median (MCMC)	σ (Hessian)	σ (MCMC)	BCI- Lower	BCI- Upper
$q_{domesticLL}$	7.63	7.63	7.63	0.11	0.22	7.19	8.06
q_{coopLL}	6.21	6.20	6.19	0.11	0.20	5.81	6.59
q_{trawl}	1.35	1.35	1.35	0.31	0.09	1.18	1.54
$F_{40\%}$	0.09	0.11	0.10	0.023	0.030	0.06	0.18
2015 SSB (kt)	90.0	89.9	89.9	4.30	4.88	83.2	97
2000 Year Class	44.4	46.2	46.3	3.91	4.22	38.1	53.6
2008 Year Class	19.8	20.1	20.1	2.16	2.18	15.9	24.5

Table 3.17. Comparison of 2014 results versus 2015 results. Biomass is in kilotons.

Year	2014 SAFE	2015 SAFE	Difference (%)	2014 SAFE	2015 SAFE	Difference (%)
	Spawning Biomass	Spawning Biomass		Total Biomass	Total Biomass	
1977	131	130	-1%	294	294	0%
1978	119	119	0%	264	265	0%
1979	114	114	0%	322	322	0%
1980	109	109	0%	355	356	0%
1981	107	107	0%	373	373	0%
1982	111	111	0%	417	418	0%
1983	123	123	0%	445	446	0%
1984	139	139	0%	488	488	0%
1985	154	154	0%	491	491	0%
1986	168	169	1%	502	502	0%
1987	175	175	0%	491	491	0%
1988	174	174	0%	457	458	0%
1989	167	167	0%	415	415	0%
1990	158	158	0%	373	372	0%
1991	147	147	0%	356	355	0%
1992	136	136	0%	326	325	0%
1993	125	125	0%	320	318	-1%
1994	115	114	-1%	297	296	0%
1995	106	106	0%	277	275	-1%
1996	101	101	0%	259	257	-1%
1997	98	98	0%	254	253	0%
1998	96	95	-1%	240	238	-1%
1999	92	91	-1%	251	249	-1%
2000	89	88	-1%	260	259	0%
2001	86	85	-1%	262	259	-1%
2002	85	84	-1%	292	290	-1%
2003	88	87	-1%	299	295	-1%
2004	91	90	-1%	303	299	-1%
2005	96	95	-1%	295	291	-1%
2006	102	101	-1%	289	285	-1%
2007	107	106	-1%	280	275	-2%
2008	109	107	-2%	271	266	-2%
2009	108	106	-2%	262	256	-2%
2010	106	104	-2%	263	259	-2%
2011	104	101	-3%	254	251	-1%
2012	101	98	-3%	246	243	-1%
2013	98	95	-3%	228	226	-1%
2014	95	92	-3%	218	208	-5%
2015		90			202	

Table 3.18. Sablefish spawning biomass (kilotons), fishing mortality, and yield (kilotons) for seven harvest scenarios. Abundance projected using 1979-2013 recruitments.

Year	Maximum permissible F	Author's F* (specified catch)	Half max. F	5-year average F	No fishing	Overfished?	Approaching overfished?
Spawning biomass (kt)							
2015	90.0	90.0	90.0	90.0	90.0	90.0	90.0
2016	86.5	86.5	86.4	86.5	86.5	86.5	86.5
2017	80.9	82.0	83.8	81.6	87.1	79.8	80.9
2018	76.5	78.5	81.1	77.4	87.9	74.6	76.5
2019	74.9	76.5	79.6	75.7	90.8	72.4	74.0
2020	76.4	77.8	79.7	77.0	96.9	73.5	74.7
2021	80.1	81.2	81.8	80.7	105.9	76.6	77.6
2022	84.4	85.3	84.8	85.4	116.2	80.3	81.1
2023	88.5	89.2	89.6	90.1	126.9	83.8	84.4
2024	92.2	92.8	93.3	94.5	137.2	86.9	87.3
2025	95.3	95.8	98.3	98.5	147.1	89.4	89.7
2026	98.0	98.3	103.6	102.1	156.5	91.5	91.8
2027	100.3	100.6	107.3	105.4	165.3	93.3	93.5
2028	102.3	102.5	110.9	108.4	173.6	94.9	95.0
Fishing mortality							
2015	0.063	0.063	0.063	0.063	0.063	0.063	0.063
2016	0.078	0.065	0.039	0.069	-	0.093	0.093
2017	0.073	0.060	0.038	0.069	-	0.086	0.086
2018	0.069	0.071	0.037	0.069	-	0.080	0.080
2019	0.067	0.069	0.036	0.069	-	0.077	0.077
2020	0.068	0.069	0.036	0.069	-	0.077	0.077
2021	0.069	0.070	0.037	0.069	-	0.079	0.079
2022	0.071	0.071	0.038	0.069	-	0.081	0.081
2023	0.072	0.073	0.041	0.069	-	0.082	0.082
2024	0.074	0.074	0.043	0.069	-	0.084	0.084
2025	0.075	0.075	0.045	0.069	-	0.086	0.086
2026	0.076	0.076	0.047	0.069	-	0.087	0.087
2027	0.078	0.078	0.047	0.069	-	0.089	0.089
2028	0.079	0.079	0.047	0.069	-	0.090	0.090
Yield (kt)							
2015	10.4	10.4	10.4	10.4	10.4	10.4	10.4
2016	11.8	11.8	6.0	10.5	-	13.9	11.8
2017	10.5	10.8	5.7	10.1	-	12.1	10.5
2018	10.4	10.9	6.0	10.5	-	11.7	12.3
2019	11.2	11.6	6.7	11.3	-	12.5	13.0
2020	12.5	12.8	7.6	12.2	-	13.8	14.2
2021	13.6	13.9	8.4	12.9	-	15.1	15.4
2022	14.7	14.9	9.1	13.5	-	16.2	16.4
2023	15.5	15.7	9.7	14.1	-	17.1	17.2
2024	16.3	16.4	10.3	14.6	-	17.8	17.9
2025	17.0	17.0	10.8	15.0	-	18.5	18.6
2026	17.5	17.6	11.3	15.5	-	19.1	19.1
2027	18.0	18.1	11.7	15.8	-	19.6	19.6
2028	18.6	18.6	12.2	16.2	-	20.2	20.2

* Projections in Author's F (Alternative 2) are based on estimated catches of 9,781 t and 8,715 t used in place of maximum permissible ABC for 2016 and 2017. This was done in response to management requests for a more accurate two-year projection.

Table 3.19. Analysis of ecosystem considerations for the sablefish fishery.

<i>Indicator</i>	<i>Observation</i>	<i>Interpretation</i>	<i>Evaluation</i>
<i>ECOSYSTEM EFFECTS ON STOCK</i>			
<i>Prey availability or abundance trends</i>			
Zooplankton	None	None	Unknown
<i>Predator population trends</i>			
Salmon	Decreasing	Increases the stock	No concern
<i>Changes in habitat quality</i>			
Temperature regime	Warm increases recruitment	Variable recruitment	No concern (can't affect)
Prevailing currents	Northerly increases recruitment	Variable recruitment	No concern (can't affect)
<i>FISHERY EFFECTS ON ECOSYSTEM</i>			
<i>Fishery contribution to bycatch</i>			
Prohibited species	Small catches	Minor contribution to mortality	No concern
Forage species	Small catches	Minor contribution to mortality	No concern
HAPC biota (seapens/whips, corals, sponges, anemones)	Small catches, except long-term reductions predicted	Long-term reductions predicted in hard corals and living structure	Possible concern
Marine mammals and birds	Bird catch about 10% total	Appears to be decreasing	Possible concern
Sensitive non-target species	Grenadier, spiny dogfish, and unidentified shark catch notable	Grenadier catch high but stable, recent shark catch is small	Possible concern for grenadiers
<i>Fishery concentration in space and time</i>	IFQ less concentrated	IFQ improves	No concern
<i>Fishery effects on amount of large size target fish</i>	IFQ reduces catch of immature	IFQ improves	No concern
<i>Fishery contribution to discards and offal production</i>	sablefish <5% in longline fishery, but 30% in trawl fishery	IFQ improves, but notable discards in trawl fishery	Trawl fishery discards definite concern
<i>Fishery effects on age-at-maturity and fecundity</i>	trawl fishery catches smaller fish, but only small part of total catch	slightly decreases	No concern

Figures





Figure 3.1. Long term and short term sablefish catch by gear type.

Catch by FMP management area




Figure 3.2. Sablefish fishery total reported catch (kt) by North Pacific Fishery Management Council area and year.




Figure 3.3. Observed and predicted sablefish relative population weight and numbers versus year. Points are observed estimates with approximate 95% confidence intervals. Solid red line is model predicted. The relative population weights are not fit in the models, but are presented for comparison.






Figure 3.4. Observed and predicted sablefish abundance indices. Fishery indices are on top two panels. GOA trawl survey is on the bottom left panel. Points are observed estimates with approximate 95% confidence intervals while solid red lines are model predictions.




Figure 3.5. Average fishery catch rate (pounds/hook) by region and data source for longline survey and fishery data. The fishery switched from open-access to individual quota management in 1995. Data is not presented for years when there were fewer than three vessels. This occurred in observer data in the Bering Sea in 1994, 1997, 2003, and 2012, in logbook data in the Bering Sea in 2002, and in East Yakutat observer data from 1990-1994.




Figure 3.5. (continued)




Figure 3.6. Average fishery catch rate (pounds/hook) and associated 95% confidence intervals by region and data source. The fishery switched from open-access to individual quota management in 1995. Data is not presented for years when there were fewer than three vessels. This occurred in observer data in the Bering Sea in 1994, 1997, 2003, and 2012, in logbook data in the Bering Sea in 2002, and in East Yakutat observer data from 1990-1994.




Figure 3.6. (continued)




Figure 3.7. Relative abundance (numbers) by region and survey. The regions Bering Sea, Aleutians Islands, and western Gulf of Alaska are combined in the first plot. The two surveys are the Japan-U.S. cooperative longline survey and the domestic (U.S.) longline survey. In this plot, the values for the U.S. survey were adjusted to account for the higher efficiency of the U.S. survey gear.




Figure 3.8 Comparison of abundance trends in GOA gully stations versus GOA slope stations.




Figure 3.9. NMFS Bering Sea Slope and Aleutian Island trawl survey biomass estimates. Bering Sea Slope years are jittered so that intervals do not overlap.




Figure 3.10a. Comparisons of IPHC and AFSC longline survey trends in relative population number of sablefish in the Gulf of Alaska.




Figure 3.10b. Comparisons of IPHC and AFSC trawl survey trends abundance of sablefish in the Gulf of Alaska. Years in which both surveys occurred have a correlation coefficient of $r = 0.9$.




Figure 3.11a. Northern Southeast Inside (NSEI) sablefish longline survey catch-per-unit-effort (CPUE) in round pounds/hook and commercial catch from 1980 to 2014. A three-hour minimum soak time was used on the NSEI sablefish longline survey (from K. Green. ADF&G, pers. comm. October, 2015).




Figure 3.11b. Northern Southeast Inside (NSEI) commercial sablefish longline catch-per-unit-effort (CPUE) in round pounds-per-hook from 1997 – 2014 and commercial catch from 1980 – 2014 (from K. Green pers. comm., ADF&G, October, 2015).




Figure 3.12. Age-length conversion matrices for sablefish. Top panels are female, bottom panel are males, left is 1960-1995, and right is 1996-2015.





Figure 3.13.--Estimated sablefish total biomass (thousands t) and spawning biomass (bottom) with 95% MCMC credible intervals.




Figure 3.14a. Estimated recruitment by year class 1958-2011 (number at age 2, millions) for 2014 and 2015 models.




Figure 3.14b. Estimates of the number of age-2 sablefish (millions) with 95% credible intervals by year class. Credible intervals are based on MCMC posterior.




Figure 3.15. Relative contribution of the last 20 year classes to next year's female spawning biomass.




Figure 3.16. Gulf of Alaska bottom trawl survey length (cm) compositions for female sablefish at depths <500 m. Bars are observed frequencies and lines are predicted frequencies.




Figure 3.17. Gulf of Alaska bottom trawl survey length (cm) compositions for male sablefish at depths <500 m. Bars are observed frequencies and lines are predicted frequencies.

Top 4 year classes by Survey and Area




Figure 3.18. Above average 1995, 1997, 2000 and potential above-average 2008 year classes relative population abundance in each survey year and area.




Figure 3.19. Domestic longline survey age compositions. Bars are observed frequencies and lines are predicted frequencies.




Figure 3.19 (cont.). Domestic longline survey age compositions. Bars are observed frequencies and lines are predicted frequencies.




Figure 3.19 (cont.). Domestic longline survey age compositions. Bars are observed frequencies and lines are predicted frequencies.




Figure 3.20. Relative abundance (number in thousands) by age and region from the domestic (U.S.) longline survey. The regions Bering Sea, Aleutian Islands, and Western Gulf of Alaska are combined.




Figure 3.20 (cont.). Relative abundance (number in thousands) by age and region from the domestic (U.S.) longline survey. The regions Bering Sea, Aleutian Islands, and Western Gulf of Alaska are combined.




Figure 3.20 (cont.). Relative abundance (number in thousands) by age and region from the domestic (U.S.) longline survey. The regions Bering Sea, Aleutian Islands, and Western Gulf of Alaska are combined.




Figure 3.21. Japanese longline survey age compositions. Bars are observed frequencies and line is predicted frequencies.




Figure 3.22. Domestic fixed gear fishery length (cm) compositions for females. Bars are observed frequencies and lines are predicted frequencies.




Figure 3.22 (cont.). Domestic fixed gear fishery length (cm) compositions for females. Bars are observed frequencies and lines are predicted frequencies.




Figure 3.23. Domestic fixed gear fishery length (cm) compositions for males. Bars are observed frequencies and lines are predicted frequencies.




Figure 3.23 (cont.). Domestic fixed gear fishery length (cm) compositions for males. Bars are observed frequencies and lines are predicted frequencies.




Figure 3.24. Domestic fishery age compositions. Bars are observed frequencies and lines are predicted frequencies.




Figure 3.24 (cont.). Domestic fishery age compositions. Bars are observed frequencies and lines are predicted frequencies.




Figure 3.25a. Domestic trawl gear fishery length (cm) compositions for females. Bars are observed frequencies and lines are predicted frequencies.




Figure 3.25b. Domestic trawl gear fishery length (cm) compositions for males. Bars are observed frequencies and lines are predicted frequencies.




Figure 3.26. Domestic longline survey length (cm) compositions for females. Bars are observed frequencies and lines are predicted frequencies.




Figure 3.26 (cont.). Domestic longline survey length (cm) compositions for females. Bars are observed frequencies and lines are predicted frequencies.




Figure 3.27. Domestic longline survey length (cm) compositions for males. Bars are observed frequencies and lines are predicted frequencies.




Figure 3.27.(cont.). Domestic longline survey length (cm) compositions for males. Bars are observed frequencies and lines are predicted frequencies.




Figure 3.28. Sablefish selectivities for fisheries. The derby longline occurred until 1994 when the fishery switched to IFQ in 1995.









Figure 3.28 (cont.). Sablefish selectivities for surveys.




Figure 3.29. Time series of combined fully-selected fishing mortality for fixed and trawl gear for sablefish.




Figure 3.30. Phase-plane diagram of time series of sablefish estimated spawning biomass relative to the unfished level and fishing mortality relative to F_{OFL} for author recommended model. Bottom is zoomed in to examine more recent years.




Figure 3.31a. Retrospective trends for spawning biomass (top) and percent difference from terminal year (bottom) from 2005-2015. Mohn's revised $\rho = 0.023$.




Figure 3.31b. Retrospective trends for spawning biomass (top) and percent difference from terminal year (bottom) from 2005-2015 with MCMC credible intervals per year. Mohn's revised $\rho = 0.023$.




Figure 3.31c. Squid plot of the development of initial estimates of age-2 recruitment since year class 2002 through year class 2012 from retrospective analysis. Number to right of terminal year indicates year class.




Figure 3.32. Posterior probability distribution for projected spawning biomass (thousands t) in 2016.




Figure 3.33. Pairwise scatterplots of key parameter MCMC runs. Red curve is loess smooth. Numbers in upper right hand panel are correlation coefficients between parameters.




Figure 3.34. Probability that projected spawning biomass (from MCMC) will fall below $B_{40\%}$, $B_{35\%}$ and $B_{17.5\%}$.




Figure 3.35. Estimates of female spawning biomass (thousands t) and their uncertainty. White line is the median and green line is the mean, shaded fills are 5% increments of the posterior probability distribution of spawning biomass based on 10,000,000 MCMC simulations. Width of shaded area is the 95% credibility interval. Harvest policy is the same as the projections in Scenario 2 (Author's F).




Figure 3.36. (A) The mean relative change in apportionment percentages across areas from 2007-2014. (B) The relative change in the apportionment share for the Bering Sea from 2007-2014. (C) The mean change in ABC for each area from 2007-2014.

Appendix 3A.--Sablefish longline survey - fishery interactions

NMFS has requested the assistance of the fishing fleet to avoid the annual sablefish longline survey since the inception of sablefish IFQ management in 1995. We requested that fishermen stay at least five nautical miles away from each survey station for 7 days before and 3 days after the planned sampling date (3 days allow for survey delays). Beginning in 1998, we also revised the longline survey schedule to avoid the July 1 rockfish trawl fishery opening as well as other short, but less intense fisheries.

History of interactions

Publicity, the revised longline survey schedule, and fishermen cooperation generally have been effective at reducing fishery interactions. Distribution of the survey schedule to all IFQ permit holders, radio announcements from the survey vessel, and the threat of a regulatory rolling closure have had intermittent success at reducing the annual number of longline fishery interactions.

Since 2000, the number of vessels fishing near survey stations has remained relatively low. During the past several surveys, many fishing vessels were contacted by the survey vessel and in most cases fishermen were aware of the survey or willing to help out by fishing other grounds to avoid potential survey interactions.

Longline Survey-Fishery Interactions

Year	<u>Longline</u>		<u>Trawl</u>		<u>Pot</u>		<u>Total</u>	
	Stations	Vessels	Stations	Vessels	Stations	Vessels	Stations	Vessels
1995	8	7	9	15	0	0	17	22
1996	11	18	15	17	0	0	26	35
1997	8	8	8	7	0	0	16	15
1998	10	9	0	0	0	0	10	9
1999	4	4	2	6	0	0	6	10
2000	10	10	0	0	0	0	10	10
2001	1	1	1	1	0	0	2	2
2002	3	3	0	0	0	0	3	3
2003	4	4	2	2	0	0	6	6
2004	5	5	0	0	1	1	6	6
2005	1	1	1	1	0	0	2	2
2006	6	6	1	2	0	0	7	8
2007	8	6	2	2	0	0	10	8
2008	2	2	2	2	0	0	4	4
2009	3	3	0	0	0	0	3	3
2010	2	2	1	1	0	0	3	3
2011	3	3	0	0	0	0	3	3
2012	5	5	0	0	0	0	5	5
2013	5	5	0	0	0	0	5	5
2014	2	2	0	0	0	0	2	2
2015	3	3	1	1			4	4

Recommendation

We have followed several practical measures to alleviate fishery interactions with the survey. Trawl fishery interactions generally have decreased; longline fishery interactions have been low but continue to occur. Discussions with vessels encountered on the survey indicated an increasing level of “hired” skippers who are unaware of the survey schedule. Publicizing the survey schedule to skippers who aren’t

quota shareholders should be improved. We will continue to work with association representatives and individual fishermen from the longline and trawl fleets to reduce fishery interactions and ensure accurate estimates of sablefish abundance.

Appendix 3B.—Supplemental catch data

In order to comply with the Annual Catch Limit (ACL) requirements, two new datasets have been generated to help estimate total catch and removals from NMFS stocks in Alaska.

The first dataset, non-commercial removals, estimates total removals that do not occur during directed groundfish fishing activities. This includes removals incurred during research, subsistence, personal use, recreational, and exempted fishing permit activities, but does not include removals taken in fisheries other than those managed under the groundfish FMP. These estimates represent additional sources of removals to the existing Catch Accounting System estimates. For sablefish, these estimates can be compared to the research removals reported in previous assessments (Hanselman et al. 2010) (Table 3B.1). The sablefish research removals are substantial relative to the fishery catch and compared to the research removals for many other species. These research removals support a dedicated longline survey. Additional sources of significant removals are bottom trawl surveys and the International Pacific Halibut Commissions longline survey. Recreational removals are relatively minor for sablefish. Total removals from activities other than directed fishery has ranged from 239-359 t in recent years. This represents ~1.5 – 2.5 percent of the recommended ABC annually. These removals represent a relatively low risk to the sablefish stock. When an assessment model is fit that includes these removals as part of the total catch, the result is an increase in ABC of comparable magnitude.

The second dataset, Halibut Fishery Incidental Catch Estimation (HFICE), is an estimate of the incidental catch of groundfish in the halibut IFQ fishery in Alaska, which is currently unobserved. To estimate removals in the Pacific halibut fishery, methods were developed by the HFICE working group and approved by the Gulf of Alaska and Bering Sea/Aleutian Islands Plan Teams and the Scientific and Statistical Committee of the North Pacific Fishery Management Council. A detailed description of the methods is available in Tribuzio et al. (2011).

These estimates are for total catch of groundfish species in the halibut IFQ fishery and do not distinguish between “retained” or “discarded” catch. These estimates should be considered a separate time series from the current CAS estimates of total catch. Because of potential overlaps HFICE removals should not be added to the CAS produced catch estimates. The overlap will apply when groundfish are retained or discarded during an IFQ halibut trip. IFQ halibut landings that also include landed groundfish are recorded as retained in eLandings and a discard amount for all groundfish is estimated for such landings in CAS. Discard amounts for groundfish are not currently estimated for IFQ halibut landings that do not also include landed groundfish. For example, catch information for a trip that includes both landed IFQ halibut and sablefish would contain the total amount of sablefish landed (reported in eLandings) and an estimate of discard based on at-sea observer information. Further, because a groundfish species was landed during the trip, catch accounting would also estimate discard for all groundfish species based on available observer information and following methods described in Cahalan et al. (2010). The HFICE method estimates all groundfish caught during a halibut IFQ trip and thus is an estimate of groundfish caught whether landed or discarded. This prevents simply adding the CAS total with the HFICE estimate because it would be analogous to counting both retained and discarded groundfish species twice. Further, there are situations where the HFICE estimate includes groundfish caught in State waters and this would need to be considered with respect to ACLs (e.g. Chatham Strait sablefish fisheries). Therefore, the HFICE estimates should be considered preliminary estimates for what is caught in the IFQ halibut fishery. With restructuring of the Observer Program improved estimates of groundfish catch in the halibut fishery began in 2013. More years of data are needed for an evaluation the effects of observer

restructuring on catch of sablefish in the halibut IFQ fishery..

The HFICE estimates of sablefish catch by the halibut fishery are substantial and represent approximately 10% of the annual sablefish ABC (Table 3B.2). Sablefish and halibut are often caught and landed in association with each other by the IFQ fishery. It is unknown what level of sablefish catch reported here is already accounted for as IFQ harvest in the CAS system because the HFICE estimates do not separate retained and discarded catch. If these were strictly additive removals, 10% would represent a significant amount of additional mortality and a potential risk to the stock, but how much is additive is unknown. The HFICE estimates may represent some valuable discard information for sablefish, but that level is unknown until these estimates are separated from the IFQ landings and CAS system.

Literature Cited

- Cahalan J., J. Mondragon., and J. Gasper. 2010. Catch Sampling and Estimation in the Federal Groundfish Fisheries off Alaska. NOAA Technical Memorandum NMFS-AFSC-205. 42 p.
- Hanselman, D. H., C. Lunsford, and C. Rodgveller. 2010. Alaskan Sablefish. *In* Stock assessment and fishery evaluation report for the groundfish resources of the GOA and BS/AI as projected for 2010. North Pacific Fishery Management Council, 605 W 4th Ave, Suite 306 Anchorage, AK 99501.pp.
- Tribuzio, C.A., S. Gaichas, J. Gasper, H. Gilroy, T. Kong, O. Ormseth, J. Cahalan, J. DiCosimo, M. Furuness, H. Shen, and K. Green. 2011. Methods for the estimation of non-target species catch in the unobserved halibut IFQ fleet. August Plan Team document. Presented to the Joint Plan Teams of the North Pacific Fishery Management Council.

Table 3B.1 Total removals of sablefish (t) from activities not related to directed fishing, since 1977. Trawl survey sources are a combination of the NMFS echo-integration, small-mesh, GOA, AI, and BS Slope bottom trawl surveys, and occasional short-term research projects. Other is recreational, personal use, and subsistence harvest.

Year	Source	Trawl surveys	Japan US longline survey	Domestic longline survey	IPHC longline survey*	Other	Total
1977		3					3
1978		14					14
1979		27	104				131
1980		70	114				184
1981		88	150				238
1982		108	240				348
1983		46	236				282
1984		127	284				412
1985		186	390				576
1986		123	396				519
1987		117	349				466
1988		15	389	303			707
1989		4	393	367			763
1990		26	272	366			664
1991		3	255	386			645
1992		0	281	393			674
1993		39	281	408			728
1994		1	271	395			667
1995		0		386			386
1996		13		430			443
1997		1		396			397
1998		26		325	50		401
1999		43		311	49		403
2000		2		290	53		345
2001		11		326	48		386
2002		3		309	58		370
2003		16		280	98		393
2004		2		288	98		387
2005		18		255	92		365
2006	Assessment of the sablefish stock in Alaska (Hanselman et al. 2010)	2		287	64		352
2007		17		266	48		331
2008		3		262	46		310
2009		14		242	47		257
2010		3		291	50	15	359
2011		9		273	39	16	312
2012		4		203	27	39	273
2013		4		178	22	35	239
2014	AKRO	<1		197	32	29	258

* IPHC survey sablefish removals are released and estimates from mark-recapture studies suggest that these removals are expected to produce low mortality. Some state removals are included.

Table 3B.2. Estimates of Alaska sablefish catch (t) from the Halibut Fishery Incidental Catch Estimation (HFICE) working group. AI = Aleutian Islands, WGOA = Western Gulf of Alaska, CGOA = Central Gulf of Alaska, EGOA = Eastern Gulf of Alaska, PWS = Prince William Sound.

<u>Area</u>	<u>2001</u>	<u>2002</u>	<u>2003</u>	<u>2004</u>	<u>2005</u>	<u>2006</u>	<u>2007</u>	<u>2008</u>	<u>2009</u>	<u>2010</u>
Western/Central AI	27	19	34	18	14	11	36	44	17	23
Eastern AI	18	16	46	26	20	6	4	13	6	7
WGOA	10	9	12	22	21	16	7	12	3	12
CGOA-Shumagin	184	27	36	65	60	47	21	38	10	37
CGOA-Kodiak/ PWS*	802	107	96	89	82	49	57	33	69	63
EGOA-Yakutat	110	324	291	258	240	149	175	103	207	195
EGOA-Southeast	339	335	389	315	269	242	230	184	242	262
Southeast Inside*	459	1,018	1,181	917	786	739	701	574	731	805
Total	1,948	2,231	2,346	2,469	2,194	2,476	1,937	1,874	1,921	1,594

*These areas include removals from the state of Alaska.