

# Assessment of the yellowfin sole stock in the Bering Sea and Aleutian Islands

Thomas K. Wilderbuer, Daniel G. Nichol and James Ianelli

## Executive Summary

### Summary of Changes in Assessment Inputs

#### *Changes to the input data*

- 1) 2016 fishery age composition.
- 2) 2016 survey age composition.
- 3) 2017 trawl survey biomass point estimate and standard error.
- 4) Estimate of the discarded and retained portions of the 2015 catch.
- 5) Estimate of total catch made through the end of 2017. Catch of 150,000 t assumed for 2018 and 2019 projection.

#### *Changes to the assessment methodology*

No changes were made to this assessment.

## Summary of Results

The assessment updates last year's with results and management quantities that are higher than the 2016 assessment primarily due to 1) including 2016 fishery weight-at-age information, 2) including two more years in the spawner-recruit time-series, and 3) a lower estimated survey catchability. Yellowfin sole continue to be well-above  $B_{MSY}$  and the annual harvest remains below the ABC level. The female spawning stock is in a slow downward trend.

Quantity	As estimated or specified last year for:		As estimated or recommended this year for:	
	2017	2018	2018	2019
$M$ (natural mortality rate)	0.12	0.12	0.12	0.12
Tier	1a	1a	1a	1a
Projected total (age 6+) biomass (t)	2,290,000	2,202,300	2,553,100	2,460,700
Female spawning biomass (t)				
Projected	778,600	770,900	895,000	890,000
$B_0$	1,202,700		1,204,000	
$B_{MSY}$	424,000		456,000	
$F_{OFL}$	0.125	0.125	0.12	0.12
$\max F_{ABC}$	0.114	0.114	0.109	0.109
$F_{ABC}$	0.114	0.114	0.109	0.109
OFL (t)	287,000	276,000	306,700	295,600
maxABC (t)	260,800	250,800	277,500	267,500
ABC (t)	260,800	250,800	277,500	267,500
Status	As determined last year for:		As determined this year for:	
	2015	2016	2016	2017
Overfishing	No	n/a	No	n/a
Overfished	n/a	No	n/a	No
Approaching overfished	n/a	No	n/a	No

Projections are based on estimated catches of 150,000 t used in place of maximum ABC for 2018 and 2019.

## **Responses to SSC and Plan Team Comments on Assessments in General**

### ***General comments for all assessments:***

**SSC encourages assessment authors to adopt the current model naming convention.**

This assessment now complies with the desired naming convention.

## **Responses to SSC and Plan Team Comments Specific to this Assessment**

The yellowfin sole stock assessment has set an example for its inclusion of ecosystem factors in the assessment. Temperature was found to be related to survey catchability and growth. Interestingly, after conducting a field experiment to further examine relationships between temperature and catchability, it was found that survey biomass is more strongly correlated to wave height than temperature, which is in turn confounded with temperature. The SSC supports the approach outlined in the SAFE to further elucidate the effects of sea state and/or bottom temperature on  $q$ , noting that these covariates may act on the assessment in different ways (i.e., availability versus gear efficiency). If an effect of wave height on catchability is confirmed, implying that rougher seas adversely affect the ability of the trawl to tend the seafloor, then this would beg the question whether other assessed species are similarly affected. So, the outcome of this work has the potential for important, far-reaching implications.

The results of last summer's sampling efficiency experiment indicated that sea-state had a much larger effect on yellowfin sole herding, and trawl efficiency in general, than bottom temperature. Dan Nichol looked at the time-series of eastern Bering Sea yellowfin sole biomass estimates and found that they also were strongly correlated to wave height. To confuse things more, wave height was correlated to temperature.

1. The experimental results were analyzed and a paper on wave height effect on yellowfin sole catchability was submitted to Fisheries Bulletin (*The effects of wave-induced vessel motion on the geometry of a bottom survey trawl and the herding of yellowfin sole by Somerton, Weinberg, Munro, Rugolo and Wilderbuer, In Review*). However, this still does not answer the question of why you find a strong temperature effect on the survey catchability function in the stock assessment model. One possibility is that trawl sampling efficiency is more influenced by waves but yellowfin sole availability to the survey is more influenced by temperature.
2. Dan Nichol has completed a CPUE versus waves analysis back to 2005. Although RACE surveys have recorded sea state data for many years it was not entered into the database earlier than 2005, so Dan will do some hand entry, then repeat the analysis with more years and considering wave height and temperature together in a model. This work is viewed as secondary to the availability research (see #3), given that availability likely has a much greater influence on biomass estimates than sea-state/temperature.
3. Dan Nichol also has made progress toward showing that yellowfin sole availability changes with temperature using sex ratio, based on the notion that males arrive first and leave the spawning areas later than females, and that the timing of annual spawning is temperature-dependent.
4. A CIE review is scheduled for May 2018 for yellowfin sole, northern rock sole and Alaska plaice. We will solicit comments from the expert panel on exploring an additional waves parameter to fit the survey biomass.

**One ongoing issue with the yellowfin sole assessment is a strong retrospective pattern illustrated by Figure 4.21. More recent assessments tend to yield higher estimates of female spawning biomass. Attempts to determine the cause have been unsuccessful so far. The PT suggested that the author consider examining the potential effects of q and M on the retrospective patterns. The SSC encourages ongoing efforts to understand this phenomenon so that appropriate model adjustments can be made. The SSC also recommends updating weight-at-age with each assessment as new data become available annually. Finally, the SSC looks forward to the use of the adopted model naming convention in next year's yellowfin sole assessment.**

The retrospective comment is addressed in this assessment under the *retrospective analysis* heading. Weight-at-age is routinely updated, both for the fishery and the survey.

## **Introduction**

The yellowfin sole (*Limanda aspera*) is one of the most abundant flatfish species in the eastern Bering Sea (EBS) and currently is the target of the largest flatfish fishery in the world. They inhabit the EBS shelf and are considered one stock. Abundance in the Aleutian Islands region is negligible.

Yellowfin sole are distributed in North American waters from off British Columbia, Canada, (approx. lat. 49° N) to the Chukchi Sea (about lat. 70° N) and south along the Asian coast to about lat. 35° N off the South Korean coast in the Sea of Japan. Adults exhibit a benthic lifestyle and occupy separate winter, spawning and summertime feeding distributions on the eastern Bering Sea shelf. From over-winter grounds near the shelf margins, adults begin a migration onto the inner shelf in April or early May each year for spawning and feeding. The directed fishery has typically occurred from winter through autumn (Wilderbuer et al. 1992). Yellowfin sole are managed as a single stock in the BSAI management area as there is presently no evidence of stock structure.

## **Fishery**

Yellowfin sole have annually been caught with bottom trawls on the Bering Sea shelf since the fishery began in 1954 and were overexploited by foreign fisheries in 1959-62 when catches averaged 404,000 t annually (Fig. 4.1, top panel). As a result of reduced stock abundance, catches declined to an annual average of 117,800 t from 1963-71 and further declined to an annual average of 50,700 t from 1972-77. The lower yield in this latter period was partially due to the discontinuation of the U.S.S.R. fishery. In the early 1980s, after the stock condition had improved, catches again increased reaching a peak of over 227,000 t in 1985.

During the 1980s, there was also a major transition in the characteristics of the fishery. Yellowfin sole were traditionally taken exclusively by foreign fisheries and these fisheries continued to dominate through 1984. However, U.S. fisheries developed rapidly during the 1980s in the form of joint ventures, and during the last half of the decade began to dominate and then take all of the catch as the foreign fisheries were phased out of the EBS. Since 1990, only domestic harvesting and processing has occurred.

The management of the yellowfin sole fishery changed significantly in 2008 with the implementation of Amendment 80 to the BSAI Fisheries Management Plan. The Amendment directly allocated fishery resources among BSAI trawl harvesters in consideration of their historic harvest patterns and future harvest needs in order to improve retention and utilization of fishery resources by the non-AFA trawl catcher/processor fleet. This was accomplished by extending the groundfish retention standards to all H&G vessels and also by providing the ability to form cooperatives within the newly formed Amendment 80 sector. In addition, Amendment 80 also mandated additional monitoring requirements which included observer coverage on all hauls, motion-compensating scales for weighing samples, flow scales to obtain accurate catch weight estimates for the entire catch, no mixing of hauls and no on-deck sorting. The partitioning of TAC and PSC (prohibited species catch) among cooperatives has significantly changed the

way the annual catch has accumulated (Fig 4.1, bottom panel) and the rate of target catch per bycatch ton. There is now a more even and slow attainment of the annual catch relative to the pre-Amendment 80 fishing behavior.

Yellowfin sole are usually headed and gutted, frozen at sea, and then shipped to Asian countries for further processing (AFSC 2016). The first wholesale value of Alaska yellowfin sole totaled \$97.8 million in 2014. In 2010, following a comprehensive assessment process, the yellowfin sole fishery was certified under the Marine Stewardship Council environmental standard for sustainable and well-managed fisheries. The certification also applies to all the major flatfish fisheries in the BSAI and GOA. The total annual catch (t) since implementation of the MFCMA in 1977 is shown in Table 4.1.

In 2011, federally permitted vessels using non-pelagic trawl gear whose harvest results in flatfish retained catch that is greater than any other retained fishery category were required to use modified trawl gear. The modifications required the use of elevating devices to raise the section of the trawl warps between the doors and the trawl wing tips by 2.5 inches off the seafloor. The purpose of the management action was to reduce damage of non-target animals, particularly those that form habitat structure or support other fisheries while not substantially reducing flatfish catch rates or causing gear handling problems (Rose et al. 2010).

The 1997 catch of 181,389 t (retained and discards) was the largest since the fishery became completely domestic but was at lower levels from 1998 – 2010, averaging 94,004 t (Table 4.2). From 2011-2014 the catch increased, averaging 155,000 t. The 2013 catch totaled 165,000 t (73% of the ABC), the highest annual catch in the past 19 years. For 2017, the catch distribution has been spread out from January through May with the majority coming from 4 BSAI management areas (509, 513, 514, 516). As of mid-October 2017, the fishing season is ongoing. In order to estimate the total 2017 catch for the stock assessment model, the average proportion of the 2010-2016 cumulative catch attained by the 37<sup>th</sup> week of the year (mid-September) was applied to the 2017 catch amount at the same time period and results in a 2017 catch estimate of 143,100 t (55% of the ABC). The size composition of the 2017 catch for both males and females, from observer sampling, are shown in Figure 4.2, the catch proportions by month and area are shown in Figure 4.3, and maps of the locations where yellowfin sole were caught in 2017, by month (through mid-September), are shown in Figure 4.4. The average age of yellowfin sole in the 2016 catch is estimated at 12.9 and 12.3 years for females and males, respectively.

The time-series of catch in Table 4.1 also includes yellowfin sole that were discarded in domestic fisheries during the period 1987 to the present. Annual discard estimates were calculated from at-sea sampling (Table 4.2). The rate of discard has ranged from a low of 2% of the total catch in 2012 (and 2015) to 30% in 1992. The trend has been toward fuller retention of the catch in recent years, and with the advent of the Amendment 80 harvest practices, discarding is at its lowest level since these estimates have become available. Historically, discarding primarily occurred in the yellowfin sole directed fishery, with lesser amounts in the Pacific cod, Pollock, rock sole, flathead sole, and “other flatfish” fisheries (Table 4.3).

## Data

The data used in this assessment include estimates of total catch, bottom trawl survey biomass estimates and their attendant 95% confidence intervals, catch-at-age from the fishery, and population age composition estimates from the bottom trawl survey. Weight-at-age and proportion mature-at-age are also available from studies conducted during the bottom trawl surveys.

Data source	Years
Fishery catch	1954-2017
Fishery age composition	1964-2016
Fishery weight-at-age	Avg wt at age from 2008-16 used for 2008-2016
Survey biomass and standard error, bottom temperature	1982-2017
Survey age composition	1979-2016
Annual length-at-age and weight-at-age from surveys	1979-2016
Maturity at age	Combined 1992 and 2012 samples

#### *Fishery Catch and Catch-at-Age*

This assessment uses fishery catch data from 1955- 2017 (shown for 1964-2017 in Table 4.1), including an estimate of the 2017 catch, and fishery catch-at-age (proportions) from 1964-2016 (Table 4.4, 1975-2016). The 2016 fishery age composition was primarily composed of fish older than 9 years with a large amount of 20+ fish.

#### *Survey Biomass Estimates and Population Age Composition Estimates*

Indices of relative abundance available from AFSC surveys have shown a major increase in the abundance of yellowfin sole during the late 1970s, increasing from 21 kg/ha in 1975 to 51 kg/ha in 1981 (Fig. 4.2 in Bakkala and Wilderbuer 1990). These increases have also been documented through Japanese commercial pair trawl data and catch-at-age modeling in past assessments (Bakkala and Wilderbuer 1990).

Since 1981, the survey CPUEs have fluctuated widely (Fig. 4.5). Biomass estimates for yellowfin sole from the annual bottom trawl survey on the eastern Bering Sea shelf are shown in Table 4.5 and Figure 4.6. The data show a doubling of survey biomass between 1975 and 1979 with a further increase to over 3.3 million t in 1981. Total survey abundance estimates fluctuated erratically from 1983 to 1990 with biomass ranging from as high as 3.5 million t in 1983 to as low as 1.9 million t in 1986. Biomass estimates since 1990 indicate an even trend at high levels of abundance for yellowfin sole, with the exception of the results from the 1999 and 2000 summer surveys, which were at lower levels. Surveys from 2001-2005 estimated an increase each year but the estimates since 2006 indicate a stable level with some annual variability. However, the 2012 estimate is a 19% decrease from 2011 and the 2013 and 2014 surveys have estimated a 17% increase over 2012. Similarly, there was a 24% decrease from 2014 to 2015 followed by a 48% increase from 2015 to 2016, the highest biomass estimate since 1984.

Fluctuations of the magnitude shown between 1980 and 1990, 1998 and 1999, 2008 and 2009, 2011and 2012, 2014 and 2015 and 2016 are unreasonable considering the elements of slow growth and long life span of yellowfin sole combined with low to moderate exploitation rate, characteristics which should produce more gradual changes in abundance.

Variability of yellowfin sole survey biomass estimates (Fig. 4.6) is in part due to the availability of yellowfin sole to the survey area (Nichol, 1998). Yellowfin sole are known to undergo annual migrations from wintering areas off the shelf-slope break to near shore waters where they spawn throughout the spring and summer months (Nichol, 1995; Wakabayashi, 1989; Wilderbuer et al., 1992). Exploratory survey sampling in coastal waters of the eastern Bering Sea during early summer indicate that yellowfin sole concentrations can be greater in these shallower areas not covered by the standard AFSC survey than in the survey proper. Commercial bottom trawlers have commonly found high concentrations of yellowfin sole in areas such as near Togiak Bay (Low and Narita, 1990) and in more recent years from Kuskokwim Bay to just south of Nunivak Island. The coastal areas are sufficiently large enough to offer a substantial refuge for yellowfin sole from the current survey.

Over the past 18 years, survey biomass estimates for yellowfin sole have shown a positive correlation with shelf bottom temperatures (Nichol, 1998); estimates have generally been lower during cold years. The 1999 survey, which was conducted in exceptionally cold waters, indicated a decline in biomass that was unrealistic. The bottom temperatures during the 2000 survey were much warmer than in 1999, and the biomass increased, but still did not approach estimates from earlier years. Average bottom temperature and biomass both increased again during the period 2001 – 2003, with the 2003 value the highest temperature and biomass observed over the 22 year time series up to that time. Given that both the 1999 and 2000 surveys were conducted two weeks earlier than previous surveys, it is possible that the time difference may also have also affected the availability of yellowfin sole to the survey. If, for example, the timing of peak yellowfin sole spawning in nearshore waters corresponded to the time of the survey, a greater proportion of the population would be unavailable to the standard survey area. This pattern was observed again in 2009 and 2012 when the temperatures and the bottom trawl survey point estimates were lower. Summer shelf bottom temperatures in 2012 were the 2<sup>nd</sup> coldest recorded by the survey and the time-series and resulted in a 19% decline from 2011. In 2016 the Bering Sea had the highest recorded bottom temperature since measurements began in 1982 and the 2016 estimate of biomass was the highest in 32 years and 48% higher than the 2015 estimate. The 2017 survey estimate of 2,787,700 t was 3% lower than 2016.

We propose two possible reasons why survey biomass estimates are lower during years when bottom temperatures are low. First, catchability may be lower because yellowfin sole may be less active when cold. Less active fish may be less susceptible to herding, and escapement under the footrope of survey gear may increase if fish are less active. Secondly, bottom temperatures may influence the timing of the inshore spawning migrations of yellowfin sole and therefore affect their availability to the survey area. Because yellowfin sole spawning grounds include nearshore areas outside the survey area, availability of fish within the survey area can vary with the timing of this migration and the timing of the survey. In the case of 2016, a very warm year in the Bering Sea, it appears that a higher portion of the adult biomass was distributed on the shelf (outside of the spawning areas) relative to the average of all previous survey years, indicating earlier spawning migration (Fig 4.7).

Yellowfin sole population numbers-at-age estimated from the annual bottom trawl surveys are shown in Table 4.6 and their occurrence in trawl survey hauls and associated collections of lengths and age structures since 1982 are shown in Table 4.7. Their total tonnage caught in the resource assessment surveys since 1982 are listed in Table 4.8 and also in an appendix table with IPHC survey catches.

#### *Age Determination*

Yellowfin sole ages have been determined at the AFSC by using the break and burn method on collected otoliths since 1979 in surveys and from fisheries. In 2016 the age determination methods for yellowfin sole were validated using the bomb-produced uptake measurement of <sup>14</sup>C method (Kastelle et al. 2016).

#### *Length and Weight-at-Age*

Past assessments of yellowfin sole have used sex-specific, time-invariant growth based on the average length-at-age and weight-at-length relationships from the time-series of survey observations summed over all years since 1982. These weight-at-age estimates were estimated from the following relationships:

Parameters of the von Bertalanffy growth curve have been estimated for yellowfin sole, by sex, from the trawl survey database as follows:

	L <sub>inf</sub>	K	t <sub>0</sub>	n
Males	33.7	0.161	-0.111	656
Females	37.8	0.137	0.112	709

A sex-specific length-weight relationship was also calculated from the survey database using the usual power function, weight (g) = a Length(cm)<sup>b</sup>, where a and b are parameters estimated to provide the best fit to the data (Fig. 4.8).

	a	b	n
males	0.00854	3.081	2,701
females	0.0054	3.227	3,662

These estimates of weight at length were applied to the annual trawl survey estimates of population length at age, by sex, to calculate the weight at each age (Fig. 4.8). Since the resulting estimates of annual weight-at-age were highly variable for fish older than 11 years, ages 11-20 were smoothed using a five year average smoothing method for 1982-2016.

Recent applications of dendrochronology (tree-ring techniques) have been used to develop biochronologies from the otolith growth increments of northern rock sole (*Lepidopsetta polyxystra*), yellowfin sole and Alaska plaice (*Pleuronectes quadrituberculatus*) in the eastern Bering Sea. These techniques ensure that all growth increments are assigned the correct calendar year, allowing for estimation of somatic growth by age and year for chronologies that span approximately 25 years (Matta et al. 2010). The analysis indicated that yellowfin sole somatic growth exhibits annual variability and has a strong positive correlation with May bottom water temperature in the Bering Sea (Fig. 4.9).

The relationship between temperature and growth was further explored by reanalyzing yellowfin sole growth by age and year. Length-weight data collected when obtaining otolith (age) samples in RACE surveys (n=7,000 from 1987, 1994 and 1999-2009) also indicate that weight at age exhibits annual variability and is highly correlated with summer bottom water temperature observations with a lag of 2-3 years for the temperature effect to be seen (shown for age 5 fish in figure 4.10). These observations were then extended back to 1979 using survey population length-at-age estimates (since weight-at-age is a power function of the length-at-age, Clark et al. 1999, Walters and Wilderbuer 2000).

In order to incorporate time-varying (year effect on growth) and temperature-dependent growth functions into the age-structured stock assessment model we used the annual observed population mean weight-at-age (time-varying) from the trawl survey. These empirical data indicate good somatic growth correspondence with annual bottom temperature anomalies from 1982-2017 (Fig. 4.11). Fishery weight at age data available from 2008-2016 were averaged across years for each age to provide updated estimates for the fishery

#### *Maturity-at-age*

Maturity information collected from yellowfin sole females during the 1992 and 1993 eastern Bering Sea trawl surveys have been used in this assessment for the past 20 years (Table 4.10). Nichol (1995) estimated the age of 50% maturity at 10.5 years based on the histological examination of 639 ovaries. Maturity has recently been re-evaluated from a histological analysis of ovaries collected in 2012 (Table 4.10). Results were very similar to the earlier study with only a 2% difference in estimates of yellowfin sole female spawning biomass (TenBrink and Wilderbuer 2015). In addition, the SSC requested that the assessment use a maturity schedule that uses estimates derived from both the 1992 and the 2012 collections (Table 4.10). For yellowfin sole sexual maturity occurs well after the age of entry into the fishery. Yellowfin sole females are 82% selected to the fishery by age 10 whereas they have been found to be only 40% mature at this age.

## Analytic Approach

### Model Structure

The abundance, mortality, recruitment and selectivity of yellowfin sole were assessed with a stock assessment model using the AD Model Builder language (Fournier et al. 2012; Ianelli and Fournier 1998). The conceptual model is a separable catch-age analysis that uses survey estimates of biomass and age composition as auxiliary information (Fournier and Archibald 1982). The assessment model simulates the dynamics of the population and compares the expected values of the population characteristics to the characteristics observed from surveys and fishery sampling programs. This is accomplished by the simultaneous estimation of the parameters in the model using the maximum likelihood estimation procedure. The fit of the simulated values to the observable characteristics is optimized by maximizing a log(likelihood) function given some distributional assumptions about the observed data.

The model starts at age one and fish older than twenty are allowed to accumulate into a plus group. Since the sex-specific weight-at-age for yellowfin sole diverges after age of maturity (about age 10 for 40% of the stock) with females growing larger than males, the current assessment model is coded to accommodate the sex-specific aspects of the population dynamics of yellowfin sole. The model allows for the input of sex-specific estimates of fishery and survey age composition and weight-at-age and provides sex-specific estimates of population numbers, fishing mortality, selectivity, fishery and survey age composition and allows for the estimation of sex-specific natural mortality and catchability. The model retains the utility to fit combined sex data inputs.

The suite of parameters estimated by the model are classified by three likelihood components:

Data component	Distributional assumption
Trawl fishery catch-at-age	Multinomial
Trawl survey population age composition	Multinomial
Trawl survey biomass estimates and S.E.	Log normal

The total log likelihood is the sum of the likelihoods for each data component (Table 4.11). The likelihood components may be weighted by an emphasis factor, however, equal emphasis was placed on fitting each likelihood component in the yellowfin sole assessment except for the catch. The AD Model Builder software fits the data components using automatic differentiation (Griewank and Corliss 1991) software developed as a set of libraries (AUTODIFF C++ library). Table 4.11 also presents the key equations used to model the yellowfin sole population dynamics in the Bering Sea and Table 4.12 provides a description of the variables used in Table 4.11.

Sharp increases in trawl survey abundance estimates for most species of Bering Sea flatfish between 1981 and 1982 indicate that the 83-112 trawl was more efficient for capturing these species than the 400-mesh eastern trawl used in 1975, and 1979-81. Allowing the model to tune to these early survey estimates would most likely underestimate the true pre-1982 biomass, thus exaggerating the degree to which biomass increased during that period. Although this underestimate would have little effect on the estimate of current yellowfin sole biomass, it would affect the spawner and recruitment estimates for the time-series. Hence, the pre-1982 survey biomass estimates were omitted from the analysis.

The model of yellowfin sole population dynamics was evaluated with respect to the observations of the time-series of survey and fishery age compositions and the survey biomass trend since 1982.

### Parameters Estimated Outside the Assessment Model

Natural mortality ( $M$ ) was initially estimated by a least squares analysis where catch-at-age data were fitted to Japanese pair trawl effort data while varying the catchability coefficient ( $q$ ) and  $M$  simultaneously. The best fit to the data (the point where the residual variance was minimized) occurred at

a M value of 0.12 (Bakkala and Wespestad 1984). This was also the value which provided the best fit to the observable population characteristics when M was profiled over a range of values in the stock assessment model using data up to 1992 (Wilderbuer 1992). Since then, natural mortality has been estimated as a free parameter in some of the stock assessment model runs which have been evaluated the past five years. A natural mortality value of 0.12 is used for both sexes in the base model presented in this assessment.

Yellowfin sole maturity schedules were estimated from in-situ observations from two studies as discussed in a previous section (Table 4.10).

## Parameters Estimated Inside the Assessment Model

The parameters estimated by the model are presented below:

Fishing mortality	Selectivity	Survey catchability	Year class strength	Spawner-recruit	Total
65	264	2	104	2	437

The increase in the number of parameters estimated in this assessment compared to last year (6) can be accounted for by the input of another year of fishery data and the entry of another year class into the observed population and four more sex-specific fishery selectivity parameters.

### *Year class strengths*

The population simulation specifies the numbers-at-age in the beginning year of the simulation, the number of recruits in each subsequent year, and the survival rate for each cohort as it moves through the population over time using the population dynamics equations given in Table 4.11.

### *Selectivity*

Fishery and survey selectivity was modeled separately for males and females using the two parameter formulation of the logistic function (Table 4.11). The model was run with an asymptotic selectivity curve for the older fish in the fishery and survey, but still was allowed to estimate the shape of the logistic curve for young fish. The oldest year classes in the surveys and fisheries were truncated at 20 and allowed to accumulate into the age category 20+ years. A single selectivity curve, for both males and females, was fit for all years of survey data.

Given that there have been annual changes in management, vessel participation and most likely gear selectivity, time-varying fishing selectivity curves were estimated. A logistic equation was used to model fishery selectivity and is a function of time-varying parameters specifying the age and slope at 50% selection,  $\varphi_t$  and  $\eta_t$ , respectively. The fishing selectivity ( $S^f$ ) for age  $a$  and year  $t$  is modeled as,

$$S_{a,t}^f = \left[ 1 + e^{\eta_t(a - \varphi_t)} \right]^{-1}$$

where  $\eta_t$  and  $\varphi_t$  are time-varying and partitioned (for estimation) into parameters representing the mean and a vector of deviations (log-scale) conditioned to sum to zero. The deviations are constrained by a lognormal prior with a variance that was iteratively estimated. The process of iterating was to first set the variance to a high value (diffuse prior) of 0.5<sup>2</sup> and estimate the deviations. The next step was to compare the variability of model estimates. The variance of the model estimates were then rounded up slightly and fixed for subsequent runs. The 2016 values were fixed as the average of the 3 most recent years.

### *Fishing Mortality*

The fishing mortality rates (F) for each age and year are calculated to approximate the catch weight by solving for F while still allowing for observation error in catch measurement. A large emphasis (300) was placed on the catch likelihood component to force the model to closely match the observed catch.

### *Survey Catchability*

A past assessment (Wilderbuer and Nichol 2001) first examined the relationship between estimates of survey biomass and bottom water temperature. To better understand how water temperature may affect the catchability of yellowfin sole to the survey trawl, catchability was estimated for each year in the stock assessment model as:

$$q = e^{-\alpha + \beta T}$$

where  $q$  is catchability,  $T$  is the average annual bottom water temperature anomaly at survey stations less than 100 m, and  $\alpha$  and  $\beta$  are parameters estimated by the model. The catchability equation has two parts. The  $e^{-\alpha}$  term is a constant or time-independent estimate of  $q$ . The second term,  $e^{\beta T}$  is a time-varying (annual)  $q$  which responds to metabolic aspects of herding or distribution (availability) which can vary annually with bottom water temperature. The result of incorporating bottom temperature to estimate annual  $q$  has resulted in an improved fit to the survey (shown in Figure 4.12 for the base model).

### *Spawner-Recruit Estimation*

Annual recruitment estimates from 1978-2012 were constrained to fit a Ricker (1958) form of the stock recruitment relationship as follows:

$$R = \alpha S e^{-\beta S}$$

where  $R$  is age 1 recruitment,  $S$  is female spawning biomass ( $t$ ) the previous year, and  $\alpha$  and  $\beta$  are parameters estimated by the model. The spawner-recruit fitting is estimated in a later phase after initial estimates of survival, numbers-at-age and selectivity are obtained.

## **Results**

### **Model Evaluation**

The model evaluation for this stock assessment involved a two-step process. The first step was to evaluate the productivity of the yellowfin sole stock by an examination of which sets of years to include for spawner-recruit fitting (increased from 1978-2010 to 1978-2012 in this assessment). The second step evaluated various hypothesized states of nature by fitting natural mortality and catchability estimates in various combinations.

The SSC determined in December 2006 that yellowfin sole would be managed under the Tier 1 harvest guidelines, and therefore future harvest recommendations would be based on MSY and  $F_{MSY}$  values calculated from a spawner-recruit relationship. MSY is an equilibrium concept and its value is dependent on both the spawner-recruit estimates which are assumed to represent the equilibrium stock size-recruitment relationship and the model used to fit the estimates. In the yellowfin sole stock assessment model, a Ricker form of the stock-recruit relationship was fit to various combinations of these data and estimates of  $F_{MSY}$  and  $B_{MSY}$  were calculated, assuming that the fit to the stock-recruitment data represents the long-term productivity of the stock.

For this assessment, 2 different stock-recruitment time-series were investigated: the full time-series 1955-2012 (Model 14\_2) and the post-regime shift era, 1978-2012 (Model 14\_1) (Fig. 4.13) (see Joint Plan Team recommendations for September 2012). Very different estimates of the long-term sustainability of the stock ( $F_{MSY}$  and  $B_{MSY}$ ) are obtained, depending on which years of stock-recruitment data are included in the fitting procedure (Table 4.13). When the entire time-series from 1955-2012 was fit, the large recruitments that occurred at low spawning stock sizes in the 1960s and early 1970s determined that the yellowfin sole stock was most productive at a smaller stock size with the result that  $F_{MSY}$  (0.19) is higher than  $F_{35\%}$  ( $F_{35\%} = 0.14$ ) and  $B_{MSY}$  is 333,700 (Model 14\_2). If we limit the analysis to consider only recruitments which occurred after the well-documented regime shift in 1977 (Model 14\_1), a lower value of  $F_{MSY}$  is obtained (0.114) and  $B_{MSY}$  is 456,200 t. Table 4.13 indicates that the ABC values from the

Model 14\_2 harvest scenario for 2018 would be 193,800 t higher than Model 14\_1. Posterior distributions of  $F_{MSY}$  for these models indicate that this parameter is estimated with less uncertainty for Model 14\_1 resulting in the reduced buffer between ABC and OFL relative to Model 14\_2 (11% for Model 14\_1 versus 1% for Model 14\_2, Table 4.13 and Fig 4.14).

It is important for the Tier 1 calculations to identify which subset of the stock recruitment data is used. Using the full time series to fit the spawner recruit curve estimates that the stock is most productive at a small stock size. Thus MSY and  $F_{MSY}$  are relatively high values and  $B_{MSY}$  is a lower value. If the stock was productive in the past at a small stock size because of non-density dependent factors (environment), then reducing the stock size to low levels could be detrimental to the long-term sustainability of the stock if the environment, and thus productivity, have changed from the earlier period. Since observations of yellowfin sole recruitment at low stock sizes are not available from multiple time periods, it is uncertain if future recruitment events at low stock conditions would be as productive as during the late 1960s-early 1970s.

Given the uncertainty of the productivity of yellowfin sole at low spawning stock sizes, and because the AFSC policy for reference point time-series selection is to use the post 1977 regime shift values unless there is a compelling reason to do otherwise, the productivity of yellowfin sole in this assessment is estimated by fitting the 1977-2012 spawner-recruit data in the model (Model 14\_1).

The second step in the model evaluation for this assessment entails the use of a single structural model to consider the uncertainty in the key parameters M and catchability. This is the Model which has been the model of choice in the past 7 assessments (Model 14\_1) and operates by fixing M at 0.12 for both sexes and then estimates q using the relationship between survey catchability and the annual average water temperature at the sea floor (from survey stations at less than 100 m). The other models used in the evaluation represented various combinations of estimating M or q as free parameters with different amounts of uncertainty in the parameter estimates (Wilderbuer et al. 2010). The results are detailed in those assessments and are not repeated here except for the following observations.

Modeling survey catchability as a nonlinear function of bottom water temperature returns q estimates > 1.0 for years when the bottom temperature is anomalously warm (greater than the mean temperature) and less than 1.0 when below the temperature mean. These values are consistent with our hypothesis that more fish are available to the survey in warm years relative to cooler years due to the timing of the annual spawning migration to nearshore areas that occurs sooner in warm years.


Experiments examining the bridle efficiency of the Bering Sea survey trawl indicate that yellowfin sole are herded into the trawl path from an area between the wing tips of the net and the point where the bridles contact the seafloor (Somerton and Munro 2001). The herding experiments suggest that the survey trawl catchability is greater than 1.0. The likelihood profile of q from the model indicated a small variance with a narrow range of likely values with a low probability of q being equal to the value of 1.0 in a past assessment (Wilderbuer and Nichol 2003).

A model that allows M to be estimated as a free parameter for males with females fixed at 0.12 provided a better fit to the sex ratio estimated from the annual trawl survey age compositions than did the base model (both sexes fixed at M = 0.12). However, since the population sex ratio annually observed at the time of the survey is a function of the timing of the annual spawning in adjacent inshore areas, it is questionable that providing the best fit to these observations is really fitting the population sex ratio better. Thus, the model configuration which utilizes the relationship between annual seafloor temperature and survey catchability with M fixed at 0.12 for both sexes (Model 14\_1) is the preferred model used to base the assessment of the condition of the Bering Sea yellowfin sole resource for the 2018 fishing season.

## Time Series Results

Before presenting the preferred model results, a brief consideration of the inputs and changes to the assessment methodology relative to last year (Model 14\_1) is given. Primary updates for Model 14\_1 were the 2017 catch, the fishery and survey age compositions from 2016 and the 2017 survey biomass (3% lower than 2016) and standard error estimates. The fishery and survey weights-at-age were also changed in a small amount to include the latest year of data and the spawner-recruit data was increased by 2 years to include the 2011 and 2012 year-class recruitment estimates. In their totality, these changes produced a Model 14\_1  $F_{ABC}$  estimate that was 5% lower than 2016 and a  $F_{FOFL}$  that was 4% lower.

This increase was mostly due to refitting the spawner-recruit curve with the averaged fishery weight-at-age data which had the effect of moving the curve and  $B_{MSY}$  to the right (higher  $B_{MSY}$ , lower  $F_{MSY}$  for Tier 1 stocks).



The 2017 overall estimate (1982 – 2017) of trawl survey catchability decreased from 0.95 to 0.9. This resulted in higher model estimates of population numbers at age and biomass for the time-series back to the mid-1960s relative to last year's assessment and increased the estimated level of female spawning biomass. The model results indicate the stock has been in a slowly declining condition since the mid-1980s. The estimates of total biomass and ABC are higher than those used to manage the stock in 2017. Seven of the past 11 years have had negative bottom temperature anomalies in the Bering Sea but the last four have been above the mean. The temperature-dependent  $q$  adjustment for 2017 was 0.92.

### *Fishing Mortality and Selectivity*

The assessment model estimates of the annual fishing mortality in terms of age-specific annual  $F$  and on fully selected ages are given in Tables 4.14 and 4.15, respectively. The full-selection  $F$  has averaged 0.08 over the period of 1978-2017 with a maximum of 0.12 in 1978 and a minimum of 0.04 in 2001.

Selectivities estimated by the model (Table 4.16, Fig. 4.15) indicate that both sexes of yellowfin sole are 50% selected by the fishery at about age 9 and nearly fully selected by age 13, with annual variability.

### *Abundance Trend*

The model estimates  $q$  at an average value of 0.9 for the period 1982-2017 which results in the model estimate of the 2016 age 2+ total biomass at 2,878,100 t (Table 4.17). Model results indicate that yellowfin sole total biomass (age 2+) was at low levels during most of the 1960s and early 1970s (700,000-1,000,000 t) after a period of high exploitation (Table 4.17, Fig. 4.16, center left panel).

Sustained above average recruitment from 1967-76 combined with light exploitation resulted in a biomass increase to a peak of 3.5 million t by 1985. The population biomass has since been in a slow decline as the strong 1981 and 1983 year-classes have passed through the population, with only the 1991, 1995 and 2003 year-classes at levels observed during the 1970s. The present biomass is estimated at 82% of the peak 1985 level.

The female spawning biomass has also declined since the peak in 1994, with a 2017 estimate of 884,800 t (22% decline). The spawning biomass has been in a gradual decline for the past 22 years and is 30% above the  $B_{40\%}$  level and 1.9 times the  $B_{MSY}$  level (Fig. 4.16). The model estimate of yellowfin sole population numbers at age for all years is shown in Table 4.18 and the resulting fit to the observed fishery and survey age compositions input into the model are shown in the Figure 4.17. The fit to the trawl survey biomass estimates are shown in Figure 4.16. Allowing  $q$  to be correlated with annual bottom temperature provides a better fit to the bottom trawl survey estimates than using a  $q$  fixed at the average value (Fig. 4.18). Table 4.19 lists the numbers of female spawners estimated by the model for all ages and years. The estimated average age of yellowfin sole in the population is 6.7 years for males and females.

Both the trawl survey and the stock assessment model indicate that the yellowfin sole resource increased during the 1970s and early 1980s to a peak level during the mid-1980s. The yellowfin sole population biomass slowly decreased over the 22 years since the mid-1990s as the majority of year-classes during those years were below average strength. Average to above average recruitment from 2006 to 2009 is expected to maintain the abundance of yellowfin sole at a level above  $B_{MSY}$  in the near future. The stock assessment projection model indicates a mildly decreasing trend in female spawning biomass through 2024 if the fishing mortality rate continues at the same level as the average of the past 5 years (Fig. 4.22).

#### *Recruitment Trends*

The primary reason for the sustained increase in abundance of yellowfin sole during the 1970s and early 1980s was the recruitment of a series of stronger than average year classes spawned in 1967-76 (Figure 4.19 and Table 4.20). The 1981 year class was the strongest observed (and estimated) during the 47 year period analyzed and the 1983 year class was also very strong. Survey age composition estimates and the assessment model also estimate that the 1987 and 1988 year classes were average and the 1991 and 1995 year classes were above average. With the exception of these 4 year classes, recruitment from 15 of the following 19 years estimated from 1984-2005 (since the strong 1983 year-class) were below the 48 year average, which caused the population to gradually decline. The 2003 year-class has now been observed multiple times in the age compositions and is clearly a strong year class, similar to some of the strong recruitment mentioned above and are contributing to the reservoir of spawning fish in the current population. In addition, recruitment from 2006-2009 appear also to be average to above average.

#### *Historical Exploitation Rates*

Based on results from the stock assessment model, annual average exploitation rates of yellowfin sole since 1977 ranged from 3 to 7% of the total biomass, and have averaged 4% (Table 4.15). Posterior distributions of selected parameters from the preferred stock assessment model used in the assessment are shown in Figure 4.20. The values and standard deviations of some selected model parameters are listed in Table 4.21.

#### *Retrospective analysis*

A within-model retrospective analysis is also included for the recommended assessment model (Model 14\_1) where retrospective female spawning biomass is calculated by working backwards in time dropping data one year at a time and then comparing the “peeled” estimate to the reference stock assessment model used in the assessment (Fig. 4.21). The resulting pattern from the current assessment model was less than desirable.

Peculiar to the yellowfin sole assessment, in comparison to the northern rock sole and Alaska plaice assessments (that have nice patterns), is the large amount of variability in the annual survey biomass

assessments for this stock due to the temperature-influenced availability to the survey. This large variability in the annual estimates can contribute to undesirable patterns since the earlier years are not fitting the same highly variable information as the current year. Exploratory model runs were made to examine the influence of fitting the survey biomass on the retrospective patterns. This was accomplished by making model runs that increased the survey biomass standard error by 10%, 20% and 30%, and also runs that attempted to decrease the influence of fitting the survey age composition by decreasing the effective n, and also a run with a fixed q (no bottom temp modeling). The models are listed below and are evaluated using Mohn's test statistic.

Model Description	Mohn's test statistic
Current stock assessment model (Model 2)	-0.193
Model 14_1 with survey standard error increased by 10%	-0.239
Model 14_1 with survey standard error increased by 20%	-0.211
Model 14_1 with survey standard error increased by 30%	-0.219
Down-weighted survey age comps, base st. dev values	-0.207
Up-weighted survey age comps (500) and down-weighted survey SE (increased st. dev value by 30%)	-0.238
survey q fixed at 1.05 for all runs	-0.186


A small gain was made by fixing q at 1.05 ( $M = 0.12$ ) whereas all the other runs had similar results, increasing the test statistic over the base value (lower is better).

The Plan Team suggested examining the effect that q and M have on the retrospective patterns. Responding to this request, model runs were made varying M from 0.9 to 0.14 and q from 0.8 to 1.2 (fixed M and q, no temperature modeling). Decreasing values of q and M gave better retrospective patterns and lower Mohn's test statistic.

		Catchability					
		0.8	0.9	1	1.1	1.2	
	0.09	0.019	0.003	0.028	0.0511	0.076	
	0.1	0.013	0.036	0.059	0.084	0.107	
Natural mortality	0.11	0.043	0.066	0.087	0.11	0.131	
	0.12	0.07	0.105	0.113	0.135	0.155	
	0.13	0.051	0.114	0.135	0.156	0.177	
	0.14	0.114	0.135	0.153	0.177	0.2	

Hotter colors correspond to lower test statistic values.

Figure below is an example of retrospective pattern with low M and q.



### **Harvest Recommendations**

The SSC has determined that yellowfin sole qualify as a Tier 1 stock and therefore the 2018 ABC is calculated using Tier 1 methodology. The Tier 1 harvest level is calculated as the product of the harmonic mean of  $F_{MSY}$  and the geometric mean of the 2018 biomass estimate, as follows:

$$B_{gm} = e^{\frac{\ln \hat{B} - \frac{cv^2}{2}}{2}}$$

, where  $B_{gm}$  is the geometric mean of the 2018 biomass estimate,  $\hat{B}$  is the point estimate of the 2018 biomass from the stock assessment model and  $cv^2$  is the coefficient of variation of the point estimate (a proxy for sigma);

and

$$\bar{F}_{har} = e^{\frac{\ln \hat{F}_{msy} - \frac{\ln sd^2}{2}}{2}}$$

, where  $\bar{F}_{har}$  is the harmonic mean,  $\hat{F}_{msy}$  is the peak mode of the  $F_{MSY}$  distribution and  $sd^2$  is the square of the standard deviation of the  $F_{MSY}$  distribution.

In 2006 the SSC selected the 1978-2001 data set for the Tier 1 harvest recommendation. Using this approach again for the 2018 harvest (now the 1978-2012 time-series) recommendation (Model 14\_1 in Table 4.13), the  $F_{ABC} = F_{harmonic\ mean} = 0.109$ . The estimate of age 6+ total biomass for 2018 is 2,553,100 t. The calculations outlined above give a Tier 1 ABC harvest recommendation of **277,500 t** and an OFL of 306,700 t for 2018. This results in an 11% (29,250 t) buffer between ABC and OFL. The ABC value is 6% higher than last year, primarily due to changes to the spawner-recruit curve from the fishery weight-at-age modeling and a lower catchability estimate.

The stock assessment analysis must also consider harvest limits, usually described as overfishing fishing mortality levels with corresponding yield amounts. Amendment 56 to the BSAI FMP sets the Tier 1 harvest limit at the  $F_{MSY}$  fishing mortality value. The overfishing fishing mortality values, ABC fishing mortality values and their corresponding yields are given as follows:

<u>Harvest level</u>	<u>F value</u>	<u>2018 Yield</u>
<b>Tier 1 <math>F_{OFL} = F_{MSY}</math></b>	<b>0.12</b>	<b>306,700 t</b>
<b>Tier 1 <math>F_{ABC} = F_{harmonic\ mean}</math></b>	<b>0.109</b>	<b>277,500 t</b>

### *Status Determination*

A standard set of projections is required for each stock managed under Tiers 1, 2, or 3 of Amendment 56. This set of projections encompasses seven harvest scenarios designed to satisfy the requirements of Amendment 56, the National Environmental Policy Act, and the Magnuson-Stevens Fishery Conservation and Management Act (MSFCMA).

For each scenario, the projections begin with the vector of 2017 numbers at age estimated in the assessment. This vector is then projected forward to the beginning of 2018 using the schedules of natural mortality and selectivity described in the assessment and the best available estimate of total (year-end) catch for 2017. In each subsequent year, the fishing mortality rate is prescribed on the basis of the spawning biomass in that year and the respective harvest scenario. In each year, recruitment is drawn from an inverse Gaussian distribution whose parameters consist of maximum likelihood estimates determined from recruitments estimated in the assessment. Spawning biomass is computed in each year based on the time of peak spawning and the maturity and weight schedules described in the assessment. Total catch is assumed to equal the catch associated with the respective harvest scenario in all years. This projection scheme is run 1000 times to obtain distributions of possible future stock sizes, fishing mortality rates, and catches.

Five of the seven standard scenarios will be used in an Environmental Assessment prepared in conjunction with the final SAFE. These five scenarios, which are designed to provide a range of harvest alternatives that are likely to bracket the final TAC for 2018, are as follows (“ $\max F_{ABC}$ ” refers to the maximum permissible value of  $F_{ABC}$  under Amendment 56):

*Scenario 1:* In all future years,  $F$  is set equal to  $\max F_{ABC}$ . (Rationale: Historically, TAC has been constrained by ABC, so this scenario provides a likely upper limit on future TACs.)

*Scenario 2:* In all future years,  $F$  is set equal to a constant fraction of  $\max F_{ABC}$ , where this fraction is equal to the ratio of the  $F_{ABC}$  value for 2018 recommended in the assessment to the  $\max F_{ABC}$  for 2018. (Rationale: When  $F_{ABC}$  is set at a value below  $\max F_{ABC}$ , it is often set at the value recommended in the stock assessment.)

*Scenario 3:* In all future years,  $F$  is set equal to the 2013-2017 average  $F$ . (Rationale: For some stocks, TAC can be well below ABC, and recent average  $F$  may provide a better indicator of  $F_{TAC}$  than  $F_{ABC}$ .)

*Scenario 4:* In all future years, the upper bound on  $F_{ABC}$  is set at  $F_{60\%}$ . (Rationale: This scenario provides a likely lower bound on  $F_{ABC}$  that still allows future harvest rates to be adjusted downward when stocks fall below reference levels.)

*Scenario 5:* In all future years,  $F$  is set equal to zero. (Rationale: In extreme cases, TAC may be set at a level close to zero.)

Two other scenarios are needed to satisfy the MSFCMA’s requirement to determine whether a stock is currently in an overfished condition or is approaching an overfished condition. These two scenarios are as follow (for Tier 3 stocks, the MSY level is defined as  $B_{35\%}$ ):

*Scenario 6:* In all future years,  $F$  is set equal to  $F_{OFL}$ . (Rationale: This scenario determines whether a stock is overfished. If the stock is expected to be above its MSY level in 2016 and above its MSY level in 2029 under this scenario, then the stock is not overfished.)

*Scenario 7:* In 2018 and 2019,  $F$  is set equal to  $\max F_{ABC}$ , and in all subsequent years,  $F$  is set equal to  $F_{OFL}$ . (Rationale: This scenario determines whether a stock is approaching an overfished condition. If the stock is expected to be above its MSY level in 2030 under this scenario, then the stock is not approaching an overfished condition.)

Simulation results shown in Table 4.22 indicate that yellowfin sole are not currently overfished and are not approaching an overfished condition. The projection of yellowfin sole female spawning biomass

through 2030 is shown in Figure 4.22 and a phase plane figure of the estimated time-series of yellowfin sole female spawning biomass relative to the harvest control rule is shown in Figure 4.23.

#### *Scenario Projections and Two-Year Ahead Overfishing Level*

In addition to the seven standard harvest scenarios, Amendments 48/48 to the BSAI and GOA Groundfish Fishery Management Plans require projections of the likely OFL two years into the future. The 2016 numbers at age from the stock assessment model are projected to 2017 given the 2016 catch and then a 2017 catch of 150,000 t is applied to the projected 2017 population biomass to obtain the 2018 OFL.


Tier 1 Projection			Geometric mean 6+ total biomass		
Year	Catch	SSB		ABC	OFL
2018	150,000	895,000	2,553,100	277,500	306,700
2019	150,000	890,000	2,460,700	267,400	295,600

## Ecosystem Considerations

### Ecosystem Effects on the stock

#### *1) Prey availability/abundance trends*

Yellowfin sole diet by life stage varies as follows: Larvae consume plankton and algae, early juveniles consume zooplankton, late juvenile stage and adults prey includes bivalves, polychaetes, amphipods, mollusks, euphausiids, shrimps, brittle stars, sculpins and miscellaneous crustaceans. Information is not available to assess the abundance trends of the benthic infauna of the Bering Sea shelf. The original description of infaunal distribution and abundance by Haflinger (1981) resulted from sampling conducted in 1975 and 1976 and has not been re-sampled since. The large populations of flatfish which have occupied the middle shelf of the Bering Sea over the past twenty-five years for summertime feeding do not appear food-limited. These populations have fluctuated due to the variability in recruitment success which suggests that the primary infaunal food source has been at an adequate level to sustain the yellowfin sole resource.



#### *2) Predator population trends*

As juveniles, it is well-documented from studies in other parts of the world that flatfish are prey for shrimp species in near shore areas. This has not been reported for Bering Sea yellowfin sole due to a lack of juvenile sampling and collections in near shore areas, but is thought to occur. As late juveniles they have been found in stomachs of Pacific cod and Pacific halibut; mostly small yellowfin sole ranging from 7 to 25 cm standard length..

Past, present and projected future population trends of these predator species can be found in their respective SAFE chapters in this volume and also from Annual reports compiled by the International

Pacific Halibut Commission. Encounters between yellowfin sole and their predators may be limited since their distributions do not completely overlap in space and time.

*3) Changes in habitat quality*

Changes in the physical environment which may affect yellowfin sole distribution patterns, recruitment success and migration timing patterns are catalogued in the Ecosystem Considerations Report of this SAFE report. Habitat quality may be enhanced during years of favorable cross-shelf advection (juvenile survival) and warmer bottom water temperatures with reduced ice cover (higher metabolism with more active feeding).

### **Fishery Effects on the ecosystem**

- 1) The yellowfin sole target fishery contribution to the total bycatch of other target species is shown for 1992-2016 in Table 4.23. The catch of non-target species from 2003-2016 is shown in Table 4.24. The yellowfin sole target fishery contribution to the total bycatch of prohibited species is shown for 2014 and 2015 in Table 13 of the Economic SAFE (Appendix C) and is summarized for 2015 as follows:

Prohibited species	Yellowfin sole fishery % of total bycatch
Halibut mortality	30
Herring	2
Red King crab	5
<u>C. bairdi</u>	25.5
Other Tanner crab	78.2
Salmon	<1

- 2) Relative to the predator needs in space and time, the yellowfin sole target fishery has a low selectivity for fish 7-25 cm and therefore has minimal overlap with removals from predation.
- 3) The target fishery is not perceived to have an effect on the amount of large size target fish in the population due to its history of light to moderate exploitation (6%) over the past 30 years. Population age composition data indicate a large 20+ age group.
- 4) Yellowfin sole fishery discards are presented in the Catch History section.
- 5) It is unknown what effect the fishery has had on yellowfin sole maturity-at-age and fecundity, but based on two maturity studies conducted 20 years apart, it is expected to be minimal.
- 6) Analysis of the benthic disturbance from the yellowfin sole fishery is available in the Preliminary draft of the Essential Fish Habitat Environmental Impact Statement.

<b>Ecosystem effects on yellowfin sole</b>			
Indicator	Observation	Interpretation	Evaluation
<i>Prey availability or abundance trends</i>			
Benthic infauna	Stomach contents	Stable, data limited	Unknown
<i>Predator population trends</i>			
Fish (Pacific cod, halibut, skates)	Stable	Possible increases to yellowfin sole mortality	
<i>Changes in habitat quality</i>			
Temperature regime	Cold years yellowfin sole catchability and herding may decrease, timing of migration may be prolonged	Likely to affect surveyed stock	No concern (dealt with in model)
Winter-spring environmental conditions	Affects pre-recruit survival	Probably a number of factors	Causes natural variability
<b>Yellowfin sole effects on ecosystem</b>			
Indicator	Observation	Interpretation	Evaluation
<i>Fishery contribution to bycatch</i>			
Prohibited species	Stable, heavily monitored	Minor contribution to mortality	No concern
Forage (including herring, Atka mackerel, cod, and pollock)	Stable, heavily monitored	Bycatch levels small relative to forage biomass	No concern
HAPC biota	Low bycatch levels of (spp)	Bycatch levels small relative to HAPC biota	No concern
Marine mammals and birds	Very minor direct-take	Safe	No concern
Sensitive non-target species	Likely minor impact	Data limited, likely to be safe	No concern
<i>Fishery concentration in space and time</i>	Low exploitation rate	No concern Little detrimental effect	
<i>Fishery effects on amount of large size target fish</i>			
	Low exploitation rate	Natural fluctuation	No concern
<i>Fishery contribution to discards and offal production</i>			
	Stable trend	Improving, but data limited	Possible concern
<i>Fishery effects on age-at-maturity and fecundity</i>			
	Unknown	NA	Possible concern

## Data Gaps and Research Priorities

Isolation by distance genetic study to define stock structure in the planning stage. NPRB proposal to collect maturity in the northern Bering Sea for comparison with recent SE Bering Sea shelf samples.

## References

- Alaska Fisheries Science Center. 2016. Wholesale market profiles for Alaska groundfish and crab fisheries. 134 p. Alaska Fish. Sci. Cent., NOAA, Natl. Mar. Fish. Serv. 7600 Sand Point Way NE. Seattle, WA.
- Bakkala, R. G. and V. Wespestad. 1984. Yellowfin sole. In R. G. Bakkala and L. resources of the eastern Bering Sea and Aleutian Islands region in 1983, p. 37-60. U.S. Dep. Commer., NOAA Tech. Memo. NMFS F/NWC-53.
- Bakkala, R. G., V. Wespestad, and L. Low. 1982. The yellowfin sole (*Limanda aspera*) resource of the eastern Bering Sea--its current and future potential for commercial fisheries. U.S. Dep. Commer., NOAA Tech. Memo. NMFS F/NWC-33, 43p.
- Bakkala, R. G., and T. K. Wilderbuer. 1990. Yellowfin sole. In Stock Assessment and Fishery Evaluation Document for Groundfish Resources in the Bering Sea/Aleutian Islands Region as Projected for 1990, p. 60-78. North Pacific Fishery Management Council, P. O. Box 103136, Anchorage, Ak 99510.
- Clark, W. G., Hare, S. R., Parmas, A. M., Sullivan, P. J., Trumble, R. J. 1999. Decadal changes in growth and recruitment of Pacific halibut (*Hipplglossus stenolepis*). Can. J. fish. Aquat. Sci. 56, 242-252.
- Fournier, D. A., H.G. Skaug, J. Ancheta, J. Ianelli, A. Magnusson, M. N. Maunder, A. Nielsen, and J. Sibert. 2012. AD Model Builder: using automatic differentiation for statistical inference of highly parameterized complex nonlinear models. Optim. Methods Softw. 27:233-239.
- Fournier, D. A. and C.P. Archibald. 1982. A general theory for analyzing catch-at-age data. Can. J. Fish Aquat. Sci. 39:1195-1207.
- Griewank, A. and G. F. Corliss (eds) 1991. Automatic differentiation of algorithms: theory, implementation and application. Proceedings of the SIAM Workshop on the Automatic Differentiation of Algorithms, held Jan. 6-8, Breckenridge, CO. Soc. Indust. And Applied Mathematics, Philadelphia.
- Haflinger, K. 1981. A survey of benthic infaunal communities of the Southeastern Bering Sea shelf. In Hood and Calder (editors) The Eastern Bering Sea Shelf: Oceanography and Resources, Vol. 2. P. 1091-1104. Office Mar. Pol. Assess., NOAA. Univ. Wash. Press, Seattle, Wa 98105.
- Ianelli, J. N. and D. A. Fournier. 1998. Alternative age-structured analyses of the NRC simulated stock assessment data. In Restrepo, V. R. [ed.] Analyses of simulated data sets in support of the NRC study on stock assessment methods. NOAA Tech. Memo. NMFS-F/SPO-30. 96 p.
- Kastelle, C., T. Helser, S. Wischniowski, T. Loher, B. Geotz and L. Kautzi. 2016. Incorporation of bomb-produced <sup>14</sup>C into fish otoliths: A novel approach for evaluating age validation and bias with an application to yellowfin sole and northern rockfish. Ecological modeling 320 (2016) 79-91.
- Low, L. and R.E. Narita. 1990. Condition of groundfish resources in the Bering Sea-Aleutian Islands region as assessed in 1988. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-F/NWC-178, 224 p.
- Matta, M. E., B. A. Black and T. K. Wilderbuer. 2010. Climate-driven synchrony in otolith growth-increment chronologies for three Bering Sea flatfish species. MEPS, Vol. 413:137-145, 2010.
- Nichol, D. R . 1995. Spawning and maturation of female yellowfin sole in the eastern Bering Sea. In Proceedings of the international flatfish symposium, October 1994, Anchorage, Alaska, p. 35-50. Univ. Alaska, Alaska Sea Grant Rep. 95-04.

- Nichol, D.R. 1998. Annual and between sex variability of yellowfin sole, *Pleuronectes asper*, spring-summer distributions in the eastern Bering Sea. Fish. Bull., U.S. 96: 547-561.
- Ricker, W. E. 1958. Handbook of computations for biological statistics of fish populations. Bull. Fish. Res. Bd. Can., (119) 300 p.
- Rose, C. S., J. R. Gauvin and C. F. Hammond. 2010. Effective herding of flatfish by cables with minimal seafloor contact. Fishery Bulletin 108(2):136-144.
- Somerton, D. A. and P. Munro. 2001. Bridle efficiency of a survey trawl for flatfish. Fish. Bull. 99:641-652 (2001).
- TenBrink, T. T. and T. K. Wilderbuer. 2015. Updated maturity estimates for flatfishes (Pleuronectidae) in the eastern Bering Sea, with notes on histology and implications to fisheries management. Coastal and Marine Fisheries: Dynamics, Management and Ecosystem Science. O:1-9. 2015. DOI: 10.1080/19425120.2015.1091411.
- Wakabayashi, K. 1989. Studies on the fishery biology of yellowfin sole in the eastern Bering Sea. [In Jpn., Engl. Summ.] Bull. Far Seas Fish. Res. Lab. 26:21-152.
- Wakabayashi, K., R. Bakkala, and L. Low. 1977. Status of the yellowfin sole resource in the eastern Bering Sea through 1976. Unpubl. manuscr., 45p. Northwest and Alaska Fish. Cent., Natl. Mar. Fish. Serv., NOAA, 7600 Sand Point Way N.E., Bin C 15700, Seattle, Wa 98115.
- Walters, G. E. and T. K. Wilderbuer. 2000. Decreasing length at age in a rapidly expanding population of northern rock sole in the eastern Bering Sea and its effect on management advice. Journal of Sea Research 44(2000)17-26.
- Wilderbuer, T. K. 1992. Yellowfin sole. In Stock Assessment and Fishery Evaluation Document for Groundfish Resources in the Bering Sea/Aleutian Islands Region as Projected for 1993, chapter 3. North Pacific Fishery Management Council, P. O. Box 103136, Anchorage, Ak 99510.
- Wilderbuer, T. K. 1993. Yellowfin sole. In Stock Assessment and Fishery Evaluation Document for Groundfish Resources in the Bering Sea/Aleutian Islands Region as Projected for 1994, chapter 3. North Pacific Fishery Management Council, P. O. Box 103136, Anchorage, Ak 99510.
- Wilderbuer, T. K. and D. Nichol. 2001. Yellowfin sole. In Stock Assessment and Fishery Evaluation Document for Groundfish Resources in the Bering Sea/Aleutian Islands Region as Projected for 2004, chapter 4. North Pacific Fishery Management Council, P. O. Box 103136, Anchorage, Ak 99510.
- Wilderbuer, T. K. and D. Nichol. 2003. Yellowfin sole. In Stock Assessment and Fishery Evaluation Document for Groundfish Resources in the Bering Sea/Aleutian Islands Region as Projected for 2004, chapter 4. North Pacific Fishery Management Council, P. O. Box 103136, Anchorage, Ak 99510.
- Wilderbuer, T. K. D. G. Nichol, and J. Ianelli. 2010. Yellowfin sole. In Stock Assessment and Fishery Evaluation Document for Groundfish Resources in the Bering Sea/Aleutian Islands Region as Projected for 2011, chapter 4. North Pacific Fishery Management Council, P. O. Box 103136, Anchorage, Ak 99510.
- Wilderbuer, T.K., G.E. Walters, and R.G. Bakkala 1992. Yellowfin sole, *Pleuronectes asper*, of the eastern Bering Sea: biological characteristics, history of exploitation, and management. Mar Fish. Rev. 54(4):1-18.

## Tables

Table 4.1--Catch (t) of yellowfin sole 1964-2017. Catch for 2017 is an estimate through the end of 2017.

Year	Foreign	Domestic		Total
		JVP	DAP	
1964	111,777			111,777
1965	53,810			53,810
1966	102,353			102,353
1967	162,228			162,228
1968	84,189			84,189
1969	167,134			167,134
1970	133,079			133,079
1971	160,399			160,399
1972	47,856			47,856
1973	78,240			78,240
1974	42,235			42,235
1975	64,690			64,690
1976	56,221			56,221
1977	58,373			58,373
1978	138,433			138,433
1979	99,019			99,019
1980	77,768	9,623		87,391
1981	81,255	16,046		97,301
1982	78,331	17,381		95,712
1983	85,874	22,511		108,385
1984	126,762	32,764		159,526
1985	100,706	126,401		227,107
1986	57,197	151,400		208,597
1987	1,811	179,613	4	181,428
1988		213,323	9,833	223,156
1989		151,501	1,664	153,165
1990		69,677	14,293	83,970
1991			115,842	115,842
1992			149,569	149,569
1993			106,101	106,101
1994			144,544	144,544
1995			124,740	124,740
1996			129,659	129,659
1997			181,389	181,389
1998			101,201	101,201
1999			67,320	67,320
2000			83,850	83,850
2001			63,395	63,395
2002			73,000	73,000
2003			74,418	74,418
2004			69,046	69,046
2005			94,383	94,383
2006			99,068	99,068
2007			121,029	121,029
2008			148,894	148,894
2009			107,528	107,528
2010			118,624	118,624
2011			151,164	151,164
2012			147,183	147,183
2013			164,944	164,944
2014			156,778	156,778
2015			126,933	126,933
2016			135,353	135,353
2017			143,100	143,100

Table 4.2. Estimates of retained and discarded (t) yellowfin sole caught in Bering Sea fisheries.

<b>Year</b>	<b>Retained</b>	<b>Discarded</b>
1987	3	1
1988	7,559	2,274
1989	1,279	385
1990	10,093	4,200
1991	89,054	26,788
1992	103,989	45,580
1993	76,798	26,838
1994	107,629	36,948
1995	96,718	28,022
1996	101,324	28,334
1997	149,570	31,818
1998	80,365	20,836
1999	55,202	12,118
2000	69,788	14,062
2001	54,759	8,635
2002	62,050	10,950
2003	63,732	10,686
2004	57,378	11,668
2005	85,321	9,062
2006	90,570	8,498
2007	109,084	11,945
2008	141,253	7,659
2009	92,488	5,733
2010	113,244	5,380
2011	146,419	4,745
2012	143,737	3,446
2013	158,781	6,163
2014	152,164	4,614
2015	123,065	3,871
2016	131,205	4,148

Table 4.3. Discarded and retained catch of non-CDQ yellowfin sole, by fishery, in 2016.

Source: AKFIN.

<b>Trip Target Name</b>	<b>Discarded</b>	<b>Retained</b>
Atka Mackerel	<1	<1
Pollock - bottom	4	297
Pacific Cod	1,636	315
Alaska Plaice - BSAI	6	260
Other Flatfish - BSAI	<1	
Halibut		
Rockfish	<1	<1
Flathead Sole	92	2,403
Kamchatka Flounder - BSAI	<1	<1
Pollock - midwater	172	415
Rock Sole - BSAI	598	21,646
Sablefish		
Greenland Turbot - BSAI	0	
Arrowtooth Flounder	<1	<1
Yellowfin Sole - BSAI	1,637	105,868

Table 4.4. Yellowfin sole fishery catch-at-age (proportions), 1975-2016.

	males										
	7	8	9	10	11	12	13	14	15	16	17+
<b>1975</b>	0.09	0.24	0.12	0.02	0.01	0.01	0.02	0.00	0.00	0.00	0.00
<b>1976</b>	0.05	0.04	0.14	0.11	0.04	0.02	0.02	0.00	0.01	0.00	0.00
<b>1977</b>	0.03	0.08	0.07	0.12	0.09	0.04	0.01	0.00	0.00	0.00	0.00
<b>1978</b>	0.05	0.09	0.07	0.07	0.07	0.03	0.02	0.01	0.00	0.00	0.00
<b>1979</b>	0.03	0.06	0.11	0.07	0.06	0.04	0.02	0.00	0.00	0.00	0.00
<b>1980</b>	0.04	0.02	0.03	0.05	0.05	0.05	0.03	0.01	0.01	0.01	0.01
<b>1981</b>	0.04	0.04	0.03	0.05	0.07	0.06	0.04	0.02	0.01	0.00	0.00
<b>1982</b>	0.04	0.06	0.05	0.05	0.07	0.07	0.04	0.02	0.01	0.00	0.00
<b>1983</b>	0.06	0.03	0.06	0.05	0.04	0.06	0.05	0.03	0.02	0.01	0.01
<b>1984</b>	0.01	0.06	0.04	0.08	0.04	0.05	0.05	0.10	0.04	0.02	0.01
<b>1985</b>	0.02	0.03	0.06	0.06	0.07	0.06	0.07	0.06	0.03	0.02	0.02
<b>1986</b>	0.03	0.02	0.05	0.06	0.04	0.04	0.04	0.02	0.04	0.04	0.05
<b>1987</b>	0.02	0.05	0.04	0.05	0.04	0.05	0.06	0.04	0.02	0.03	0.13
<b>1988</b>	0.03	0.04	0.09	0.02	0.04	0.04	0.02	0.05	0.04	0.01	0.08
<b>1989</b>	0.00	0.04	0.04	0.06	0.02	0.02	0.04	0.03	0.03	0.03	0.16
<b>1990</b>	0.05	0.01	0.18	0.02	0.05	0.02	0.03	0.04	0.00	0.04	0.07
<b>1991</b>	0.01	0.09	0.01	0.19	0.03	0.06	0.01	0.01	0.02	0.02	0.07
<b>1992</b>	0.01	0.03	0.10	0.03	0.14	0.02	0.02	0.02	0.01	0.01	0.04
<b>1993</b>	0.02	0.01	0.01	0.08	0.01	0.10	0.02	0.02	0.01	0.02	0.09
<b>1994</b>	0.02	0.04	0.03	0.02	0.11	0.01	0.09	0.01	0.03	0.01	0.05
<b>1995</b>	0.03	0.06	0.03	0.01	0.02	0.10	0.00	0.10	0.01	0.01	0.05
<b>1996</b>	0.02	0.06	0.04	0.02	0.01	0.03	0.10	0.02	0.07	0.01	0.06
<b>1997</b>	0.02	0.02	0.05	0.04	0.03	0.04	0.02	0.10	0.02	0.05	0.09
<b>1998</b>	0.02	0.02	0.05	0.04	0.03	0.03	0.01	0.02	0.03	0.01	0.12
<b>1999</b>	0.00	0.02	0.01	0.02	0.04	0.05	0.03	0.02	0.03	0.04	0.11
<b>2000</b>	0.00	0.02	0.05	0.01	0.02	0.05	0.03	0.03	0.02	0.05	0.13
<b>2001</b>	0.01	0.02	0.01	0.03	0.03	0.01	0.02	0.03	0.03	0.02	0.16
<b>2002</b>	0.00	0.02	0.03	0.04	0.09	0.03	0.02	0.05	0.01	0.01	0.14
<b>2003</b>	0.01	0.08	0.04	0.03	0.02	0.04	0.02	0.02	0.03	0.02	0.15
<b>2004</b>	0.01	0.02	0.09	0.02	0.02	0.02	0.03	0.01	0.01	0.02	0.19
<b>2005</b>	0.01	0.04	0.02	0.08	0.02	0.02	0.03	0.04	0.02	0.01	0.14
<b>2006</b>	0.06	0.05	0.03	0.05	0.06	0.02	0.01	0.02	0.03	0.01	0.09
<b>2007</b>	0.02	0.06	0.03	0.03	0.05	0.06	0.02	0.02	0.03	0.02	0.12
<b>2008</b>	0.02	0.03	0.06	0.02	0.03	0.02	0.07	0.01	0.02	0.02	0.11
<b>2009</b>	0.01	0.04	0.03	0.05	0.03	0.01	0.02	0.04	0.02	0.02	0.13
<b>2010</b>	0.06	0.03	0.06	0.02	0.05	0.04	0.02	0.01	0.02	0.01	0.10
<b>2011</b>	0.02	0.07	0.03	0.03	0.03	0.04	0.01	0.01	0.01	0.03	0.14
<b>2012</b>	0.02	0.04	0.08	0.04	0.03	0.03	0.04	0.04	0.01	0.02	0.16
<b>2013</b>	0.02	0.00	0.03	0.08	0.05	0.06	0.04	0.04	0.03	0.01	0.09
<b>2014</b>	0.02	0.04	0.03	0.04	0.06	0.02	0.04	0.04	0.04	0.01	0.13
<b>2015</b>	0.01	0.03	0.04	0.02	0.02	0.05	0.02	0.05	0.02	0.04	0.16
<b>2016</b>	0.02	0.03	0.03	0.05	0.04	0.02	0.05	0.03	0.02	0.01	0.17

Table 4.4- continued.

<b>females</b>	<b>7</b>	<b>8</b>	<b>9</b>	<b>10</b>	<b>11</b>	<b>12</b>	<b>13</b>	<b>14</b>	<b>15</b>	<b>16</b>	<b>17+</b>
<b>1975</b>	0.05	0.14	0.09	0.05	0.02	0.02	0.02	0.00	0.01	0.00	0.00
<b>1976</b>	0.04	0.07	0.17	0.10	0.07	0.01	0.01	0.00	0.01	0.00	0.00
<b>1977</b>	0.07	0.16	0.11	0.02	0.01	0.01	0.00	0.00	0.00	0.00	0.00
<b>1978</b>	0.05	0.13	0.12	0.09	0.09	0.03	0.01	0.01	0.00	0.00	0.00
<b>1979</b>	0.03	0.07	0.12	0.12	0.08	0.06	0.03	0.01	0.01	0.00	0.00
<b>1980</b>	0.06	0.04	0.06	0.11	0.10	0.07	0.07	0.04	0.02	0.01	0.03
<b>1981</b>	0.06	0.06	0.03	0.05	0.09	0.11	0.07	0.05	0.02	0.01	0.01
<b>1982</b>	0.03	0.07	0.06	0.05	0.09	0.09	0.05	0.03	0.02	0.01	0.00
<b>1983</b>	0.07	0.05	0.08	0.04	0.05	0.07	0.06	0.05	0.03	0.02	0.01
<b>1984</b>	0.03	0.04	0.05	0.09	0.04	0.06	0.05	0.06	0.03	0.01	0.02
<b>1985</b>	0.02	0.02	0.06	0.05	0.07	0.06	0.05	0.06	0.05	0.01	0.01
<b>1986</b>	0.03	0.03	0.04	0.09	0.07	0.06	0.04	0.03	0.04	0.03	0.06
<b>1987</b>	0.01	0.03	0.02	0.04	0.05	0.05	0.06	0.05	0.03	0.03	0.09
<b>1988</b>	0.02	0.03	0.07	0.02	0.04	0.05	0.04	0.06	0.05	0.02	0.12
<b>1989</b>	0.00	0.04	0.05	0.05	0.03	0.03	0.05	0.02	0.05	0.05	0.15
<b>1990</b>	0.02	0.01	0.13	0.03	0.06	0.03	0.02	0.01	0.01	0.08	0.09
<b>1991</b>	0.01	0.07	0.01	0.11	0.04	0.04	0.01	0.03	0.04	0.02	0.09
<b>1992</b>	0.01	0.02	0.09	0.02	0.14	0.04	0.04	0.01	0.03	0.02	0.12
<b>1993</b>	0.02	0.01	0.02	0.09	0.01	0.12	0.03	0.03	0.02	0.03	0.18
<b>1994</b>	0.02	0.03	0.03	0.03	0.16	0.00	0.10	0.01	0.04	0.02	0.13
<b>1995</b>	0.04	0.06	0.02	0.01	0.02	0.10	0.00	0.16	0.01	0.03	0.12
<b>1996</b>	0.01	0.04	0.06	0.02	0.03	0.03	0.07	0.01	0.11	0.01	0.11
<b>1997</b>	0.02	0.02	0.06	0.03	0.02	0.03	0.03	0.10	0.01	0.06	0.10
<b>1998</b>	0.02	0.03	0.08	0.04	0.02	0.04	0.04	0.12	0.01	0.07	0.13
<b>1999</b>	0.01	0.02	0.03	0.02	0.08	0.05	0.03	0.04	0.04	0.07	0.23
<b>2000</b>	0.00	0.01	0.05	0.03	0.03	0.07	0.08	0.05	0.02	0.04	0.22
<b>2001</b>	0.01	0.02	0.05	0.08	0.05	0.04	0.07	0.05	0.04	0.02	0.16
<b>2002</b>	0.01	0.02	0.03	0.04	0.06	0.04	0.03	0.04	0.05	0.02	0.21
<b>2003</b>	0.00	0.05	0.04	0.03	0.04	0.08	0.03	0.02	0.02	0.03	0.19
<b>2004</b>	0.01	0.01	0.10	0.05	0.03	0.02	0.05	0.02	0.01	0.05	0.19
<b>2005</b>	0.02	0.03	0.03	0.08	0.03	0.03	0.04	0.06	0.03	0.01	0.19
<b>2006</b>	0.07	0.05	0.03	0.04	0.14	0.02	0.00	0.01	0.03	0.01	0.10
<b>2007</b>	0.01	0.04	0.03	0.02	0.04	0.08	0.03	0.03	0.03	0.03	0.17
<b>2008</b>	0.02	0.04	0.04	0.05	0.03	0.03	0.07	0.04	0.02	0.03	0.17
<b>2009</b>	0.02	0.03	0.06	0.06	0.03	0.05	0.04	0.05	0.03	0.03	0.17
<b>2010</b>	0.04	0.03	0.04	0.04	0.04	0.03	0.04	0.04	0.06	0.03	0.17
<b>2011</b>	0.02	0.05	0.04	0.05	0.04	0.06	0.03	0.03	0.03	0.04	0.16
<b>2012</b>	0.02	0.03	0.05	0.04	0.05	0.02	0.05	0.02	0.01	0.02	0.16
<b>2013</b>	0.01	0.02	0.04	0.07	0.06	0.07	0.05	0.04	0.02	0.03	0.14
<b>2014</b>	0.01	0.02	0.04	0.05	0.05	0.06	0.04	0.03	0.04	0.03	0.18
<b>2015</b>	0.01	0.02	0.02	0.04	0.03	0.08	0.04	0.04	0.03	0.04	0.18
<b>2016</b>	0.02	0.04	0.04	0.03	0.04	0.03	0.06	0.03	0.02	0.01	0.21

Table 4.5 Yellowfin sole biomass estimates (t) from the annual Bering Sea shelf bottom trawl survey and upper and lower 95% confidence intervals.

Year	Total	Lower CI	Upper CI
1982	3,377,800	2,571,000	4,184,600
1983	3,535,300	2,958,100	4,112,400
1984	3,141,200	2,636,800	3,645,600
1985	2,443,700	1,563,400	3,324,000
1986	1,909,900	1,480,700	2,339,000
1987	2,613,100	2,051,800	3,174,400
1988	2,402,400	1,808,400	2,996,300
1989	2,316,300	1,836,700	2,795,800
1990	2,183,800	1,886,200	2,479,400
1991	2,393,300	2,116,000	2,670,700
1992	2,172,900	1,898,900	2,690,600
1993	2,465,400	2,151,500	2,779,300
1994	2,610,500	2,266,800	2,954,100
1995	2,009,700	1,724,800	2,294,600
1996	2,298,600	1,749,900	2,847,300
1997	2,163,400	1,907,900	2,418,900
1998	2,329,600	2,033,130	2,626,070
1999	1,306,470	1,118,800	1,494,150
2000	1,581,900	1,382,000	1,781,800
2001	1,863,700	1,605,000	2,122,300
2002	2,016,700	1,740,700	2,292,700
2003	2,239,600	1,822,700	2,656,600
2004	2,530,600	2,147,900	2,913,300
2005	2,823,500	2,035,800	3,499,800
2006	2,133,070	1,818,253	2,447,932
2007	2,152,738	1,775,191	2,530,285
2008	2,099,521	1,599,100	2,600,000
2009	1,739,238	1,435,188	2,043,288
2010	2,367,830	1,807,430	2,928,230
2011	2,403,021	1,926,371	2,879,671
2012	1,951,400	1,675,982	2,226,819
2013	2,279,004	1,934,134	2,623,874
2014	2,512,250	2,058,018	2,966,482
2015	1,932,347	1,644,043	2,220,651
2016	2,859,811	2,532,202	3,187,421
2017	2,787,688	2,310,198	3,265,178

Table 4.6. Yellowfin sole population numbers-at-age (millions) estimated from the annual bottom trawl surveys, 1982-2016.

<b>females</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>5</b>	<b>6</b>	<b>7</b>	<b>8</b>	<b>9</b>	<b>10</b>	<b>11</b>	<b>12</b>	<b>13</b>	<b>14</b>	<b>15</b>	<b>16</b>	<b>17+</b>
<b>1979</b>	21	113	150	442	616	386	555	801	626	528	219	274	59	35	29	15
<b>1980</b>	1	92	342	518	800	1055	413	661	880	651	765	285	113	33	23	23
<b>1981</b>	0	20	195	839	692	1321	1155	261	477	744	527	311	168	55	23	45
<b>1982</b>	38	183	349	1211	1485	1424	1619	843	829	832	704	409	246	159	51	84
<b>1983</b>	0	5	59	154	751	1413	843	1065	936	753	1155	866	295	160	60	54
<b>1984</b>	0	53	278	264	427	745	841	1111	1080	941	541	583	480	239	174	133
<b>1985</b>	0	3	105	442	587	406	632	915	441	518	545	384	298	321	205	127
<b>1986</b>	0	8	24	219	349	666	279	574	519	377	284	318	196	250	136	259
<b>1987</b>	0	0	70	120	803	458	843	259	376	599	356	449	243	270	247	688
<b>1988</b>	0	0	7	370	71	1495	560	557	184	239	351	208	360	273	219	886
<b>1989</b>	0	0	14	98	718	234	1337	593	446	74	179	308	234	238	183	565
<b>1990</b>	0	0	70	102	325	1066	192	1257	408	482	101	72	107	78	231	605
<b>1991</b>	0	10	127	248	123	405	896	151	1263	213	525	63	128	87	123	807
<b>1992</b>	0	19	247	485	520	213	286	938	94	825	75	309	129	137	170	715
<b>1993</b>	0	24	100	357	634	434	269	224	1314	78	866	157	165	69	68	674
<b>1994</b>	0	54	95	223	518	905	555	482	284	1170	516	44	274	142	42	588
<b>1995</b>	0	19	153	288	181	889	627	274	135	25	634	21	561	104	80	512
<b>1996</b>	0	16	154	809	288	279	434	517	206	146	151	602	116	637	47	619
<b>1997</b>	0	18	324	502	725	256	239	506	228	114	176	184	500	44	314	533
<b>1998</b>	0	10	83	479	420	900	260	203	370	413	369	170	176	265	67	1167
<b>1999</b>	0	3	65	198	175	185	727	104	107	245	190	186	72	102	175	425
<b>2000</b>	0	11	54	248	208	304	444	537	189	198	237	219	65	117	145	572
<b>2001</b>	0	1	71	239	522	248	403	415	654	374	83	191	154	127	189	617
<b>2002</b>	0	16	123	170	255	778	346	290	229	457	221	91	307	116	152	805
<b>2003</b>	0	15	115	241	251	287	1143	225	279	286	251	103	115	170	168	943
<b>2004</b>	10	33	192	430	560	441	217	966	221	212	218	219	106	20	167	1020
<b>2005</b>	0	53	167	194	602	433	213	487	834	196	144	191	324	170	53	1332
<b>2006</b>	0	67	302	376	276	634	470	176	325	738	133	133	71	156	175	514
<b>2007</b>	0	37	515	348	376	277	504	308	124	227	504	119	137	127	105	724
<b>2008</b>	0	24	115	736	621	546	359	355	198	117	259	350	153	79	85	732
<b>2009</b>	5	38	204	204	1187	609	488	259	210	218	129	138	196	88	43	444
<b>2010</b>	0	33	328	386	438	895	554	517	329	335	155	166	135	173	99	684
<b>2011</b>	0	14	243	539	707	463	769	410	457	204	226	149	142	145	186	619
<b>2012</b>	10	50	229	394	503	293	243	752	256	334	106	156	37	150	128	547
<b>2013</b>	0	4	88	269	420	531	256	221	409	406	358	119	135	133	133	770
<b>2014</b>	0	0	37	421	384	248	420	231	228	523	341	160	144	228	34	819
<b>2015</b>	0	23	3	167	467	350	308	287	249	149	282	258	135	99	80	592
<b>2016</b>	1	32	71	45	164	743	565	403	364	300	144	245	230	140	163	1027

Table 4.6.(continued)

year/age	males															
	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17+
1979	21	115	143	390	381	303	583	847	604	406	349	247	54	76	29	36
1980	20	78	306	632	853	1221	457	558	616	568	444	370	147	18	8	8
1981	0	50	200	1047	640	1280	858	394	372	546	534	266	66	83	55	12
1982	89	193	428	1780	1781	1059	1673	644	774	463	471	482	302	8	24	8
1983	0	1	65	183	724	1729	808	1049	676	699	722	566	425	550	77	51
1984	0	68	246	323	497	734	830	612	788	718	358	379	201	316	122	106
1985	0	41	172	419	559	263	652	527	401	451	360	224	260	157	112	65
1986	0	13	47	108	373	652	262	327	284	335	211	205	115	210	82	252
1987	0	5	41	106	838	467	673	445	328	277	210	147	106	142	185	600
1988	0	2	10	435	49	1163	553	443	85	187	28	177	336	189	28	599
1989	0	2	23	181	788	177	1306	513	357	135	50	103	54	204	35	478
1990	0	11	47	121	316	888	195	1144	318	263	40	65	67	24	55	389
1991	0	0	103	354	139	275	1046	68	1137	328	244	74	64	60	53	420
1992	0	0	146	445	566	262	226	812	114	907	193	213	12	12	61	607
1993	0	20	52	233	646	393	279	247	1096	69	842	53	53	50	0	341
1994	4	22	71	166	427	953	656	308	191	822	26	622	46	132	11	303
1995	0	0	169	120	270	667	565	94	179	75	478	13	603	49	24	418
1996	0	76	95	837	244	227	425	344	331	141	139	399	61	449	125	495
1997	0	10	214	425	798	181	184	446	245	194	214	108	514	79	264	416
1998	0	48	70	351	569	832	159	226	204	272	346	140	157	191	113	814
1999	0	5	100	142	225	243	575	146	94	309	269	75	53	28	119	425
2000	0	0	36	219	259	143	509	583	78	215	133	77	92	78	66	547
2001	0	0	87	141	652	341	375	357	562	208	87	158	65	73	140	432
2002	0	58	72	158	309	758	318	333	262	442	194	120	220	161	133	507
2003	0	24	95	178	258	251	1074	238	363	53	284	173	10	71	57	682
2004	4	63	114	469	447	199	395	993	263	81	195	223	103	47	249	456
2005	0	49	166	187	474	476	204	288	972	123	142	121	133	69	93	726
2006	0	101	173	348	332	505	393	288	298	384	116	155	89	39	11	590
2007	0	58	481	352	405	284	545	209	166	252	338	101	133	72	59	620
2008	0	10	99	662	462	483	344	453	225	144	185	329	63	66	35	581
2009	0	65	144	289	946	462	555	248	249	217	78	31	195	30	29	363
2010	0	78	199	418	371	1032	462	510	171	189	159	53	117	151	78	678
2011	1	7	150	385	482	358	792	398	224	176	77	81	136	103	157	440
2012	0	69	274	352	344	273	238	425	297	179	98	67	91	34	100	2
2013	0	7	92	366	384	481	211	268	445	200	200	33	89	100	118	612
2014	0	0	0	9	366	396	286	338	310	251	400	206	193	20	192	841
2015	1	29	36	131	426	332	301	312	318	48	180	131	80	1	80	492
2016	0	43	85	20	142	704	544	402	367	125	117	227	180	88	35	859

Table 4.7 Occurrence of yellowfin sole in the Bering Sea trawl survey and collections of length and age structures and the number of otoliths aged from each survey.

Year	Total Hauls	Hauls w/Len	Number lengths	Hauls w/otoliths	Hauls w/ages	Number of otoliths	Number ages
<b>1982</b>	334	246	37023	35	35	744	744
<b>1983</b>	353	256	33924	37	37	709	709
<b>1984</b>	355	271	33894	56	56	821	796
<b>1985</b>	357	261	33824	44	43	810	802
<b>1986</b>	354	249	30470	34	34	739	739
<b>1987</b>	357	224	31241	16	16	798	798
<b>1988</b>	373	254	27138	14	14	543	543
<b>1989</b>	374	236	29672	24	24	740	740
<b>1990</b>	371	251	30257	28	28	792	792
<b>1991</b>	372	248	27986	26	26	742	742
<b>1992</b>	356	229	23628	16	16	606	606
<b>1993</b>	375	242	26651	20	20	549	549
<b>1994</b>	375	269	24448	14	14	526	522
<b>1995</b>	376	254	22116	20	20	654	647
<b>1996</b>	375	247	27505	16	16	729	721
<b>1997</b>	376	262	26034	11	11	470	466
<b>1998</b>	375	310	34509	15	15	575	570
<b>1999</b>	373	276	28431	31	31	777	770
<b>2000</b>	372	255	24880	20	20	517	511
<b>2001</b>	375	251	26558	25	25	604	593
<b>2002</b>	375	246	26309	32	32	738	723
<b>2003</b>	376	241	27135	37	37	699	695
<b>2004</b>	375	251	26103	26	26	725	712
<b>2005</b>	373	251	24658	34	34	644	635
<b>2006</b>	376	246	28470	39	39	428	426
<b>2007</b>	376	247	24790	66	66	779	772
<b>2008</b>	375	238	25848	65	65	858	830
<b>2009</b>	376	235	22018	70	70	784	752
<b>2010</b>	376	228	20619	77	77	841	827
<b>2011</b>	376	228	21665	65	64	784	753
<b>2012</b>	376	242	23519	72	72	993	973
<b>2013</b>	376	232	23261	70	70	821	803
<b>2014</b>	376	219	20229	52	52	799	790
<b>2015</b>	376	223	20830	73	73	878	875
<b>2016</b>	376	242	26674	69	69	884	876
<b>2017</b>	376	258	25767	78		896	

Table 4.8 Total tonnage of yellowfin sole caught in resource assessment surveys in the eastern Bering Sea from 1977-2017.

<b>Year</b>	<b>Research catch (t)</b>
1977	60
1978	71
1979	147
1980	92
1981	74
1982	158
1983	254
1984	218
1985	105
1986	68
1987	92
1988	138
1989	148
1990	129
1991	118
1992	60
1993	95
1994	91
1995	95
1996	72
1997	76
1998	79
1999	61
2000	72
2001	75
2002	76
2003	78
2004	114
2005	94
2006	74
2007	74
2008	69
2009	60
2010	79
2011	77
2012	64
2013	75
2014	81
2015	64
2016	98
2017	98

Table 4.9. Mean length and weight at age for yellowfin sole (unsmoothed).

	average mean length at age (cm)																			
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
males	7	11	12	14	17	20	22	24	26	27	28	29	30	30	31	31	31	32	32	32
females	10	13	15	17	20	22	25	27	29	30	31	32	33	33	33	34	34	33	34	34
	weight at age (g) males																			
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
1954	0	4	15	34	60	91	125	160	195	230	263	294	322	348	372	393	412	429	444	481
1955	0	4	15	34	60	91	125	160	195	230	263	294	322	348	372	393	412	429	444	481
	0	4	15	34	60	91	125	160	195	230	263	294	322	348	372	393	412	429	444	481
1974	0	4	15	34	60	91	125	160	195	230	263	294	322	348	372	393	412	429	444	481
1975	4	14	18	32	54	85	120	156	193	225	253	280	303	324	330	344	355	366	390	423
1976	4	14	18	32	54	85	120	156	193	225	253	280	303	324	330	344	355	366	390	423
1977	4	14	18	32	54	85	120	156	193	225	253	280	303	324	330	344	355	366	390	423
1978	4	14	18	32	54	85	120	156	193	225	253	280	303	324	330	344	355	366	390	423
1979	4	14	18	32	54	85	120	156	193	225	253	280	303	324	330	344	355	366	390	423
1980	4	14	18	32	54	85	120	156	193	225	253	280	303	324	330	344	355	366	390	423
1981	4	14	18	32	54	85	120	156	193	225	253	280	303	324	330	344	355	366	390	423
1982	4	11	25	50	83	112	133	142	158	182	196	212	218	249	403	386	386	455	532	408
1983	4	5	5	23	57	95	156	156	155	176	212	227	227	254	262	287	271	370	370	408
1984	4	10	20	31	57	121	150	181	202	193	202	213	246	252	257	262	282	415	290	370
1985	4	11	23	32	51	84	148	186	214	227	228	246	277	267	283	305	407	389	532	387
1986	4	9	18	27	34	61	98	176	217	233	239	229	271	263	258	324	265	318	300	370
1987	4	8	14	17	27	53	97	157	211	226	260	267	311	309	276	291	307	296	329	394
1988	4	7	10	18	45	75	76	138	207	242	261	304	301	297	339	304	308	315	326	386
1989	4	7	10	27	47	72	142	130	179	244	270	351	338	352	317	302	391	309	361	348
1990	4	9	16	22	44	64	98	120	175	197	273	323	341	326	337	286	348	353	343	388
1991	4	9	17	29	51	75	100	132	180	212	266	267	325	355	326	359	352	304	532	381
1992	4	9	17	28	53	86	97	125	174	208	239	264	306	508	407	395	344	360	406	360
1993	4	9	18	45	56	93	135	145	206	209	238	265	387	303	349	363	376	349	342	384
1994	4	23	32	53	76	92	116	182	198	207	259	336	311	345	345	407	356	479	349	424
1995	4	10	19	32	59	88	110	154	177	207	249	258	336	294	319	377	367	383	401	448
1996	4	10	19	32	54	107	134	163	184	215	221	264	281	295	314	326	333	418	326	435
1997	4	8	14	37	64	75	149	174	185	239	231	248	261	303	349	336	384	370	346	444
1998	4	10	20	27	49	79	113	156	208	207	259	262	289	301	291	332	330	354	350	392
1999	4	6	7	18	37	63	95	123	170	171	245	281	269	269	347	330	395	350	350	450
2000	4	10	20	36	32	64	88	133	161	284	233	271	302	255	291	331	351	349	373	385
2001	4	9	16	27	38	51	91	152	161	198	268	240	280	299	292	320	343	357	430	434
2002	4	9	18	21	57	59	81	134	188	204	241	248	269	306	303	343	336	304	368	414
2003	4	11	22	39	53	83	109	161	179	251	248	304	263	468	330	339	305	339	352	405
2004	4	7	20	40	64	94	157	157	213	266	334	310	297	356	360	338	387	414	443	446
2005	4	11	24	44	77	110	136	170	201	262	278	332	366	308	328	350	375	347	349	434
2006	4	10	19	36	71	124	139	180	207	237	233	315	330	380	385	446	369	335	382	390
2007	4	10	19	36	63	107	140	181	208	248	291	286	311	340	375	342	353	369	422	430
2008	4	8	13	29	50	91	113	181	194	252	262	289	306	364	366	369	372	374	417	481
2009	4	7	11	20	39	74	112	133	194	273	270	302	348	321	379	320	405	370	391	460
2010	4	14	18	32	54	85	120	156	193	225	253	280	303	324	330	344	355	366	390	423
2011	4	14	17	25	47	81	134	164	174	305	283	330	291	346	332	344	389	364	375	400
2012	4	14	12	27	48	83	126	181	214	249	274	296	295	341	342	382	380	388	396	400
2013	4	14	13	21	40	72	122	179	227	259	278	320	273	379	357	379	407	390	366	400
2014	4	8	11	44	34	75	150	195	246	296	313	314	330	273	385	387	400	478	436	400
2015	4	8	11	44	34	75	150	195	246	296	313	314	330	273	385	387	400	478	436	400
2016	4	8	43	57	63	82	116	171	253	319	318	331	338	346	381	383	408	434	413	460
2017	4	8	43	57	63	82	116	171	253	319	318	331	338	346	381	383	408	434	413	460

Table 4.9—(continued) Mean length and weight at age for yellowfin sole (unsmoothed).

	females																			
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
1954	4	15	34	60	91	125	160	195	230	263	294	322	348	372	393	412	429	444	481	590
1955	4	15	34	60	91	125	160	195	230	263	294	322	348	372	393	412	429	444	481	590
1974	4	15	34	60	91	125	160	195	230	263	294	322	348	372	393	412	429	444	481	590
1975	8	20	31	55	84	124	165	217	266	301	341	374	407	428	443	480	483	499	590	590
1976	8	20	31	55	84	124	165	217	266	301	341	374	407	428	443	480	483	499	590	590
1977	8	20	31	55	84	124	165	217	266	301	341	374	407	428	443	480	483	499	590	590
1978	8	20	31	55	84	124	165	217	266	301	341	374	407	428	443	480	483	499	590	590
1979	8	20	31	55	84	124	165	217	266	301	341	374	407	428	443	480	483	499	590	590
1980	8	20	31	55	84	124	165	217	266	301	341	374	407	428	443	480	483	499	590	590
1981	8	20	31	55	84	124	165	217	266	301	341	374	407	428	443	480	483	499	590	590
1982	8	20	42	75	98	139	176	214	233	235	289	300	339	336	406	490	417	386	568	590
1983	10	14	26	60	103	162	185	201	243	255	280	329	395	477	539	583	578	630	685	590
1984	14	26	33	57	110	156	177	222	246	294	338	332	325	422	436	458	497	665	654	590
1985	11	16	28	46	77	177	202	251	286	302	323	371	370	421	425	499	624	600	620	590
1986	14	27	23	41	71	103	173	239	284	338	342	350	402	351	391	422	440	455	611	590
1987	10	14	20	47	55	127	179	256	317	324	373	373	385	384	422	412	458	436	523	590
1988	9	12	16	34	66	85	159	237	286	307	378	396	404	388	415	437	429	485	578	590
1989	12	21	33	67	71	112	133	197	279	339	402	430	449	456	456	456	578	476	516	590
1990	11	17	24	38	65	99	126	197	243	321	449	450	416	446	464	455	471	523	569	590
1991	11	16	23	58	56	100	142	156	238	310	370	457	446	473	474	490	492	484	598	590
1992	12	21	29	55	85	121	177	176	283	305	284	352	435	516	459	484	519	459	547	590
1993	15	28	35	64	93	155	165	232	244	301	333	368	442	452	497	499	471	538	586	590
1994	20	46	53	86	87	125	155	235	276	284	337	396	351	461	464	480	476	514	553	590
1995	12	20	28	60	84	123	160	217	284	332	340	443	384	414	454	439	619	482	589	590
1996	11	16	36	51	108	137	167	202	222	311	318	334	405	399	432	534	462	523	558	590
1997	16	34	33	72	85	157	200	236	260	292	353	373	401	469	440	490	431	515	600	590
1998	10	14	36	51	90	104	177	237	278	279	318	370	416	405	403	448	407	532	581	590
1999	9	12	18	37	67	103	131	239	284	296	328	348	384	396	416	461	502	477	639	590
2000	11	16	33	33	91	81	158	175	237	306	310	373	401	440	422	494	506	483	636	590
2001	6	6	32	41	57	83	148	179	255	305	357	372	447	415	420	422	476	522	598	590
2002	11	18	27	48	65	87	120	224	243	261	337	346	374	408	434	452	505	489	585	590
2003	9	12	31	53	86	124	156	213	289	303	344	407	425	399	434	365	438	457	536	590
2004	9	18	43	63	101	168	172	245	299	346	380	407	483	543	450	461	464	500	604	590
2005	14	26	44	78	114	152	213	238	277	337	347	397	439	461	531	522	438	539	629	590
2006	9	13	40	82	125	153	204	245	319	314	375	370	533	460	476	865	480	537	691	590
2007	11	16	36	66	115	173	198	244	316	311	362	358	417	461	462	497	491	611	640	590
2008	13	24	28	54	98	129	199	226	286	320	355	384	442	434	471	530	530	552	630	590
2009	6	9	18	45	69	127	163	239	306	322	375	416	381	413	473	736	539	491	679	590
2010	8	20	31	55	84	124	165	217	266	301	341	374	407	428	443	480	483	499	590	590
2011	8	18	25	56	80	126	188	205	327	332	372	403	415	440	426	369	491	542	590	590
2012	8	12	26	49	81	144	169	256	313	341	349	445	459	471	476	444	527	525	590	590
2013	8	12	21	35	92	125	182	261	305	364	410	426	464	456	451	507	494	532	590	590
2014	6	8	11	18	34	74	145	203	260	305	376	367	405	410	488	519	483	581	548	590
2015	6	8	11	18	34	74	145	203	260	305	376	367	405	410	488	519	483	581	548	590
2016	6	8	32	50	66	74	112	186	338	372	412	408	455	456	485	508	515	532	555	590
2017	6	8	32	50	66	74	112	186	338	372	412	408	455	456	485	508	515	532	555	590

Table 4.10. Female yellowfin sole proportion mature at age from Nichol (1995) and TenBrink and Wilderbuer (2015).

Age	1992, 1993 samples	2012 samples	Combined
1	0.00	0	0
2	0.00	0	0
3	0.001	0	0
4	0.004	0	0
5	0.008	0	0
6	0.020	0.01	0.01
7	0.046	0.03	0.04
8	0.104	0.09	0.10
9	0.217	0.21	0.21
10	0.397	0.43	0.41
11	0.612	0.68	0.65
12	0.790	0.86	0.83
13	0.899	0.94	0.92
14	0.955	0.98	0.97
15	0.981	0.99	0.99
16	0.992	1.0	1.0
17	0.997	1.0	1.0
18	1.0	1.0	1.0
19	1.0	1.0	1.0
20	1.0	1.0	1.0

Table 4.11. Key equations used in the population dynamics model.

$$N_{t,1} = R_t = R_0 e^{\tau_t}, \quad \tau_t \sim N(0, \sigma^2_R) \quad \text{Recruitment 1956-75}$$

$$N_{t,1} = R_t = R_\gamma e^{\tau_t}, \quad \tau_t \sim N(0, \sigma^2_R) \quad \text{Recruitment 1976-96}$$

$$C_{t,a} = \frac{F_{t,a}}{Z_{t,a}} (1 - e^{-z_{t,a}}) N_{t,a} \quad \text{Catch in year } t \text{ for age } a \text{ fish}$$

$$N_{t+1,a+1} = N_{t,a} e^{-z_{t,a}} \quad \text{Numbers of fish in year } t+1 \text{ at age } a$$

$$N_{t+1,A} = N_{t,A-1} e^{-z_{t,A-1}} + N_{t,A} e^{-z_{t,A}} \quad \text{Numbers of fish in the "plus group"}$$

$$S_t = \sum N_{t,a} W_{t,a} \phi_a \quad \text{Spawning biomass}$$

$$Z_{t,a} = F_{t,a} + M \quad \text{Total mortality in year } t \text{ at age } a$$

$$F_{t,a} = s_a \mu^F \exp^{\varepsilon^F_t}, \quad \varepsilon^F_t \sim N(o, \sigma^2_F) \quad \text{Fishing mortality}$$

$$s_a = \frac{1}{1 + (e^{-\alpha + \beta a})} \quad \text{Age-specific fishing selectivity}$$

$$C_t = \sum C_{t,a} \quad \text{Total catch in numbers}$$

$$P_{t,a} = \frac{C_{t,a}}{C_t} \quad \text{Proportion at age in catch}$$

$$SurB_t = q \sum N_{t,a} W_{t,a} v_a \quad \text{Survey biomass}$$

Table 4.11—continued.

$$qprior = \lambda \frac{0.5(\ln q_{est,t} - \ln q_{prior})^2}{\sigma_q^2} \quad \text{survey catchability prior (when estimated)}$$

$$mprior = \lambda \frac{0.5(\ln m_{est} - \ln m_{prior})^2}{\sigma_m^2} \quad \text{natural mortality prior (when estimated)}$$

$$reclike = \lambda \left( \sum_{i=1965}^{endyear} \bar{R} - R_i \right)^2 + \sum_{a=1}^{20} (\bar{R}_{init} - R_{init,a})^2 + \frac{1}{2((\sum_{i=1965}^{endyear} \bar{R} - R_i) \frac{1}{n+1})} \quad \text{recruitment likelihood}$$

$$catchlike = \lambda \sum_{i=startyear}^{endyear} (\ln C_{obs,i} - \ln C_{est,i})^2 \quad \text{catch likelihood}$$

$$surveylike = \lambda \frac{(\ln B - \ln \hat{B})^2}{2\sigma^2} \quad \text{survey likelihood}$$

$$SurvAgelike = \sum_{i,t} m_t P_{t,a} \ln \frac{\hat{P}_{t,a}}{P_{t,a}} \quad \text{survey age composition likelihood}$$

$$FishAgelike = \sum_{i,t} m_t P_{t,a} \ln \frac{\hat{P}_{t,a}}{P_{t,a}} \quad \text{fishery age composition likelihood}$$

Table 4.12. Variables used in the population dynamics model.

<b>Variables</b>	<b>Description</b>
$R_t$	Age 1 recruitment in year $t$
$R_0$	Geometric mean value of age 1 recruitment, 1956-75
$R_\gamma$	Geometric mean value of age 1 recruitment, 1976-2014
$\tau_t$	Recruitment deviation in year $t$
$N_{t,a}$	Number of fish in year $t$ at age $a$
$C_{t,a}$	Catch numbers of fish in year $t$ at age $a$
$P_{t,a}$	Proportion of the numbers of fish age $a$ in year $t$
$C_t$	Total catch numbers in year $t$
$W_{t,a}$	Mean body weight (kg) of fish age $a$ in year $t$
$\phi_a$	Proportion of mature females at age $a$
$F_{t,a}$	Instantaneous annual fishing mortality of age $a$ fish in year $t$
$M$	Instantaneous natural mortality, assumed constant over all ages and years
$Z_{t,a}$	Instantaneous total mortality for age $a$ fish in year $t$
$s_a$	Age-specific fishing gear selectivity
$\mu^F$	Median year-effect of fishing mortality
$\varepsilon_t^F$	The residual year-effect of fishing mortality
$v_a$	Age-specific survey selectivity
$\alpha$	Slope parameter in the logistic selectivity equation
$\beta$	Age at 50% selectivity parameter in the logistic selectivity equation
$\sigma_t$	Standard error of the survey biomass in year $t$

Table 4.13. Models evaluated for stock productivity in the 2017 stock assessment of yellowfin sole

	<b>Model 14_2</b>	<b>Model 14_1</b>
<b>Years included</b>	1955-2012	1978-2012
<b>Fmsy</b>	0.19	0.12
<b>Bmsy (t)</b>	333,700	454,000
<b>ABC (t)</b>	471,300	277,500
<b>OFL (t)</b> <b>Buffer between ABC and OFL</b>	476,800	306,700
	1%	11%

Table 4-14. Model estimates of annual average fishing mortality for male and female yellowfin sole.

year/age	Females													
	7	8	9	10	11	12	13	14	15	16	17	18	19	20
<b>1964</b>	0.082	0.179	0.269	0.315	0.330	0.335	0.337	0.337	0.337	0.337	0.337	0.337	0.337	0.337
<b>1965</b>	0.012	0.044	0.114	0.189	0.225	0.237	0.240	0.240	0.241	0.241	0.241	0.241	0.241	0.241
<b>1966</b>	0.018	0.053	0.139	0.270	0.375	0.426	0.444	0.449	0.451	0.452	0.452	0.452	0.452	0.452
<b>1967</b>	0.027	0.078	0.194	0.362	0.497	0.562	0.586	0.594	0.597	0.597	0.598	0.598	0.598	0.598
<b>1968</b>	0.011	0.027	0.062	0.129	0.228	0.328	0.397	0.433	0.449	0.455	0.458	0.458	0.458	0.458
<b>1969</b>	0.057	0.205	0.442	0.578	0.617	0.625	0.627	0.627	0.628	0.628	0.628	0.628	0.628	0.628
<b>1970</b>	0.347	0.608	0.650	0.654	0.655	0.655	0.655	0.655	0.655	0.655	0.655	0.655	0.655	0.655
<b>1971</b>	0.383	0.583	0.605	0.607	0.607	0.607	0.607	0.607	0.607	0.607	0.607	0.607	0.607	0.607
<b>1972</b>	0.000	0.000	0.000	0.000	0.000	0.001	0.003	0.010	0.034	0.106	0.250	0.250	0.250	0.250
<b>1973</b>	0.018	0.064	0.178	0.318	0.394	0.419	0.426	0.427	0.428	0.428	0.428	0.428	0.428	0.428
<b>1974</b>	0.004	0.014	0.038	0.078	0.116	0.135	0.143	0.145	0.146	0.146	0.146	0.146	0.146	0.146
<b>1975</b>	0.018	0.056	0.101	0.123	0.129	0.130	0.131	0.131	0.131	0.131	0.131	0.131	0.131	0.131
<b>1976</b>	0.011	0.029	0.059	0.091	0.111	0.121	0.124	0.126	0.126	0.126	0.126	0.126	0.126	0.126
<b>1977</b>	0.026	0.036	0.046	0.052	0.055	0.057	0.058	0.059	0.059	0.059	0.059	0.059	0.059	0.059
<b>1978</b>	0.032	0.063	0.091	0.107	0.113	0.115	0.115	0.115	0.116	0.116	0.116	0.116	0.116	0.116
<b>1979</b>	0.019	0.037	0.053	0.062	0.065	0.067	0.067	0.067	0.067	0.067	0.067	0.067	0.067	0.067
<b>1980</b>	0.014	0.024	0.038	0.051	0.062	0.069	0.073	0.075	0.076	0.076	0.076	0.076	0.076	0.076
<b>1981</b>	0.014	0.024	0.036	0.046	0.053	0.057	0.058	0.059	0.060	0.060	0.060	0.060	0.060	0.060
<b>1982</b>	0.015	0.026	0.035	0.041	0.043	0.044	0.044	0.044	0.044	0.044	0.044	0.044	0.044	0.044
<b>1983</b>	0.022	0.034	0.042	0.044	0.045	0.046	0.046	0.046	0.046	0.046	0.046	0.046	0.046	0.046
<b>1984</b>	0.017	0.034	0.052	0.063	0.068	0.070	0.071	0.071	0.071	0.071	0.071	0.071	0.071	0.071
<b>1985</b>	0.020	0.041	0.067	0.088	0.099	0.103	0.105	0.106	0.106	0.106	0.106	0.106	0.106	0.106
<b>1986</b>	0.025	0.058	0.084	0.094	0.097	0.098	0.098	0.098	0.098	0.098	0.098	0.098	0.098	0.098
<b>1987</b>	0.009	0.021	0.042	0.066	0.082	0.091	0.094	0.096	0.096	0.096	0.096	0.096	0.096	0.096
<b>1988</b>	0.011	0.032	0.068	0.100	0.115	0.120	0.122	0.122	0.122	0.122	0.122	0.122	0.122	0.122
<b>1989</b>	0.004	0.013	0.035	0.064	0.082	0.089	0.091	0.091	0.092	0.092	0.092	0.092	0.092	0.092
<b>1990</b>	0.004	0.012	0.024	0.034	0.038	0.039	0.040	0.040	0.040	0.040	0.040	0.040	0.040	0.040
<b>1991</b>	0.006	0.014	0.025	0.035	0.041	0.043	0.044	0.045	0.045	0.045	0.045	0.045	0.045	0.045
<b>1992</b>	0.011	0.025	0.044	0.059	0.068	0.071	0.073	0.073	0.073	0.073	0.074	0.074	0.074	0.074
<b>1993</b>	0.008	0.014	0.022	0.032	0.041	0.047	0.051	0.053	0.054	0.055	0.055	0.055	0.055	0.055
<b>1994</b>	0.014	0.031	0.049	0.060	0.064	0.065	0.066	0.066	0.066	0.066	0.066	0.066	0.066	0.066
<b>1995</b>	0.016	0.033	0.047	0.054	0.056	0.057	0.057	0.057	0.057	0.057	0.057	0.057	0.057	0.057
<b>1996</b>	0.017	0.029	0.042	0.051	0.057	0.060	0.061	0.062	0.062	0.062	0.062	0.062	0.062	0.062
<b>1997</b>	0.021	0.037	0.056	0.072	0.082	0.088	0.091	0.093	0.093	0.093	0.094	0.094	0.094	0.094
<b>1998</b>	0.016	0.029	0.041	0.048	0.051	0.053	0.053	0.053	0.053	0.054	0.054	0.054	0.054	0.054
<b>1999</b>	0.002	0.007	0.016	0.027	0.035	0.039	0.041	0.041	0.041	0.041	0.041	0.041	0.041	0.041
<b>2000</b>	0.003	0.008	0.019	0.033	0.043	0.048	0.050	0.050	0.050	0.050	0.050	0.050	0.050	0.050
<b>2001</b>	0.006	0.013	0.023	0.031	0.035	0.037	0.037	0.037	0.038	0.038	0.038	0.038	0.038	0.038
<b>2002</b>	0.003	0.008	0.018	0.030	0.037	0.040	0.041	0.042	0.042	0.042	0.042	0.042	0.042	0.042
<b>2003</b>	0.005	0.012	0.023	0.032	0.037	0.039	0.040	0.040	0.040	0.040	0.040	0.040	0.040	0.040
<b>2004</b>	0.006	0.013	0.022	0.030	0.034	0.036	0.037	0.037	0.037	0.037	0.037	0.037	0.037	0.037
<b>2005</b>	0.010	0.019	0.029	0.038	0.044	0.047	0.048	0.049	0.049	0.049	0.049	0.049	0.049	0.049
<b>2006</b>	0.032	0.042	0.046	0.046	0.046	0.046	0.046	0.046	0.046	0.046	0.046	0.046	0.046	0.046
<b>2007</b>	0.013	0.026	0.042	0.053	0.059	0.062	0.063	0.063	0.063	0.063	0.063	0.063	0.063	0.063
<b>2008</b>	0.016	0.034	0.053	0.065	0.070	0.071	0.072	0.072	0.072	0.072	0.072	0.072	0.072	0.072
<b>2009</b>	0.010	0.025	0.043	0.051	0.054	0.055	0.055	0.055	0.055	0.055	0.055	0.055	0.055	0.055
<b>2010</b>	0.011	0.023	0.039	0.050	0.055	0.057	0.058	0.058	0.058	0.058	0.058	0.058	0.058	0.058
<b>2011</b>	0.012	0.028	0.049	0.064	0.072	0.075	0.076	0.076	0.076	0.076	0.076	0.076	0.076	0.076
<b>2012</b>	0.012	0.027	0.045	0.060	0.069	0.072	0.073	0.074	0.074	0.074	0.074	0.074	0.074	0.074
<b>2013</b>	0.005	0.018	0.044	0.072	0.086	0.090	0.092	0.092	0.092	0.092	0.092	0.092	0.092	0.092
<b>2014</b>	0.006	0.017	0.040	0.066	0.082	0.089	0.091	0.092	0.092	0.092	0.092	0.092	0.092	0.092
<b>2015</b>	0.004	0.009	0.020	0.038	0.059	0.075	0.083	0.087	0.089	0.090	0.090	0.090	0.090	0.090
<b>2016</b>	0.007	0.015	0.028	0.045	0.063	0.075	0.083	0.086	0.088	0.089	0.089	0.089	0.089	0.089
<b>2017</b>	0.009	0.022	0.043	0.061	0.070	0.074	0.075	0.075	0.075	0.076	0.076	0.076	0.076	0.076

Table 4.14 continued.

Table 4.15. Model estimates of yellowfin sole full selection fishing mortality and exploitation rate (catch/total biomass).

<b>Year</b>	<b>Full selection F</b>	<b>Exploitation Rate</b>
1974	0.15	0.03
1975	0.13	0.04
1976	0.13	0.03
1977	0.06	0.02
1978	0.12	0.05
1979	0.07	0.03
1980	0.08	0.03
1981	0.06	0.03
1982	0.04	0.03
1983	0.05	0.03
1984	0.07	0.05
1985	0.11	0.06
1986	0.10	0.06
1987	0.10	0.06
1988	0.12	0.07
1989	0.09	0.05
1990	0.04	0.03
1991	0.04	0.03
1992	0.07	0.05
1993	0.06	0.03
1994	0.07	0.04
1995	0.06	0.04
1996	0.06	0.04
1997	0.09	0.06
1998	0.05	0.04
1999	0.04	0.03
2000	0.05	0.03
2001	0.04	0.02
2002	0.04	0.03
2003	0.04	0.03
2004	0.04	0.02
2005	0.05	0.03
2006	0.05	0.03
2007	0.06	0.04
2008	0.07	0.05
2009	0.05	0.04
2010	0.06	0.04
2011	0.08	0.05
2012	0.07	0.05
2013	0.09	0.06
2014	0.09	0.06
2015	0.09	0.05
2016	0.09	0.05
2017	0.08	0.05

Table 4.16 Model estimates of yellowfin sole age-specific selectivities for the survey and fishery (ages 4 to 20 from left to right).

Table 4.16 (continued) Model estimates of yellowfin sole age-specific selectivities for the survey and fishery (ages 4 to 20 from left to right).

Table 4.17. Model estimates of yellowfin sole age 2+ total biomass (t) and begin-year female spawning biomass (t) from the 2016 and 2017 stock assessments.

Year	2017 Assessment						2016 Assessment	
	Female spawning biomass	Lower 95% C.I.	Upper 95% C.I.	Total biomass	Lower 95% C.I.	Upper 95% C.I.	Female spawning biomass	Total biomass
1964	142,369	125,963	158,775	926,625	884,358	968,892	17,615	839,742
1965	169,571	150,673	188,469	901,174	859,348	943,000	36,707	835,631
1966	207,104	181,164	233,044	936,828	894,264	979,392	61,107	879,970
1967	218,877	183,738	254,016	911,246	868,430	954,062	75,435	863,836
1968	213,928	172,597	255,259	823,838	782,109	865,567	74,113	782,492
1969	197,480	159,473	235,487	850,296	806,122	894,470	69,945	810,677
1970	135,238	107,904	162,572	813,647	767,283	860,011	65,929	786,026
1971	87,750	68,217	107,284	862,710	810,862	914,558	59,749	846,563
1972	71,363	52,436	90,290	936,838	876,587	997,089	41,419	909,321
1973	78,284	55,940	100,627	1,188,580	1,117,114	1,260,046	48,795	1,155,700
1974	84,991	62,369	107,612	1,433,880	1,350,985	1,516,775	64,968	1,397,720
1975	134,520	104,490	164,550	1,786,070	1,687,158	1,884,982	114,443	1,739,260
1976	194,457	159,757	229,157	2,091,310	1,979,840	2,202,780	175,884	2,035,950
1977	285,756	244,715	326,797	2,400,420	2,276,574	2,524,266	267,530	2,336,020
1978	400,220	353,682	446,758	2,694,380	2,558,675	2,830,085	381,170	2,620,880
1979	515,975	465,098	566,852	2,853,520	2,707,666	2,999,374	494,992	2,771,670
1980	644,788	589,440	700,136	3,034,890	2,879,623	3,190,157	620,393	2,944,900
1981	763,587	704,136	823,038	3,200,280	3,035,796	3,364,764	735,618	3,103,120
1982	830,159	769,672	890,646	3,312,110	3,142,696	3,481,524	799,728	3,208,630
1983	930,442	866,389	994,495	3,289,120	3,117,812	3,460,428	895,921	3,183,300
1984	1,009,100	942,343	1,075,857	3,509,840	3,324,897	3,694,783	971,246	3,393,380
1985	1,054,930	984,923	1,124,937	3,516,680	3,324,359	3,709,001	1,013,620	3,394,790
1986	1,040,470	969,028	1,111,912	3,228,140	3,041,303	3,414,977	996,871	3,108,320
1987	1,032,280	957,877	1,106,683	3,183,200	2,990,263	3,376,137	986,120	3,058,180
1988	973,826	899,973	1,047,679	3,082,600	2,889,542	3,275,658	927,181	2,955,620
1989	944,673	869,393	1,019,953	3,130,950	2,926,591	3,335,309	896,354	2,994,610
1990	953,446	877,432	1,029,460	2,993,920	2,791,865	3,195,975	903,497	2,858,500
1991	1,030,640	950,675	1,110,605	3,106,100	2,897,247	3,314,953	977,265	2,964,280
1992	1,108,300	1,022,799	1,193,801	3,303,480	3,081,401	3,525,559	1,051,080	3,150,890
1993	1,139,450	1,049,929	1,228,971	3,328,080	3,099,130	3,557,030	1,079,390	3,167,600
1994	1,139,730	1,049,752	1,229,708	3,362,290	3,130,320	3,594,260	1,078,550	3,194,550
1995	1,135,560	1,044,219	1,226,901	3,135,510	2,911,867	3,359,153	1,072,910	2,972,780
1996	1,068,660	980,760	1,156,560	3,047,450	2,827,286	3,267,614	1,007,480	2,881,690
1997	1,029,840	942,754	1,116,926	3,056,250	2,832,945	3,279,555	968,198	2,881,350
1998	964,342	879,785	1,048,899	2,782,490	2,569,490	2,995,490	903,440	2,612,680
1999	953,817	869,562	1,038,072	2,599,520	2,395,766	2,803,274	891,856	2,434,810
2000	939,002	855,138	1,022,866	2,645,900	2,441,086	2,850,714	876,306	2,472,620
2001	932,963	849,579	1,016,347	2,571,130	2,369,891	2,772,369	868,908	2,397,250
2002	929,799	846,867	1,012,731	2,612,290	2,410,173	2,814,407	864,250	2,430,820
2003	937,347	854,321	1,020,373	2,820,560	2,606,587	3,034,533	869,246	2,618,170
2004	966,374	881,636	1,051,112	3,028,200	2,801,621	3,254,779	893,962	2,805,970
2005	981,900	896,204	1,067,596	3,138,190	2,904,329	3,372,051	905,612	2,906,560
2006	1,001,710	914,164	1,089,256	3,121,280	2,886,123	3,356,437	920,437	2,887,410
2007	1,007,020	918,540	1,095,500	3,131,800	2,892,122	3,371,478	921,166	2,895,700
2008	984,018	895,586	1,072,450	3,002,470	2,766,978	3,237,962	894,603	2,771,050
2009	950,705	863,010	1,038,400	2,834,320	2,603,218	3,065,422	858,763	2,606,720
2010	928,940	841,575	1,016,305	2,903,440	2,665,625	3,141,255	835,187	2,670,900
2011	910,298	823,033	997,563	2,936,040	2,690,395	3,181,685	815,498	2,701,250
2012	897,681	809,011	986,351	2,927,810	2,676,751	3,178,869	802,230	2,693,730
2013	894,342	802,405	986,279	2,883,080	2,627,632	3,138,528	799,268	2,647,970
2014	858,579	764,911	952,247	2,680,370	2,430,474	2,930,266	764,979	2,449,450
2015	861,299	763,454	959,144	2,699,440	2,438,694	2,960,186	767,803	2,451,240
2016	876,687	775,173	978,201	2,852,570	2,552,951	3,152,189	775,148	2,457,260
2017	884,750	780,860	988,640	2,878,150	2,535,096	3,221,204		

Table 4.18—Model estimates of yellowfin sole population numbers at age (billions) for 1954-2017.

Female s		1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19
<b>1954</b>		1.04	0.70	0.30	0.22	0.22	0.20	0.20	0.20	0.20	0.20	0.20	0.20	0.20	0.20	0.20	0.20	0.20	0.20	
<b>1955</b>		0.84	0.92	0.62	0.27	0.20	0.19	0.18	0.18	0.18	0.18	0.18	0.18	0.17	0.17	0.17	0.17	0.17	0.17	
<b>1956</b>		0.73	0.75	0.81	0.55	0.24	0.18	0.17	0.16	0.16	0.16	0.16	0.16	0.15	0.15	0.15	0.15	0.15	0.15	
<b>1957</b>		2.95	0.65	0.66	0.72	0.49	0.21	0.16	0.15	0.14	0.14	0.14	0.14	0.13	0.13	0.13	0.13	0.13	0.13	
<b>1958</b>		2.33	2.62	0.58	0.59	0.64	0.43	0.19	0.14	0.13	0.12	0.12	0.12	0.12	0.12	0.12	0.12	0.12	0.12	
<b>1959</b>		1.49	2.07	2.32	0.51	0.52	0.57	0.38	0.16	0.12	0.12	0.11	0.11	0.10	0.10	0.10	0.10	0.10	0.10	
<b>1960</b>		1.00	1.32	1.83	2.06	0.45	0.46	0.50	0.33	0.14	0.10	0.09	0.08	0.08	0.08	0.08	0.08	0.08	0.08	
<b>1961</b>		0.51	0.89	1.17	1.62	1.82	0.40	0.41	0.44	0.29	0.11	0.07	0.05	0.04	0.04	0.04	0.04	0.04	0.04	
<b>1962</b>		0.95	0.46	0.79	1.04	1.44	1.62	0.36	0.36	0.39	0.24	0.09	0.04	0.02	0.01	0.01	0.01	0.01	0.01	
<b>1963</b>		0.46	0.85	0.40	0.70	0.92	1.28	1.44	0.32	0.32	0.34	0.21	0.07	0.02	0.00	0.00	0.00	0.00	0.00	
<b>1964</b>		0.42	0.41	0.75	0.36	0.62	0.81	1.09	1.16	0.23	0.22	0.22	0.13	0.04	0.02	0.00	0.00	0.00	0.00	
<b>1965</b>		0.59	0.37	0.37	0.67	0.32	0.54	0.70	0.89	0.86	0.16	0.14	0.14	0.09	0.03	0.01	0.00	0.00	0.00	
<b>1966</b>		0.61	0.52	0.33	0.32	0.59	0.28	0.48	0.61	0.76	0.68	0.12	0.10	0.10	0.06	0.02	0.01	0.00	0.00	
<b>1967</b>		1.25	0.54	0.46	0.29	0.29	0.52	0.25	0.42	0.51	0.59	0.46	0.07	0.06	0.06	0.03	0.01	0.00	0.00	
<b>1968</b>		1.91	1.11	0.48	0.41	0.26	0.25	0.46	0.21	0.34	0.38	0.36	0.25	0.04	0.03	0.03	0.02	0.01	0.00	
<b>1969</b>		1.96	1.70	0.99	0.42	0.36	0.23	0.22	0.40	0.19	0.29	0.29	0.26	0.16	0.02	0.02	0.02	0.01	0.00	
<b>1970</b>		2.58	1.74	1.50	0.88	0.38	0.32	0.20	0.19	0.29	0.11	0.14	0.14	0.12	0.08	0.01	0.01	0.01	0.00	
<b>1971</b>		2.87	2.29	1.54	1.33	0.78	0.33	0.27	0.13	0.09	0.13	0.05	0.07	0.06	0.06	0.03	0.00	0.00	0.00	
<b>1972</b>		2.25	2.54	2.03	1.37	1.18	0.68	0.28	0.16	0.06	0.04	0.07	0.02	0.03	0.03	0.03	0.02	0.00	0.00	
<b>1973</b>		1.56	2.00	2.26	1.80	1.21	1.05	0.61	0.24	0.14	0.06	0.04	0.06	0.02	0.03	0.03	0.02	0.01	0.00	
<b>1974</b>		2.10	1.38	1.77	2.00	1.60	1.07	0.93	0.53	0.20	0.11	0.04	0.02	0.03	0.01	0.02	0.02	0.01	0.00	
<b>1975</b>		2.47	1.86	1.23	1.57	1.78	1.42	0.95	0.82	0.46	0.17	0.09	0.03	0.02	0.03	0.01	0.01	0.01	0.01	
<b>1976</b>		1.62	2.19	1.65	1.09	1.39	1.57	1.25	0.83	0.69	0.37	0.14	0.07	0.02	0.01	0.02	0.01	0.01	0.01	
<b>1977</b>		2.04	1.44	1.94	1.46	0.96	1.23	1.39	1.10	0.71	0.57	0.30	0.11	0.05	0.02	0.01	0.02	0.01	0.01	
<b>1978</b>		1.34	1.81	1.28	1.72	1.29	0.85	1.08	1.20	0.94	0.60	0.48	0.25	0.09	0.04	0.01	0.01	0.00	0.01	
<b>1979</b>		0.85	1.18	1.61	1.13	1.52	1.14	0.74	0.93	1.00	0.76	0.48	0.38	0.20	0.07	0.04	0.01	0.01	0.01	
<b>1980</b>		1.64	0.76	1.05	1.42	1.00	1.34	1.00	0.65	0.79	0.84	0.63	0.40	0.32	0.17	0.06	0.03	0.01	0.01	

Table 4.18—Model estimates of yellowfin sole population numbers at age (billions) for 1954-2017 (continued).

	<b>1</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>5</b>	<b>6</b>	<b>7</b>	<b>8</b>	<b>9</b>	<b>10</b>	<b>11</b>	<b>12</b>	<b>13</b>	<b>14</b>	<b>15</b>	<b>16</b>	<b>17</b>	<b>18</b>	<b>19</b>	<b>20</b>
<b>1981</b>	1.21	1.45	0.67	0.93	1.26	0.89	1.18	0.88	0.56	0.68	0.71	0.53	0.33	0.26	0.14	0.05	0.02	0.01	0.00	0.02
<b>1982</b>	3.51	1.08	1.29	0.59	0.82	1.11	0.78	1.04	0.76	0.48	0.57	0.60	0.44	0.28	0.22	0.11	0.04	0.02	0.01	0.02
<b>1983</b>	0.65	3.11	0.95	1.14	0.53	0.73	0.98	0.68	0.89	0.65	0.41	0.49	0.51	0.38	0.24	0.19	0.10	0.03	0.02	0.02
<b>1984</b>	2.89	0.58	2.76	0.85	1.01	0.46	0.64	0.85	0.58	0.76	0.55	0.35	0.41	0.43	0.32	0.20	0.16	0.08	0.03	0.04
<b>1985</b>	1.00	2.56	0.51	2.45	0.75	0.89	0.41	0.56	0.73	0.49	0.63	0.46	0.29	0.34	0.35	0.26	0.16	0.13	0.07	0.05
<b>1986</b>	0.76	0.88	2.27	0.45	2.17	0.66	0.79	0.36	0.48	0.61	0.40	0.51	0.37	0.23	0.27	0.28	0.21	0.13	0.10	0.10
<b>1987</b>	1.04	0.68	0.78	2.02	0.40	1.92	0.58	0.68	0.30	0.39	0.49	0.32	0.41	0.29	0.18	0.22	0.23	0.17	0.11	0.16
<b>1988</b>	1.42	0.92	0.60	0.70	1.79	0.36	1.70	0.51	0.59	0.25	0.32	0.40	0.26	0.33	0.24	0.15	0.18	0.18	0.14	0.21
<b>1989</b>	1.42	1.26	0.82	0.53	0.62	1.58	0.31	1.49	0.44	0.49	0.20	0.25	0.31	0.20	0.26	0.19	0.12	0.14	0.14	0.27
<b>1990</b>	0.71	1.26	1.12	0.73	0.47	0.55	1.40	0.28	1.30	0.38	0.41	0.17	0.21	0.25	0.17	0.21	0.15	0.09	0.11	0.34
<b>1991</b>	0.79	0.63	1.12	0.99	0.64	0.42	0.48	1.24	0.24	1.13	0.32	0.35	0.14	0.18	0.22	0.14	0.18	0.13	0.08	0.38
<b>1992</b>	1.76	0.70	0.56	0.99	0.88	0.57	0.37	0.43	1.08	0.21	0.96	0.28	0.30	0.12	0.15	0.18	0.12	0.15	0.11	0.39
<b>1993</b>	1.05	1.56	0.62	0.49	0.88	0.78	0.50	0.33	0.37	0.92	0.18	0.80	0.23	0.24	0.10	0.12	0.15	0.10	0.13	0.41
<b>1994</b>	0.89	0.93	1.38	0.55	0.44	0.78	0.69	0.44	0.28	0.32	0.79	0.15	0.68	0.19	0.20	0.08	0.10	0.13	0.08	0.45
<b>1995</b>	0.90	0.79	0.83	1.22	0.49	0.39	0.68	0.60	0.38	0.24	0.27	0.66	0.12	0.56	0.16	0.17	0.07	0.09	0.11	0.44
<b>1996</b>	2.24	0.80	0.70	0.73	1.09	0.43	0.34	0.60	0.52	0.32	0.20	0.22	0.55	0.10	0.47	0.13	0.14	0.06	0.07	0.46
<b>1997</b>	0.97	1.98	0.71	0.62	0.65	0.96	0.38	0.30	0.52	0.44	0.27	0.17	0.19	0.46	0.09	0.39	0.11	0.12	0.05	0.44
<b>1998</b>	0.80	0.86	1.76	0.63	0.55	0.57	0.84	0.33	0.25	0.43	0.36	0.22	0.14	0.15	0.37	0.07	0.32	0.09	0.10	0.40
<b>1999</b>	1.00	0.71	0.76	1.56	0.56	0.49	0.50	0.73	0.28	0.22	0.37	0.31	0.19	0.12	0.13	0.31	0.06	0.27	0.08	0.41
<b>2000</b>	1.41	0.89	0.63	0.68	1.38	0.49	0.43	0.45	0.65	0.25	0.19	0.31	0.26	0.16	0.10	0.11	0.27	0.05	0.23	0.42
<b>2001</b>	0.90	1.25	0.79	0.56	0.60	1.22	0.44	0.38	0.39	0.56	0.21	0.16	0.26	0.22	0.13	0.08	0.09	0.22	0.04	0.54
<b>2002</b>	1.25	0.80	1.11	0.70	0.50	0.53	1.08	0.39	0.33	0.34	0.48	0.18	0.14	0.23	0.19	0.11	0.07	0.08	0.19	0.50
<b>2003</b>	1.21	1.11	0.71	0.98	0.62	0.44	0.47	0.96	0.34	0.29	0.29	0.41	0.16	0.12	0.19	0.16	0.10	0.06	0.07	0.59
<b>2004</b>	1.98	1.08	0.98	0.63	0.87	0.55	0.39	0.42	0.84	0.29	0.25	0.25	0.35	0.13	0.10	0.16	0.14	0.08	0.05	0.56
<b>2005</b>	0.91	1.76	0.96	0.87	0.56	0.77	0.49	0.34	0.36	0.73	0.25	0.21	0.21	0.30	0.11	0.08	0.14	0.12	0.07	0.52
<b>2006</b>	1.12	0.80	1.56	0.85	0.77	0.49	0.68	0.43	0.30	0.31	0.62	0.22	0.18	0.18	0.25	0.10	0.07	0.12	0.10	0.50
<b>2007</b>	1.44	1.00	0.71	1.38	0.75	0.68	0.43	0.58	0.36	0.25	0.27	0.53	0.18	0.15	0.15	0.22	0.08	0.06	0.10	0.51
<b>2008</b>	1.33	1.28	0.88	0.63	1.22	0.66	0.60	0.38	0.50	0.31	0.21	0.22	0.44	0.15	0.13	0.13	0.18	0.07	0.05	0.50
<b>2009</b>	1.47	1.18	1.13	0.78	0.56	1.08	0.59	0.53	0.32	0.42	0.26	0.18	0.18	0.36	0.13	0.11	0.11	0.15	0.06	0.46
<b>2010</b>	1.73	1.30	1.05	1.00	0.70	0.50	0.96	0.51	0.45	0.27	0.36	0.22	0.15	0.15	0.30	0.11	0.09	0.09	0.12	0.43
<b>2011</b>	0.62	1.53	1.16	0.93	0.89	0.62	0.44	0.84	0.45	0.39	0.23	0.30	0.18	0.12	0.13	0.25	0.09	0.07	0.07	0.46
<b>2012</b>	0.47	0.55	1.36	1.03	0.82	0.79	0.54	0.38	0.72	0.38	0.32	0.19	0.25	0.15	0.10	0.11	0.21	0.07	0.06	0.44
<b>2013</b>	1.05	0.42	0.48	1.20	0.91	0.73	0.70	0.48	0.33	0.61	0.31	0.27	0.16	0.20	0.12	0.08	0.09	0.17	0.06	0.41
<b>2014</b>	1.31	0.93	0.37	0.43	1.07	0.81	0.64	0.61	0.42	0.28	0.51	0.26	0.22	0.13	0.16	0.10	0.07	0.07	0.14	0.38
<b>2015</b>	1.13	1.16	0.83	0.33	0.38	0.95	0.71	0.57	0.53	0.35	0.23	0.41	0.21	0.18	0.10	0.13	0.08	0.05	0.06	0.42
<b>2016</b>	1.20	1.00	1.03	0.73	0.29	0.34	0.84	0.63	0.50	0.46	0.30	0.20	0.34	0.17	0.14	0.08	0.11	0.06	0.04	0.39
<b>2017</b>	1.21	1.06	0.89	0.91	0.65	0.26	0.30	0.74	0.55	0.43	0.39	0.25	0.16	0.28	0.14	0.12	0.07	0.09	0.05	0.35

Table 4.18. Model estimates of yellowfin sole population numbers at age (billions) for 1954-2017  
 (continued).

	<b>1</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>5</b>	<b>6</b>	<b>7</b>	<b>8</b>	<b>9</b>	<b>10</b>	<b>11</b>	<b>12</b>	<b>13</b>	<b>14</b>	<b>15</b>	<b>16</b>	<b>17</b>	<b>18</b>	<b>19</b>	<b>20</b>
<b>1954</b>	1.04	0.37	0.25	0.21	0.21	0.20	0.20	0.20	0.20	0.20	0.20	0.20	0.20	0.20	0.20	0.20	0.20	0.20	0.20	0.20
<b>1955</b>	0.84	0.92	0.33	0.22	0.19	0.19	0.18	0.18	0.18	0.18	0.18	0.18	0.17	0.17	0.17	0.17	0.17	0.17	0.17	0.35
<b>1956</b>	0.73	0.75	0.81	0.29	0.20	0.17	0.17	0.16	0.16	0.16	0.16	0.16	0.15	0.15	0.15	0.15	0.15	0.15	0.15	0.46
<b>1957</b>	2.95	0.65	0.66	0.72	0.26	0.17	0.15	0.15	0.14	0.14	0.14	0.14	0.13	0.13	0.13	0.13	0.13	0.13	0.13	0.53
<b>1958</b>	2.33	2.62	0.58	0.59	0.64	0.23	0.15	0.13	0.13	0.12	0.12	0.12	0.12	0.12	0.12	0.12	0.12	0.12	0.12	0.58
<b>1959</b>	1.49	2.07	2.32	0.51	0.52	0.57	0.20	0.14	0.12	0.11	0.11	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.60
<b>1960</b>	1.00	1.32	1.83	2.06	0.45	0.46	0.50	0.18	0.11	0.09	0.09	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.53
<b>1961</b>	0.51	0.89	1.17	1.62	1.82	0.40	0.39	0.39	0.12	0.06	0.05	0.04	0.04	0.04	0.04	0.04	0.04	0.04	0.04	0.30
<b>1962</b>	0.95	0.46	0.79	1.03	1.41	1.51	0.28	0.19	0.11	0.02	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.05
<b>1963</b>	0.46	0.83	0.38	0.50	0.23	0.02	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>1964</b>	0.42	0.41	0.74	0.34	0.45	0.20	0.02	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>1965</b>	0.59	0.38	0.37	0.65	0.24	0.28	0.13	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>1966</b>	0.61	0.52	0.33	0.32	0.58	0.21	0.25	0.11	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>1967</b>	1.25	0.54	0.46	0.29	0.29	0.51	0.19	0.22	0.09	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>1968</b>	1.91	1.11	0.48	0.41	0.26	0.25	0.44	0.14	0.13	0.05	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>1969</b>	1.96	1.70	0.99	0.42	0.36	0.23	0.22	0.38	0.11	0.08	0.03	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>1970</b>	2.58	1.74	1.50	0.88	0.38	0.32	0.21	0.20	0.32	0.08	0.05	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>1971</b>	2.87	2.29	1.54	1.33	0.78	0.33	0.28	0.18	0.17	0.28	0.07	0.04	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>1972</b>	2.25	2.54	2.03	1.37	1.18	0.69	0.28	0.17	0.09	0.08	0.14	0.03	0.02	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>1973</b>	1.56	2.00	2.26	1.80	1.21	1.05	0.60	0.22	0.11	0.05	0.05	0.07	0.02	0.01	0.00	0.00	0.00	0.00	0.00	0.00
<b>1974</b>	2.10	1.38	1.77	2.00	1.60	1.07	0.92	0.47	0.14	0.06	0.03	0.03	0.04	0.01	0.01	0.00	0.00	0.00	0.00	0.00
<b>1975</b>	2.47	1.86	1.23	1.57	1.77	1.40	0.92	0.75	0.37	0.11	0.05	0.02	0.02	0.03	0.01	0.00	0.00	0.00	0.00	0.00
<b>1976</b>	1.62	2.19	1.65	1.09	1.39	1.57	1.23	0.78	0.60	0.29	0.08	0.04	0.02	0.02	0.03	0.01	0.00	0.00	0.00	0.00
<b>1977</b>	2.04	1.44	1.94	1.46	0.97	1.23	1.39	1.08	0.67	0.50	0.23	0.07	0.03	0.01	0.01	0.02	0.00	0.00	0.00	0.00
<b>1978</b>	1.34	1.81	1.28	1.72	1.30	0.86	1.09	1.22	0.94	0.57	0.42	0.19	0.06	0.03	0.01	0.01	0.02	0.00	0.00	0.00
<b>1979</b>	0.85	1.18	1.61	1.13	1.52	1.15	0.75	0.94	1.03	0.77	0.46	0.33	0.15	0.04	0.02	0.01	0.01	0.01	0.00	0.00
<b>1980</b>	1.64	0.76	1.05	1.42	1.00	1.35	1.01	0.65	0.81	0.86	0.64	0.38	0.28	0.13	0.04	0.02	0.01	0.01	0.01	0.00
<b>1981</b>	1.21	1.45	0.67	0.93	1.26	0.89	1.19	0.89	0.57	0.70	0.75	0.55	0.32	0.23	0.11	0.03	0.01	0.01	0.01	0.01
<b>1982</b>	3.51	1.08	1.29	0.59	0.82	1.12	0.78	1.04	0.77	0.49	0.60	0.63	0.46	0.27	0.19	0.09	0.03	0.01	0.01	0.01
<b>1983</b>	0.65	3.11	0.95	1.14	0.53	0.73	0.98	0.68	0.90	0.66	0.42	0.51	0.54	0.39	0.23	0.16	0.08	0.02	0.01	0.02
<b>1984</b>	2.89	0.58	2.76	0.85	1.01	0.46	0.64	0.85	0.58	0.76	0.56	0.36	0.43	0.45	0.33	0.19	0.14	0.06	0.02	0.02
<b>1985</b>	1.00	2.56	0.51	2.45	0.75	0.89	0.41	0.55	0.72	0.48	0.63	0.46	0.29	0.36	0.38	0.27	0.16	0.11	0.05	0.03
<b>1986</b>	0.76	0.88	2.27	0.45	2.17	0.66	0.78	0.35	0.46	0.58	0.39	0.50	0.37	0.23	0.28	0.30	0.22	0.13	0.09	0.07
<b>1987</b>	1.04	0.68	0.78	2.02	0.40	1.92	0.58	0.67	0.29	0.37	0.47	0.31	0.41	0.30	0.19	0.23	0.24	0.17	0.10	0.13
<b>1988</b>	1.42	0.92	0.60	0.70	1.79	0.36	1.70	0.51	0.57	0.24	0.30	0.38	0.25	0.33	0.24	0.15	0.18	0.19	0.14	0.19
<b>1989</b>	1.42	1.26	0.82	0.53	0.62	1.59	0.31	1.48	0.43	0.46	0.19	0.23	0.29	0.20	0.26	0.19	0.12	0.14	0.15	0.26
<b>1990</b>	0.71	1.26	1.12	0.73	0.47	0.55	1.41	0.28	1.29	0.36	0.37	0.15	0.19	0.24	0.16	0.21	0.15	0.10	0.12	0.33
<b>1991</b>	0.79	0.63	1.12	0.99	0.64	0.42	0.48	1.24	0.24	1.11	0.31	0.32	0.13	0.16	0.20	0.14	0.18	0.13	0.08	0.38
<b>1992</b>	1.76	0.70	0.56	0.99	0.88	0.57	0.37	0.43	1.07	0.21	0.94	0.26	0.27	0.11	0.14	0.17	0.12	0.15	0.11	0.39
<b>1993</b>	1.05	1.56	0.62	0.49	0.88	0.78	0.50	0.32	0.36	0.89	0.17	0.78	0.22	0.22	0.09	0.11	0.14	0.09	0.12	0.41
<b>1994</b>	0.89	0.93	1.38	0.55	0.44	0.78	0.69	0.44	0.28	0.31	0.77	0.14	0.66	0.18	0.19	0.08	0.09	0.12	0.08	0.45
<b>1995</b>	0.90	0.79	0.83	1.22	0.49	0.39	0.68	0.60	0.38	0.24	0.26	0.64	0.12	0.55	0.15	0.16	0.06	0.08	0.10	0.44
<b>1996</b>	2.24	0.80	0.70	0.73	1.09	0.43	0.34	0.60	0.51	0.32	0.20	0.22	0.53	0.10	0.46	0.13	0.13	0.05	0.07	0.45
<b>1997</b>	0.97	1.98	0.71	0.62	0.65	0.96	0.38	0.29	0.51	0.43	0.27	0.17	0.18	0.44	0.08	0.38	0.11	0.11	0.04	0.43

Table 4.18. Model estimates of yellowfin sole population numbers at age (billions) for 1954-2017  
 (continued).

	<b>1</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>5</b>	<b>6</b>	<b>7</b>	<b>8</b>	<b>9</b>	<b>10</b>	<b>11</b>	<b>12</b>	<b>13</b>	<b>14</b>	<b>15</b>	<b>16</b>	<b>17</b>	<b>18</b>	<b>19</b>	<b>20</b>
<b>1998</b>	0.80	0.86	1.76	0.63	0.55	0.57	0.84	0.33	0.25	0.42	0.35	0.22	0.13	0.15	0.36	0.07	0.31	0.09	0.09	0.38
<b>1999</b>	1.00	0.71	0.76	1.56	0.56	0.49	0.51	0.74	0.28	0.21	0.35	0.30	0.18	0.11	0.12	0.30	0.06	0.26	0.07	0.40
<b>2000</b>	1.41	0.89	0.63	0.68	1.38	0.49	0.43	0.45	0.65	0.25	0.19	0.31	0.25	0.16	0.10	0.11	0.26	0.05	0.22	0.40
<b>2001</b>	0.90	1.25	0.79	0.56	0.60	1.23	0.44	0.38	0.40	0.57	0.21	0.16	0.26	0.21	0.13	0.08	0.09	0.22	0.04	0.52
<b>2002</b>	1.25	0.80	1.11	0.70	0.50	0.53	1.09	0.39	0.34	0.35	0.50	0.19	0.14	0.22	0.18	0.11	0.07	0.08	0.19	0.48
<b>2003</b>	1.21	1.11	0.71	0.98	0.62	0.44	0.47	0.96	0.34	0.29	0.30	0.42	0.16	0.12	0.19	0.16	0.10	0.06	0.06	0.57
<b>2004</b>	1.98	1.08	0.98	0.63	0.87	0.55	0.39	0.42	0.84	0.29	0.25	0.25	0.36	0.13	0.10	0.16	0.13	0.08	0.05	0.54
<b>2005</b>	0.91	1.76	0.96	0.87	0.56	0.77	0.49	0.34	0.36	0.73	0.25	0.21	0.22	0.31	0.12	0.08	0.14	0.11	0.07	0.50
<b>2006</b>	1.12	0.80	1.56	0.85	0.77	0.49	0.68	0.43	0.30	0.31	0.62	0.21	0.18	0.18	0.26	0.10	0.07	0.12	0.10	0.48
<b>2007</b>	1.44	1.00	0.71	1.38	0.75	0.68	0.43	0.59	0.36	0.25	0.26	0.52	0.18	0.15	0.16	0.22	0.08	0.06	0.10	0.49
<b>2008</b>	1.33	1.28	0.88	0.63	1.22	0.67	0.60	0.38	0.50	0.30	0.21	0.22	0.44	0.15	0.13	0.13	0.18	0.07	0.05	0.49
<b>2009</b>	1.47	1.18	1.13	0.78	0.56	1.08	0.59	0.52	0.32	0.42	0.25	0.17	0.18	0.36	0.12	0.11	0.11	0.15	0.06	0.45
<b>2010</b>	1.73	1.30	1.05	1.00	0.70	0.50	0.96	0.52	0.46	0.28	0.35	0.21	0.15	0.15	0.30	0.10	0.09	0.09	0.13	0.42
<b>2011</b>	0.62	1.53	1.16	0.93	0.89	0.62	0.44	0.84	0.44	0.39	0.23	0.30	0.18	0.12	0.13	0.25	0.09	0.07	0.07	0.46
<b>2012</b>	0.47	0.55	1.36	1.03	0.82	0.79	0.54	0.38	0.72	0.37	0.32	0.19	0.24	0.15	0.10	0.10	0.21	0.07	0.06	0.44
<b>2013</b>	1.05	0.42	0.48	1.20	0.91	0.73	0.69	0.47	0.33	0.60	0.31	0.26	0.16	0.20	0.12	0.08	0.09	0.17	0.06	0.41
<b>2014</b>	1.31	0.93	0.37	0.43	1.07	0.81	0.64	0.61	0.41	0.28	0.49	0.25	0.21	0.13	0.16	0.10	0.07	0.07	0.14	0.38
<b>2015</b>	1.13	1.16	0.83	0.33	0.38	0.95	0.71	0.57	0.53	0.35	0.23	0.40	0.20	0.17	0.10	0.13	0.08	0.05	0.06	0.42
<b>2016</b>	1.20	1.00	1.03	0.73	0.29	0.34	0.84	0.63	0.50	0.46	0.30	0.19	0.33	0.17	0.14	0.08	0.11	0.06	0.04	0.39
<b>2017</b>	1.21	1.06	0.89	0.91	0.65	0.26	0.30	0.74	0.55	0.43	0.39	0.25	0.16	0.27	0.13	0.11	0.07	0.09	0.05	0.35

Table 4.19. Model estimates of the number of female spawners (millions) 1964-2017.

	<b>6</b>	<b>7</b>	<b>8</b>	<b>9</b>	<b>10</b>	<b>11</b>	<b>12</b>	<b>13</b>	<b>14</b>	<b>15</b>	<b>16</b>	<b>17</b>	<b>18</b>	<b>19</b>	<b>20</b>
<b>1964</b>	8.9	41.8	112.8	49.8	89.4	141.7	110.9	41.0	14.9	2.5	0.1	0.0	0.0	0.0	0.0
<b>1965</b>	6.0	26.6	86.8	184.1	65.4	90.5	115.4	78.4	27.3	9.6	1.6	0.1	0.0	0.0	0.0
<b>1966</b>	3.1	18.3	59.2	162.1	282.3	75.0	81.8	90.0	57.6	19.4	6.8	1.1	0.0	0.0	0.0
<b>1967</b>	5.7	9.5	40.6	109.5	242.3	298.8	58.4	52.8	53.9	33.2	11.1	3.8	0.6	0.0	0.0
<b>1968</b>	2.8	17.5	20.8	73.3	155.0	233.8	206.0	32.9	27.4	26.9	16.4	5.4	1.9	0.3	0.0
<b>1969</b>	2.5	8.6	39.1	39.5	118.3	188.8	210.8	146.7	20.7	16.1	15.4	9.2	3.0	1.1	0.2
<b>1970</b>	3.5	7.7	18.2	62.1	43.6	92.0	115.4	111.5	73.1	10.0	7.7	7.3	4.4	1.4	0.6
<b>1971</b>	3.6	10.2	12.3	19.4	55.7	31.4	54.1	59.3	54.1	34.3	4.6	3.6	3.4	2.0	0.9
<b>1972</b>	7.5	10.5	15.7	13.4	18.2	42.1	19.4	29.2	30.2	26.6	16.8	2.2	1.7	1.6	1.4
<b>1973</b>	11.5	23.2	23.7	30.7	23.0	25.2	47.7	19.2	27.2	27.0	23.1	13.4	1.6	1.2	2.1
<b>1974</b>	11.8	35.4	51.3	43.4	44.1	23.2	19.2	31.0	11.7	16.0	15.8	13.4	7.8	0.9	1.9
<b>1975</b>	15.6	36.3	79.3	98.8	71.8	56.5	23.4	16.6	25.1	9.1	12.4	12.1	10.3	6.0	2.2
<b>1976</b>	17.3	47.8	80.3	146.4	153.4	88.0	56.2	20.3	13.6	19.9	7.2	9.7	9.4	8.0	6.3
<b>1977</b>	13.6	53.1	106.3	152.3	237.3	194.2	89.2	49.3	16.8	10.8	15.7	5.6	7.6	7.4	11.2
<b>1978</b>	9.3	41.2	116.5	200.1	250.0	312.2	208.1	83.2	43.4	14.3	9.2	13.2	4.7	6.3	15.5
<b>1979</b>	12.6	28.4	89.8	213.5	313.8	311.3	316.0	183.4	69.2	34.9	11.4	7.3	10.4	3.7	17.3
<b>1980</b>	14.8	38.4	62.7	168.9	347.9	408.9	330.4	292.3	160.1	58.5	29.3	9.5	6.0	8.6	17.4
<b>1981</b>	9.7	45.2	85.2	119.4	279.5	458.0	435.4	304.9	253.6	134.2	48.6	24.1	7.8	5.0	21.4
<b>1982</b>	12.3	29.8	100.4	162.2	197.9	369.7	492.1	406.7	268.4	216.0	113.4	40.7	20.2	6.5	22.0
<b>1983</b>	8.0	37.5	66.1	190.9	269.0	263.3	401.2	465.5	363.1	232.0	185.2	96.4	34.6	17.1	24.2
<b>1984</b>	5.1	24.5	82.6	124.6	314.6	356.6	285.0	378.9	415.0	313.4	198.6	157.3	81.8	29.3	35.0
<b>1985</b>	9.8	15.6	54.2	155.8	203.2	409.2	377.3	262.7	329.5	349.2	261.6	164.5	130.1	67.6	53.1
<b>1986</b>	7.3	30.1	34.5	101.5	250.3	257.9	420.0	336.3	220.7	267.8	281.5	209.2	131.4	103.8	96.3
<b>1987</b>	21.1	22.3	66.0	63.6	160.2	315.6	265.2	376.6	284.6	180.8	217.6	227.0	168.5	105.7	160.9
<b>1988</b>	3.9	64.8	49.8	126.1	104.7	208.0	329.3	239.4	319.8	233.7	147.2	175.7	183.1	135.7	214.7

Table 4.19—Model estimates of the number of female spawners (millions) 1964-2017 (CONTINUED)

year/age	<b>6</b>	<b>7</b>	<b>8</b>	<b>9</b>	<b>10</b>	<b>11</b>	<b>12</b>	<b>13</b>	<b>14</b>	<b>15</b>	<b>16</b>	<b>17</b>	<b>18</b>	<b>19</b>	<b>20</b>
<b>1989</b>	17.4	12.0	144.2	94.1	202.5	131.3	210.0	288.6	197.8	255.7	185.3	115.8	138.1	143.7	275.0
<b>1990</b>	6.0	53.6	26.9	277.7	156.1	263.2	137.1	189.9	245.9	163.1	209.1	150.4	93.8	111.8	338.8
<b>1991</b>	4.6	18.5	120.3	51.9	465.6	209.1	287.0	130.3	170.4	213.5	140.4	178.7	128.3	80.0	384.0
<b>1992</b>	6.3	14.2	41.4	231.5	87.0	623.1	227.4	271.7	116.3	147.2	183.0	119.4	151.7	108.8	393.4
<b>1993</b>	8.6	19.3	31.6	78.9	380.8	113.6	659.7	209.3	235.8	97.6	122.6	151.2	98.5	125.0	413.8
<b>1994</b>	8.5	26.3	43.0	60.8	132.5	511.0	123.5	622.0	185.6	201.9	82.9	103.2	127.1	82.7	452.2
<b>1995</b>	4.3	26.2	58.4	81.4	99.4	172.9	543.0	114.4	543.5	156.9	169.5	69.0	85.8	105.5	444.1
<b>1996</b>	4.8	13.0	58.0	110.3	133.5	130.5	185.1	506.9	100.8	463.6	132.8	142.2	57.9	71.8	460.2
<b>1997</b>	10.5	14.5	28.9	110.0	181.8	175.7	139.6	172.4	444.9	85.6	390.5	111.0	118.7	48.2	443.4
<b>1998</b>	6.3	32.1	32.0	54.3	178.7	234.5	183.3	126.4	146.8	366.4	69.9	316.2	89.8	95.9	397.1
<b>1999</b>	5.3	19.3	71.2	60.7	89.6	236.1	252.4	171.9	111.8	125.8	311.3	58.9	266.3	75.5	414.4
<b>2000</b>	5.4	16.4	43.3	138.0	102.7	120.8	258.1	239.9	154.1	97.0	108.2	265.6	50.2	226.6	416.9
<b>2001</b>	13.5	16.7	36.9	83.7	232.6	137.7	131.1	243.3	213.0	132.4	82.6	91.5	224.4	42.3	542.7
<b>2002</b>	5.8	41.4	37.4	71.1	140.6	312.5	150.6	124.9	218.7	185.4	114.3	70.8	78.2	191.6	499.7
<b>2003</b>	4.8	18.0	93.0	72.5	120.0	189.1	341.0	143.0	111.9	189.5	159.4	97.5	60.3	66.6	588.1
<b>2004</b>	6.0	14.9	40.3	179.4	121.8	161.1	206.5	324.2	128.3	97.1	163.2	136.2	83.2	51.4	557.8
<b>2005</b>	8.5	18.6	33.4	77.6	301.4	163.7	176.3	196.8	291.6	111.7	83.9	139.9	116.5	71.1	520.7
<b>2006</b>	5.4	26.0	41.4	63.9	129.5	401.9	177.5	166.3	175.0	250.9	95.3	71.0	118.2	98.4	499.5
<b>2007</b>	7.5	16.4	56.7	77.4	104.9	171.3	434.7	167.5	148.1	150.9	214.7	80.9	60.2	100.1	506.2
<b>2008</b>	7.3	23.0	36.5	107.7	127.5	137.8	182.9	403.9	146.8	125.6	127.0	179.1	67.4	50.1	504.7
<b>2009</b>	11.9	22.4	50.9	68.8	175.4	165.7	145.6	168.4	350.9	123.4	104.8	105.1	148.1	55.6	457.9
<b>2010</b>	5.5	36.6	49.9	96.9	113.3	231.0	177.8	136.3	148.8	300.1	104.7	88.2	88.4	124.3	431.2
<b>2011</b>	6.8	16.7	81.5	95.1	160.2	149.4	247.6	166.0	120.1	126.8	253.8	87.8	73.9	73.9	464.8
<b>2012</b>	8.7	20.8	37.2	154.6	155.7	208.1	157.5	227.2	143.6	100.5	105.3	209.1	72.3	60.7	442.7
<b>2013</b>	8.0	26.6	46.2	70.8	253.9	203.1	220.2	144.9	197.0	120.5	83.7	86.9	172.4	59.5	414.6
<b>2014</b>	8.9	24.6	59.5	88.6	116.3	327.4	211.2	198.8	123.4	162.3	98.5	67.8	70.4	139.5	383.5
<b>2015</b>	10.4	27.3	55.1	114.1	146.3	150.9	341.6	190.9	169.4	101.6	132.7	79.9	54.9	57.0	423.0
<b>2016</b>	3.7	32.0	61.2	106.5	192.2	195.1	161.1	313.3	163.9	140.2	83.3	107.9	64.9	44.5	389.0
<b>2017</b>	2.9	11.4	71.5	117.6	177.9	254.5	207.6	147.7	269.2	135.8	115.0	67.8	87.6	52.6	351.6

Table 4.20. Model estimates of yellowfin sole age 5 recruitment (millions) from the 2016 and 2017 stock assessments.

<b>Year class</b>	<b>2017 assessment</b>	<b>2016 assessment</b>	<b>Year class</b>	<b>2017 assessment</b>	<b>2016 assessment</b>
<b>1964</b>	733	694	<b>1988</b>	1,646	1,673
<b>1965</b>	734	742	<b>1989</b>	817	831
<b>1966</b>	1,520	1,533	<b>1990</b>	908	925
<b>1967</b>	2,316	2,331	<b>1991</b>	2,004	2,045
<b>1968</b>	2,361	2,380	<b>1992</b>	1,191	1,217
<b>1969</b>	3,100	3,126	<b>1993</b>	1,002	1,027
<b>1970</b>	3,432	3,461	<b>1994</b>	1,010	1,037
<b>1971</b>	2,691	2,716	<b>1995</b>	2,489	2,560
<b>1972</b>	1,862	1,878	<b>1996</b>	1,076	1,104
<b>1973</b>	2,496	2,519	<b>1997</b>	910	926
<b>1974</b>	2,932	2,958	<b>1998</b>	1,103	1,131
<b>1975</b>	1,929	1,947	<b>1999</b>	1,553	1,603
<b>1976</b>	2,424	2,447	<b>2000</b>	1,013	1,040
<b>1977</b>	1,584	1,599	<b>2001</b>	1,419	1,457
<b>1978</b>	1,008	1,018	<b>2002</b>	1,401	1,410
<b>1979</b>	1,937	1,956	<b>2003</b>	2,431	2,373
<b>1980</b>	1,435	1,450	<b>2004</b>	1,201	1,124
<b>1981</b>	4,138	4,182	<b>2005</b>	1,352	1,374
<b>1982</b>	765	774	<b>2006</b>	1,726	1,732
<b>1983</b>	3,407	3,448	<b>2007</b>	1,414	1,478
<b>1984</b>	1,173	1,187	<b>2008</b>	1,529	1,510
<b>1985</b>	897	909	<b>2009</b>	1,662	1,767
<b>1986</b>	1,219	1,236	<b>2010</b>	618	802
<b>1987</b>	1,659	1,684			

Table 4.21. Selected parameter estimates and their standard deviation from the preferred stock assessment model.

parameter	value	std dev	parameter	value	std dev
alpha (q-temp model)	0.11	0.04	1976 total biomass	2,091,300	56,204
beta (q-temp model)	0.10	0.01	1977 total biomass	2,400,400	63,225
mean_log_rec	0.88	0.09	1978 total biomass	2,694,400	69,811
mean sel_slope_fsh (females)	1.15	0.08	1979 total biomass	2,853,500	75,286
mean sel50_fsh (females)	8.80	0.25	1980 total biomass	3,034,900	80,233
mean sel_slope_fsh_males	1.35	0.10	1981 total biomass	3,200,300	84,440
mean sel50_fsh_males	8.10	0.24	1982 total biomass	3,312,100	86,507
sel_slope_srv (females)	1.63	0.09	1983 total biomass	3,289,100	87,202
sel50_srv (females)	5.03	0.07	1984 total biomass	3,509,800	92,839
sel_slope_srv_males	-0.07	0.08	1985 total biomass	3,516,700	95,769
sel50_srv_males	0.02	0.02	1986 total biomass	3,228,100	92,811
Ricker SR logalpha	-4.31	0.52	1987 total biomass	3,183,200	95,063
Ricker SR logbeta	-6.30	0.32	1988 total biomass	3,082,600	94,948
Fmsy	0.11	0.04	1989 total biomass	3,131,000	99,488
log (Fmsy)	-1.44	0.50	1990 total biomass	2,993,900	97,766
ABC_biomass 2017	2,557,000	144,000	1991 total biomass	3,106,100	100,820
ABC_biomass 2018	2,465,000	160,000	1992 total biomass	3,303,500	106,140
msy	404,500	146,700	1993 total biomass	3,328,100	108,860
Bmsy	456,200	83,946	1994 total biomass	3,362,300	110,410
1954 total biomass	1,995,500	112,580	1995 total biomass	3,135,500	106,550
1955 total biomass	1,975,200	101,780	1996 total biomass	3,047,400	105,130
1956 total biomass	1,952,900	91,398	1997 total biomass	3,056,200	106,830
1957 total biomass	1,934,000	82,116	1998 total biomass	2,782,500	101,640
1958 total biomass	1,948,700	73,518	1999 total biomass	2,599,500	97,406
1959 total biomass	1,986,000	64,547	2000 total biomass	2,645,900	98,153
1960 total biomass	1,915,100	54,498	2001 total biomass	2,571,100	96,456
1961 total biomass	1,594,900	41,437	2002 total biomass	2,612,300	97,314
1962 total biomass	1,206,600	24,553	2003 total biomass	2,820,600	103,810
1963 total biomass	906,560	18,470	2004 total biomass	3,028,200	110,740
1964 total biomass	926,620	18,621	2005 total biomass	3,138,200	114,760
1965 total biomass	901,170	18,555	2006 total biomass	3,121,300	115,760
1966 total biomass	936,830	19,341	2007 total biomass	3,131,800	118,000
1967 total biomass	911,250	19,602	2008 total biomass	3,002,500	116,170
1968 total biomass	823,840	19,079	2009 total biomass	2,834,300	113,680
1969 total biomass	850,300	20,485	2010 total biomass	2,903,400	116,550
1970 total biomass	813,650	21,815	2011 total biomass	2,936,000	119,650
1971 total biomass	862,710	24,803	2012 total biomass	2,927,800	122,440
1972 total biomass	936,840	28,696	2013 total biomass	2,883,100	124,240
1973 total biomass	1,188,600	34,496	2014 total biomass	2,680,400	121,130
1974 total biomass	1,433,900	40,835	2015 total biomass	2,699,400	127,400
1975 total biomass	1,786,100	48,903	2016 total biomass	2,852,600	146,360
			2017 total biomass	2,878,200	162,720

Table 4.22. Projections of yellowfin sole female spawning biomass (1,000s t), catch (1,000s t) and full selection fishing mortality rate for seven future harvest scenarios.

Scenarios 1 and 2				Scenario 4			
Maximum Tier 3 ABC harvest permissible				Maximum Tier 3 ABC harvest permissible set at F60			
Female				Female			
Year	spawning biomass	catch	F	Year	spawning biomass	catch	F
2017	872.648	143.093	0.07	2017	872.648	143.093	0.07
2018	872.377	228.409	0.12	2018	884.502	145.995	0.07
2019	844.954	212.064	0.12	2019	890.139	149.873	0.08
2020	793.190	229.442	0.14	2020	870.836	141.006	0.08
2021	721.945	211.323	0.14	2021	835.775	135.962	0.08
2022	664.625	195.463	0.13	2022	804.738	134.927	0.08
2023	632.540	181.232	0.13	2023	790.202	134.842	0.08
2024	615.882	174.126	0.12	2024	784.202	135.041	0.08
2025	609.235	171.959	0.12	2025	784.782	135.805	0.08
2026	607.575	173.433	0.12	2026	787.834	137.642	0.08
2027	610.905	177.186	0.12	2027	794.830	139.832	0.08
2028	618.306	181.607	0.12	2028	806.787	141.935	0.08
2029	626.624	185.758	0.13	2029	820.842	144.020	0.08
2030	632.742	188.606	0.13	2030	832.341	145.736	0.08
Scenario 3				Scenario 5			
Harvest at average F over the past 5 years				No fishing			
Female				Female			
Year	spawning biomass	catch	F	Year	spawning biomass	catch	F
2017	872.648	143.093	0.07	2017	775.899	130.497	0.08
2018	883.837	150.570	0.08	2018	805.239	0	0
2019	887.644	153.257	0.08	2019	869.301	0	0
2020	866.878	143.954	0.08	2020	929.637	0	0
2021	830.620	138.617	0.08	2021	970.871	0	0
2022	798.650	137.418	0.08	2022	995.493	0	0
2023	783.345	137.213	0.08	2023	1018.59	0	0
2024	776.705	137.318	0.08	2024	1049.43	0	0
2025	776.723	138.021	0.08	2025	1082.95	0	0
2026	779.310	139.833	0.08	2026	1121.4	0	0
2027	785.911	142.014	0.08	2027	1162.29	0	0
2028	797.485	144.113	0.08	2028	1203.18	0	0
2029	811.164	146.197	0.08	2029	1246.29	0	0
2030	822.360	147.910	0.08	2030	1288.69	0	0

Table 4.22—continued.

**Scenario 6**

**Determination of whether yellowfin sole are currently overfished**

**B35=601.8**

**Female**

<b>Year</b>	<b>spawning biomass</b>	<b>catch</b>	<b>F</b>
2017	872.648	143.093	0.072055
2018	865.966	271.152	0.139932
2019	821.097	246.624	0.139936
2020	760.904	220.596	0.139936
2021	694.424	204.082	0.139936
2022	642.696	183.566	0.130276
2023	616.917	173.173	0.124755
2024	605.026	168.701	0.122208
2025	601.876	168.354	0.121533
2026	602.741	171.081	0.121719
2027	607.842	175.708	0.122789
2028	616.436	180.737	0.124382
2029	625.533	185.267	0.125918
2030	632.152	188.352	0.126934

**Scenario 7**

**Determination of whether the stock is approaching an overfished condition**

**B35=601.8**

**Female**

<b>Year</b>	<b>spawning biomass</b>	<b>catch</b>	<b>F</b>
2017	872.648	143.093	0.072055
2018	872.377	228.409	0.116667
2019	844.954	212.064	0.116679
2020	793.19	229.442	0.139936
2021	721.945	211.323	0.139936
2022	664.625	195.463	0.134973
2023	632.54	181.232	0.128101
2024	615.882	174.126	0.124533
2025	609.235	171.959	0.12311
2026	607.575	173.433	0.122754
2027	610.905	177.186	0.123439
2028	618.306	181.607	0.124763
2029	626.624	185.758	0.126129
2030	632.742	188.606	0.127043

Table 4-23. Catch and bycatch (t) of other BSAI target species in the yellowfin sole directed fishery from 1992-2016 estimated from a combination of regional office reported catch and observer sampling of the catch.

Table 4-23. (continued).

Table 4.23 (continued).

Species	2010	2011	2012	2013	2014	2015	2016
Pollock	3,749	8,685	11,226	20,246	24,712	21,282	22,324
Arrowtooth Flounder	868	2,338	995	2,012	2,216	1,686	3,252
Pacific Cod	8,649	16,300	19,230	24,382	15,217	12,169	11,988
Groundfish, General	3,048						
Rock Sole	9,030	9,762	8,959	7,737	7,031	9,773	7,948
Flathead Sole	1,895	3,236	2,109	4,191	3,999	3,337	4,105
Sablefish		<1			<1	<1	<1
Atka Mackerel		<1	<1	<1	<1	<1	<1
Pacific ocean Perch		<1		17	<1	<1	3
Rex Sole							
Other flatfish			1,201	388	2,887	1,041	1,136
Squid		<1		<1			
Northern Rockfish							
Dusky Rockfish							
Yellowfin Sole	90,008	136,905	133,719	147,777	139,480	107,955	107,505
English Sole							
Unsp.demersal rockfish							
Bocaccio		1,808	1,924	1,922	1,261		
Rockfish, General		1,969	2,270	2,686	1,969		
Octopus							
Smelt, general							
Chilipepper				<1			
Eels			1.3				
Lingcod							
Jellyfish (unspecified)							
Snails							
Sea cucumber							
Korean horsehair crab							
Kamchatka flounder			110	147		427	285
Sharks						1	11

Table 4-24. Estimated non-target species catch (t) in the yellowfin sole fishery, 2003-2015 (PSC not included).

<b>Row Labels</b>	<b>2003</b>	<b>2004</b>	<b>2005</b>	<b>2006</b>	<b>2007</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>	<b>2011</b>	<b>2012</b>	<b>2013</b>	<b>2014</b>	<b>2015</b>
Benthic urochordata	1671.6	1701.5	674.5	520.1	114.5	347.6	204.7	156.0	133.0	140.8	197.4	116.1	230.1
Birds													
Bivalves	1.5	1.1	1.3	0.3	0.5	1.5	1.3	1.8	1.7	0.7	1.2	0.9	1.4
Brittle star unidentified	34.3	32.3	28.7	20.0	7.6	19.0	5.2	4.2	14.0	13.1	5.9	11.6	2.9
Capelin	0.0	4.5	0.0	0.1	0.3	0.2	0.3	0.7	3.8	2.3	0.2	1.3	1.8
Corals Bryozoans	0.2	0.0	1.2	9.4	0.2	8.3	0.3	0.5	0.9	0.7	3.0	0.8	0.1
Eelpouts	19.1	12.3	7.7	4.5	2.3	5.6	5.2	5.1	29.3	14.3	51.6	69.8	21.1
Eulachon	0.0	0.3	0.0	0.1	5.1	0.0	0.1	0.1	0.5	0.1	0.0	0.7	0.2
Greenlings	0.6	0.7	0.3	0.7	0.5	0.2	0.0	0.1	0.0	0.1	0.0	0.0	0.2
Grenadier					0.3		0.4						
Gunnels					0.0						0.0		0.0
Hermit crab unidentified	87.9	52.0	83.6	26.9	35.8	36.6	15.4	17.0	15.9	9.9	6.3	8.6	4.1
Invertebrate unidentified	556.5	625.8	421.2	177.2	40.0	70.4	30.6	25.9	65.4	121.3	25.2	44.4	6.2
Misc crabs	14.4	21.6	11.9	10.6	28.0	14.1	11.0	11.7	20.2	18.2	39.7	19.8	18.8
Misc crustaceans	0.0	0.2	0.2	2.3	1.4	0.7	1.3	0.9	0.5	0.4	0.6	0.2	0.6
Misc fish	95.8	91.2	66.2	42.5	71.2	66.3	48.8	29.2	40.0	86.2	48.2	69.3	34.8
Misc inverts (worms etc)	0.0	0.1	0.0	0.0	0.0	0.2	0.2	0.1	0.2	0.1	0.2	0.1	0.0
Other osmerids	4.2	4.3	0.5	0.6	35.8	9.8	0.8	2.8	2.1	4.7	1.0	9.2	4.8
Pacific Sand lance	0.0	0.2	0.1	0.0	0.0	0.0	0.0	0.0	0.4	0.2	0.0	0.0	0.1
Pacific Sandfish							0.0	0.0	0.0	0.0	0.0	0.0	0.1
Pandalid shrimp	0.2	0.9	0.1	0.8	0.1	0.3	0.5	0.7	2.3	0.6	2.1	1.0	0.2
Polychaete unidentified	0.0	0.1	0.0	0.4	0.1	0.2	0.1	0.1	0.2	0.1	2.0	0.1	0.1
Scypho jellies	111.9	298.7	115.6	46.8	42.4	145.8	223.2	152.4	307.2	179.3	463.2	805.0	352.0
Sea anemone unidentified	6.3	6.2	2.6	4.9	8.8	24.8	25.5	20.5	14.7	6.2	23.4	5.7	4.2
Sea pens whips	0.0	0.0	0.2	0.0	0.0	0.3	0.2	0.6	0.0	0.1	0.1	0.0	0.0
Sea star	1941.3	1868.0	1611.8	1308.6	1462.0	1829.0	683.7	795.6	1674.0	1732.7	1372.4	2106.5	1816.7
Snails	118.3	191.1	69.7	141.5	95.3	139.6	57.7	57.7	74.7	33.7	46.4	33.7	30.0
Sponge unidentified	11.3	6.8	12.2	3.1	0.4	6.8	69.4	16.5	15.1	14.1	16.6	1.5	2.2
Stichaeidae	0.1	0.0		0.0	0.8	0.2	0.0	0.2	0.4	0.1	0.1	0.4	0.5
Surf smelt					0.0								
urchins dollars cucumbers	2.3	0.3	2.5	0.8	3.4	4.9	7.5	1.3	1.0	0.7	0.8	0.5	0.4
<b>Grand Total</b>	<b>4678</b>	<b>4920</b>	<b>3112</b>	<b>2322</b>	<b>1957</b>	<b>2732</b>	<b>1393</b>	<b>1302</b>	<b>2417</b>	<b>2381</b>	<b>2308</b>	<b>3307</b>	<b>2534</b>

Table 4.25--Yellowfin sole TAC and ABC levels, 1980- 2017.

Year	TAC	ABC	Total catch
1980	117,000	169,000	87,391
1981	117,000	214,500	97,301
1982	117,000	214,500	95,712
1983	117,000	214,500	108,385
1984	230,000	310,000	159,526
1985	229,900	310,000	227,107
1986	209,500	230,000	208,597
1987	187,000	187,000	181,428
1988	254,000	254,000	223,156
1989	182,675	241,000	153,170
1990	207,650	278,900	80,584
1991	135,000	250,600	95,000
1992	235,000	372,000	159,038
1993	220,000	238,000	106,101
1994	150,325	230,000	144,544
1995	190,000	277,000	124,740
1996	200,000	278,000	129,659
1997	230,000	233,000	181,389
1998	220,000	220,000	101,201
1999	207,980	212,000	67,320
2000	123,262	191,000	83,850
2001	113,000	176,000	63,395
2002	86,000	115,000	72,999
2003	83,750	114,000	74,418
2004	86,075	114,000	69,046
2005	90,686	124,000	94,683
2006	95,701	121,000	99,068
2007	136,000	225,000	121,029
2008	225,000	248,000	148,894
2009	210,000	210,000	107,528
2010	219,000	219,000	118,624
2011	196,000	239,000	151,164
2012	202,000	203,000	147,183
2013	198,000	206,000	164,944
2014	184,000	239,800	156,778
2015	149,000	248,800	126,933
2016	144,000	211,700	130,500
2017	154,000	260,800	143,000





Figure 4.1—Yellowfin sole annual catch (1,000s t) in the Eastern Bering Sea from 1954-20167 (top panel) and catch by week (non CDQ) from 2010 – September 2017 (bottom panel).





Figure 4.2--Size composition of the yellowfin sole catch in 2017 (through mid-September), by subarea and total.

**yellowfin sole catch by month in 2017 through September 20**



**yellowfin sole catch by area in 2017 (through September 20)**




Figure 4.3     Yellowfin sole catch by month and area in the Eastern Bering Sea in 2017.




Figure 4.4 (Fishery locations by month).




Figure 4.5. Yellowfin sole CPUE (catch per unit effort in kg/ha) from the annual Bering Sea shelf trawl surveys, 1982-2017.




Figure 4.6. Annual bottom trawl survey biomass point-estimates and 95% confidence intervals for yellowfin sole, 1982-2017.




Figure 4.7. Difference between the 1985-2016 average trawl survey CPUE for yellowfin sole and the 2017 survey CPUE. Open circles indicate that the magnitude of the catch was greater in 2017 than the long-term average, closed circles indicate the catch was greater in the long-term average than in 2017.




Figure 4.8 Estimates of average yellowfin sole weight-at-age (g) from trawl survey observations.




Figure 4.9. Master chronology for yellowfin sole and time series of mean summer bottom temperature and May sea surface temperature for the southeastern Bering Sea (Panel A). All data re normalized to a mean of 0 and standard deviation of 1. Correlations of chronologies with bottom temperature and sea surface temperature are shown in panels B and C, respectively. From Matta et al. 2010.




Figure 4.10. Yellowfin sole length-at-age anomalies, for males and females, and bottom temperature anomalies. Correspondence in these residuals is apparent with a 2-3 year lag effect from the mid-1990s to 2009. Late 1980s and early 1990s pattern may be a density-dependent response in growth from the large 1981 and 1983 year-classes.




Figure 4.11. Results show the temperature anomalies (second row at top as bars) and observed values by age and year. Shadings within the matrix reflects relative weight-at-age (within a row) with darker red being heavier than average.

### temperature-catchability model result




Figure 4.12. Average bottom water temperature from stations less than or equal to 100 m in the Bering Sea trawl survey (bars) and the stock assessment model estimate of q for each year 1982-2017.





Figure 4.13. Fit of the Ricker (1958) stock recruitment model to two distinct stock recruitment time-series data sets (top panel), and the fit to the assessment preferred model (model B, lower panel).




Figure 4.14. Posterior distributions of F<sub>msy</sub> for the two models considered in the stock productivity analysis.




Figure 4.15a    Estimated male fishery selectivity by age and year.




Figure 4.15b. Estimated female fishery selectivity by age and year.




Figure 4.16. Model fit to the survey biomass estimates (top left panel), model estimate of the full selection fishing mortality rate throughout the time-series (top right panel), model estimate of total biomass (middle left panel), the model estimate of survey selectivity (middle right panel) and the estimate of female spawning biomass (bottom left panel).




Figure 4.17. Stock assessment model fit to the time-series of fishery and survey age composition, by sex.




Figure 4.17 (continued).




Figure 4.17 (continued).




Figure 4.17 (continued).




Figure 4.18.--Comparison of the fit to the survey biomass using a fixed  $q$  and the  $q$ -bottom temperature relationship.




Figure 4.19--Year class strength of age 5 yellowfin sole estimated by the stock assessment model. The dotted line is the average of the estimates from 62 years of recruitment.




Figure 4.20.--Posterior distributions of some important parameters estimated by the preferred stock assessment model (from mcmc integration).




Figure 4.21—Retrospective plot of yellowfin sole female spawning biomass estimates (1,000s t), 2008–2016, from the recommended assessment model.




Figure 4.22. Projection of yellowfin sole female spawning biomass (1,000s t) at the average full-selection F from the past 5 years (0.104) through 2030 with  $B_{40\%}$  and  $B_{\text{msy}}$  levels indicated.




Figure 4.23. Phase plane figure of the time-series of yellowfin sole female spawning biomass relative to the harvest control rule with 1975 and 2018 indicated.

## Appendix

IPHC research catch of yellowfin sole		
	number	weight (kg)
2007	707	502
2008	0	0
2009	0	0
2010	898	741



