

## 11. Assessment of the Shortraker Rockfish Stock in the Gulf of Alaska

by  
David M. Clausen and Katy B. Echave

### EXECUTIVE SUMMARY

#### Introduction

For the first time, this year's assessment of shortraker rockfish is presented in its own separate SAFE chapter. In previous assessments since 2005, the assessment for shortraker rockfish was combined with that for the "other slope rockfish" management group in a joint chapter. The rationale for this was that both shortraker rockfish and "other slope rockfish" shared a similar assessment methodology based on the NPFMC's "tier 5" definition of overfishing. However, after the last major assessment for these two management categories in 2009, the Gulf of Alaska Groundfish Plan Team and the NPFMC Scientific and Statistical Committee recommended that they be presented in separate chapters in future SAFE reports.

#### Summary of Changes in Assessment Inputs and 2012 ABC/OFL Recommendations

Assessment methodology in this report is identical to that used in the last full assessment for Gulf of Alaska shortraker rockfish in 2009. The only major new information is biomass estimates from the 2011 trawl survey. This survey showed a large increase in the Gulfwide biomass for shortraker rockfish, which is now at a higher level than in any previous year. As in all previous assessments for shortraker rockfish, current exploitable biomass is based on averaging the biomass estimates in the last three Gulf of Alaska trawl surveys (currently 2007, 2009, and 2011). This results in an exploitable biomass of 48,048 mt for shortraker rockfish, which is an increase of 18% compared to the exploitable biomass in the 2009 assessment. The increase is due to the high biomass in the 2011 survey.

Shortraker rockfish has always been classified into tier 5 in the NPFMC's ABC and OFL definitions. The tier 5 definitions state that  $F_{ABC} \leq 0.75M$ . Using an  $M$  of 0.03 and applying this definition to the exploitable biomass of shortraker rockfish results in a recommended ABC of 1,081 mt in 2012. Geographic apportionment of the ABC amongst management areas of the Gulf of Alaska is based on a weighted average of the percent exploitable biomass distribution for each area in the three most recent trawl surveys. In these computations, each successive survey is given a progressively heavier weighting using factors of 4, 6, and 9, respectively. The new apportionment values for shortraker rockfish are: Western area, 9.59%; Central area, 41.82%; and Eastern area, 48.59%. Applying these percentages to the recommended ABC of 1,081 mt yields the following apportionments for the Gulf in 2012: Western area, 104 mt; Central area, 452 mt; and Eastern area, 525 mt.

Overfishing for a tier 5 species such as shortraker rockfish is defined to occur at a harvest rate of  $F=M$ . Therefore, applying the estimate of  $M$  for shortraker rockfish (0.03) to the estimate of current exploitable biomass (48,048 mt) yields an overfishing catch limit of 1,441 mt for 2012.

**Summary of ABCs and Overfishing Levels for 2012 (mt)**

Shortraker rockfish ABC: Gulfwide, 1,081; Western Area, 104; Central Area, 452; Eastern Area, 525.

Shortraker rockfish overfishing level: Gulfwide, 1,441.

**Summary of Results for Shortaker Rockfish in the Gulf of Alaska**

<b>Quantity</b>	As estimated or <i>specified last year for:</i>		As estimated or <i>recommended this year for:</i>	
	2011	2012	2012	2013
<i>M</i> (natural mortality rate)	0.03	0.03	0.03	0.03
Tier	5	5	5	5
Biomass (t)	40,626	40,626	48,048	48,048
<i>F</i> <sub>OFL</sub>	<i>F=M=0.03</i>	<i>F=M=0.03</i>	<i>F=M=0.03</i>	<i>F=M=0.03</i>
<i>maxF</i> <sub>ABC</sub>	<i>0.75M=0.0225</i>	<i>0.75M=0.0225</i>	<i>0.75M=0.0225</i>	<i>0.75M=0.0225</i>
<i>F</i> <sub>ABC</sub>	0.0225	0.0225	0.0225	0.0225
OFL (t)	1,219	1,219	1,441	1,441
maxABC (t)	914	914	1,081	1,081
ABC (t)	914	914	1,081	1,081
<b>Status</b>	As determined <i>last year for:</i>		As determined <i>this year for:</i>	
	2009	2010	2010	2011
Overfishing	No	n/a	No	n/a

### Summaries for Plan Team

All values are in metric tons.

<b>Stock Assemblage</b>	<b>Year</b>	<b>Biomass</b>	<b>OFL</b>	<b>ABC</b>	<b>TAC</b>	<b>Catch<sup>1</sup></b>
Shortraker Rockfish	2010	40,626	1,219	914	914	457
	2011		1,219	914	914	522
	2012	48,048	1,441	1,081		
	2013		1,441	1,081		

<b>Stock Assemblage</b>	<b>Area</b>	<b>2011 OFL</b>	<b>ABC</b>	<b>TAC</b>	<b>Catch<sup>1</sup></b>	<b>2012 OFL</b>	<b>ABC</b>	<b>2013 OFL</b>	<b>ABC</b>
Shortraker Rockfish	W		134	134	78		104		104
	C		325	325	228		452		452
	E		455	455	216		525		525
	<i>Total</i>	1,219	914	914	522	1,441	1,081	1,441	1,081

<sup>1</sup>Current as of October 3, 2011 (National Marine Fisheries Service, Alaska Region, Sustainable Fisheries Division, P.O. Box 21668, Juneau, AK 99802.).

### Responses to SSC Comments

The only SSC comment specific to this assessment in their Dec. 2009 or Dec. 2010 minutes was to recommend that assessment of shortraker rockfish be presented in its own SAFE chapter separate from “other slope rockfish”.

## INTRODUCTION

The North Pacific Fishery Management Council (NPFMC) established shortraker rockfish (*Sebastes borealis*) as a separate management category in the Gulf of Alaska (GOA) in 2005. Previously, shortraker rockfish had been grouped from 1991 to 2004 with rougheye rockfish (*Sebastes aleutianus*) in the “shortraker/rougheye” management category because the two species are similar in appearance, share the same habitat on the upper continental slope, and often co-occur in hauls. Both species were assigned a single overall ABC (acceptable biological catch) and TAC (total allowable catch), and fishermen were free to harvest either species within this TAC. However, evidence from the NMFS Alaska Groundfish Observer Program indicated that shortraker rockfish were being harvested disproportionately within the shortraker/rougheye group, which raised the possibility that shortraker could become overexploited (Clausen 2004). Because of this concern, the NPFMC decided to establish separate management categories for shortraker and rougheye rockfish starting with the 2005 fishing season.

From 2005 to 2010, the assessment for shortraker rockfish was combined with that for another management group of rockfish in the GOA, “other slope rockfish”. Although shortraker rockfish and “other slope rockfish” were distinct management entities, their assessments were presented in a single SAFE chapter because each group was assessed usually a similar methodology based on the NPFMC’s “tier 5” definition of overfishing. However, in 2010 both the GOA Groundfish Plan Team and the NPFMC Scientific and Statistical Committee (SSC) recommended that future assessments for shortraker rockfish and “other slope rockfish” be presented in separate SAFE chapters. This new chapter focusing only on shortraker rockfish responds to this request.

Shortraker rockfish ranges from Hokkaido Island, Japan, north into the Sea of Okhotsk and the Bering Sea, and through the Aleutian Islands and Gulf of Alaska south to southern California. Its center of abundance appears to be Alaskan waters. In the GOA, adults of this species inhabit a narrow band along the upper continental slope at depths of 300-500 m; outside of this depth interval, abundance decreases considerably (Ito 1999). Shortraker rockfish attains the largest size of all *Sebastes*, with a maximum reported total length of 120 cm (Mecklenburg et al. 2002).

Life history information on shortraker rockfish is extremely sparse. The fish are presumed to be viviparous, as other *Sebastes* appear to be, with internal fertilization and incubation of eggs and with the embryos receiving at least some maternal nourishment. There have been no fecundity studies on shortraker rockfish. One study on reproductive biology of the fish in the northeastern Pacific (most samples were from the GOA) indicated they had a protracted reproductive period, and that parturition (larval release) may take place from February through August (McDermott 1994). Another study indicated the peak month of parturition in Southeast Alaska was April (Westrheim 1975). There is no information on when males inseminate females or if migrations occur for spawning/breeding. Genetic techniques have been used recently to identify a small number of post-larval shortraker rockfish from samples collected in epipelagic waters far offshore in the GOA, which is the only documentation of habitat for this life stage (Kondzela et al. 2007). No data exist on when juvenile fish become demersal in the GOA; in fact, few specimens of juvenile shortraker rockfish <35 cm fork length have ever been caught in this region, so information on this life stage is virtually unknown. Off Kamchatka, juvenile shortraker are reported to become demersal starting at a length of about 10 cm (Orlov 2001). Orlov (2001) has also suggested that shortraker rockfish may undergo extensive migrations in the north Pacific. In his theory, which is mostly based on size compositions of shortraker rockfish in various regions, larvae/post-larvae of this species are transported by currents from the GOA to nursery areas in the Aleutian Islands, where they grow and subsequently migrate back to the GOA Alaska as young adults. More research is needed to substantiate this scenario. As mentioned previously, adults are particularly concentrated in a narrow band along the 300-500 m depth interval of the continental slope. Much of this

habitat is steep and difficult to trawl in the GOA, and observations from a manned submersible also indicated that shortraker rockfish seemed to prefer steep slopes with frequent boulders (Krieger and Ito 1999). Adult shortraker rockfish may also be associated with *Primnoa* spp. corals that are used for shelter (Krieger and Wing 2002). Within the slope habitat, shortraker rockfish tend to have a relatively even distribution when compared with the highly aggregated and patchy distribution of many other rockfish such as Pacific ocean perch (Clausen and Fujioka 2007).

Genetic studies of shortraker rockfish have indicated evidence of stock structure in the GOA (Matala et al. 2004; Gharrett et al. 2003), but additional research is needed to better define this structure. Although not conclusive, the genetic studies do not support Orlov's theory of extensive migrations for shortraker rockfish. No research has been done on the stock structure for any of the "other slope rockfish" species.

In practice, the NPFMC apportions the ABCs and TACs for shortraker rockfish in the GOA into three geographic management areas: the Western, Central, and Eastern Gulf of Alaska. This apportionment is to disperse the catch across the Gulf and prevent possible depletion in one area.

## FISHERY

### Catch History

Official fishery catch statistics for shortraker rockfish in the GOA are only available for 2005-2011, when the species catch was first reported separately for management purposes (Table 11-1). However, catch statistics are available for shortraker and roughey rockfish combined for the years 1991-2004, when both species were classified together into one management group, and these are also listed in Table 11-1. Previous to 1991, shortraker rockfish was classified into larger management groups that included Pacific ocean perch and other species of *Sebastes*, and it is generally not possible to separate out the shortraker catches.

Although official catch statistics for shortraker rockfish started only in 2005, estimates of the Gulfwide catch of shortraker rockfish for the years 1993-2003 were computed in Clausen (2004). These estimates are shown in Table 11-2. The estimates are based on a combination of data from the observer program and the NMFS Alaska regional office, and take into account differences in catch by area and by gear type. The estimates indicate that annual shortraker catch was generally around 1,000-1,500 mt during these years. Annual TACs for the shortraker/roughey group were the major determining factor of these catch amounts; as shown in Table 11-1, the total Gulfwide catch of shortraker/roughey for a given year was generally very similar to the corresponding TAC. The 2005-2011 shortraker rockfish official catches have been consistently lower than any of the estimated catches in previous years. These recent low catches correspond to the years when shortraker rockfish has been in its own management category separate from roughey rockfish. This suggests that the breakup of the shortraker/roughey group may have caused the subsequent reduction in catch of shortraker rockfish, but the exact reasons for the lower catches are unclear. The Central Gulf of Alaska Rockfish Pilot Program (see discussion below in "Description of the Fishery"), in effect since 2007, also caused a reduction in catches.

Non-commercial (research and sport) catches of shortraker rockfish and incidental catch in the Pacific halibut longline fishery are reported and discussed in Appendix 11A.

**Description of the Fishery**

Throughout the 1991-2004 period that shortraker/rougheye rockfish existed as a management category in the GOA, directed fishing was not allowed, and the fish could only be retained as “incidentally-caught” species. This incidental catch status has continued for shortraker rockfish since it became a separate category in 2005. In the years since 2005, shortraker rockfish have been taken mostly in fisheries targeting on rockfish, sablefish, and Pacific halibut, with lesser amounts taken in the walleye pollock and other fisheries (Table 11-3). Shortraker rockfish can both be caught with either trawls or longlines. The percent caught in each gear type is listed in the following tables for the years 1993-2011<sup>1</sup>. Note that for 1993-2004, information on catch by gear is only available for the shortraker/rougheye category and not for shortraker alone.

Shortraker/Rougheye Rockfish												
Gear	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004
Trawl	66.1	51.9	68.9	67.2	66.3	52.8	55.5	57.1	40.1	57.1	61.1	41.8
Longline	33.9	48.1	31.1	32.8	33.7	47.2	44.5	42.9	59.9	42.9	38.9	58.2

Shortraker Rockfish							
Gear	2005	2006	2007	2008	2009	2010	2011
Trawl	48.9	51.8	53.9	46.0	53.5	43.5	66.5
Longline	51.1	48.2	46.1	54.0	46.5	56.5	33.5

From 2005 to 2010, trawl and longline gear each comprised about half the annual catch, but trawl catches have predominated in 2011. Nearly all the longline catch of shortraker rockfish appears to have come as “true” incidental catch in the sablefish or halibut longline fisheries. In rockfish trawl fisheries, however, some of the shortraker is taken by actual targeting that some fishermen call “topping off” (Ackley and Heifetz 2001). “Topping off” works in this way: fishery managers assign all vessels in a directed fishery a maximum retainable amount (MRA) for certain species that may be encountered as incidental catch. If a vessel does not catch its MRA during the course of a directed fishing trip, or the MRA is set overly high (as data presented in Ackley and Heifetz [2001] suggest), before returning to port the vessel may be able to make some target hauls on the incidental species and still not exceed its MRA. Such instances of “topping off” for shortraker rockfish appear to take place in the Pacific ocean perch trawl fishery, especially because shortraker rockfish is the most valuable trawl-caught *Sebastes* rockfish in terms of landed price.

<sup>1</sup> 1993-2008: National Marine Fisheries Service, Alaska Region, Sustainable Fisheries Division, P.O. Box 21668, Juneau, AK 99802. Catches updated through October 3, 2009. 2009-2011: National Marine Fisheries Service, Alaska Region, Catch Accounting System, accessed via the Alaska Fishery Information Network (AKFIN). Catches updated through October 3, 2011.

In 2007, the Central Gulf of Alaska Rockfish Pilot Program was initiated to enhance resource conservation and improve economic efficiency for harvesters and processors who participate in the Central GOA rockfish fishery. This is a five-year rationalization program that establishes cooperatives among trawl vessels which receive exclusive harvest privileges for rockfish management groups (for details, see North Pacific Fishery Management Council 2008). The primary rockfish management groups for the program are Pacific ocean perch, northern rockfish, and pelagic shelf rockfish, but there is a small allocation for shortraker rockfish. As a result of this program, catches of shortraker rockfish taken by trawlers in the Central GOA decreased considerably in 2007 (North Pacific Fishery Management Council 2008), and the catches have remained relatively low there in following years. Catches of shortraker rockfish in the Central GOA are now at some of their lowest levels in the whole time series of catch data since 1991. Other effects of the pilot program include: 1) mandatory at-sea and plant observer coverage for vessels participating in the program, which greatly improves observer data for rockfish in the Central GOA; and 2) extending the season when most trawl-caught shortraker rockfish are taken. Previously, most were taken as incidental catch during the directed “derby-style” trawl fisheries for Pacific ocean perch, northern rockfish, and pelagic shelf rockfish, which mostly occurred during July. In the pilot program, trawling can occur anytime between May 1 and November 15, and catches are now spread over this period.

### **Bycatch**

The only analysis of bycatch in shortraker/rougheye rockfish fisheries of the Gulf of Alaska is that of Ackley and Heifetz (2001), in which they examined data for 1994-96 only. In the hauls they identified as targeting on shortraker/rougheye (most of which were presumably “topping off” hauls as described previously), the major bycatch was arrowtooth flounder, sablefish, and shortspine thornyhead, in descending order by percent.

## Discards

Gulfwide discard rates<sup>2</sup> (% of the total catch discarded within a management category) of shortraker rockfish are listed as follows for the years 1991-2011:

Year	Shortraker/ Rougheye
1991	42.0%
1992	10.4%
1993	26.8%
1994	44.8%
1995	30.7%
1996	22.2%
1997	22.0%
1998	27.9%
1999	30.6%
2000	21.2%
2001	29.1%
2002	20.8%
2003	28.3%
2004	27.6%
	Shortraker
2005	15.1%
2006	23.0%
2007	22.2%
2008	18.1%
2009	31.2%
2010	28.1%
2011	21.5%

These discard rates are higher than those for the three species of *Sebastes* in the GOA that have directed fisheries (Pacific ocean perch, northern rockfish, and dusky rockfish), but are less than those for the “other slope rockfish” category in this region (see chapters in this SAFE report for Pacific ocean perch, northern rockfish, dusky rockfish, and other rockfish).

---

<sup>2</sup> 1991-2008: National Marine Fisheries Service, Alaska Region, Sustainable Fisheries Division, P.O. Box 21668, Juneau, AK 99802. Updated through October 3, 2009. 2009-2011: National Marine Fisheries Service, Alaska Region, Catch Accounting System, accessed via the Alaska Fishery Information Network (AKFIN). Updated through October 3, 2011.



## Management Measures

A timeline of management measures that have affected shortraker rockfish is listed in the following table.

Year	Management Measures
1988	The NPFMC implements the slope rockfish assemblage, which includes shortraker rockfish and the species that will become “other slope rockfish”, together with Pacific ocean perch, northern rockfish, and rougheye rockfish. Previously, <i>Sebastes</i> in Alaska were managed as the “Pacific ocean perch complex” or “other rockfish”.
1988	Apportionment of ABC among management areas in the Gulf (Western, Central, and Eastern) for slope rockfish assemblage is determined based on average percent biomass in previous NMFS trawl surveys.
1991	Slope rockfish assemblage is split into three management subgroups with separate ABCs and TACs: Pacific ocean perch, shortraker/rougheye rockfish, and “other slope rockfish”.
1997	Area apportionment procedure for shortraker/rougheye is changed. Apportionment is now based on 4:6:9 weighting of biomass in the most recent three NMFS trawl surveys.
1999	Trawling is prohibited in the Eastern Gulf east of 140 degrees W. longitude. Eastern Gulf trawl closure becomes permanent with the implementation of FMP Amendments 41 and 58 in 2000 and 2001, respectively.
2005	Shortraker rockfish is split as a separate management entity from rougheye rockfish and now has its own ABC and TAC.
2007	Amendment 68 creates the Central Gulf Rockfish Pilot Program, which affects trawl catches of rockfish in this area.

## DATA

### Fishery Data

#### *Catch*

Detailed catch information for shortraker/rougheye and shortraker rockfish is listed in Table 11-1.

#### *Age Composition*

No shortraker rockfish from the commercial have been aged. Few age samples for this species were collected from the fishery prior to 2003, but since then, annual age collections have ranged between 313 and 747 fish.

### Survey Data

#### *Longline Surveys in the Gulf of Alaska*

Two longline surveys of the continental slope of the Gulf of Alaska provide data on the relative abundance of shortraker rockfish in this region: the earlier Japan-U.S. cooperative longline survey, and

the ongoing Alaska Fisheries Science Center (AFSC) longline survey. These surveys compute relative population numbers (RPNs) and relative population weights (RPWs) for fish on the continental slope as indices of stock abundance. The surveys are primarily directed at sablefish, but also catch considerable numbers of shortraker and rougheye rockfish. Results for both surveys concerning rockfish, however, should be viewed with some caution, as the RPNs and RPWs do not take into account possible effects of competition for hooks with other species caught on the longline, especially sablefish. An analysis of the survey data indicated there was a negative correlation between catch rates of sablefish and shortraker rockfish in the Gulf of Alaska, and that there was likely competition for hooks between species in the surveys (Rodgveller et al. 2008). The study concluded that further research and experiments are needed to better quantify the effects of hook competition and to compute adjustment factors for the surveys' catch rates. Recently, another study compared catch rates of shortraker and rougheye rockfish on survey longline gear with observed densities of these fish around the longline from a manned submersible (Rodgveller et al. 2011). Although the relationship between density and catch rates was not statistically significant, there was a positive slope where  $q$  (catchability) = 0.91. Data on shortraker and rougheye catch rates and behavior from this study suggest that soak times need to be tested to determine if longline survey soak times are adequate for assessing shortraker rockfish abundance trends. Soak times were designed for sablefish abundance indices and may not be long enough to accurately assess all other species encountered.

The cooperative longline survey was conducted annually during 1979-94, but RPNs for rockfish are only available for the years 1979-87 (Sasaki and Teshima 1988). These data are highly variable and difficult to interpret, but suggest that abundance of shortraker rockfish remained stable in the Gulf of Alaska (Clausen and Heifetz 1989). The data also indicate that shortraker rockfish are most abundant in the eastern Gulf of Alaska.

The domestic longline survey has been conducted annually since 1988, and RPNs and RPWs have been computed for each year (Table 11-4). For shortraker rockfish, Gulfwide RPNs have ranged from a low of ~11,000 in 1994 to a high of ~32,000 in 2000. Similarly, lowest and highest Gulfwide RPW values were in these same years. Definite trends in these data over the years are difficult to discern, and the Gulfwide values of RPN and RPW sometimes fluctuate considerably between adjacent years. For example, RPW in 2008 was 39,416, dropped to 25,147 in 2010, and increased to 37,698 in 2011. Some of the fluctuations may be related to changes in the abundance of sablefish, as discussed in the previous paragraph regarding competition for hooks among species.

Similar to the cooperative longline survey, the domestic survey results show that abundance of shortraker rockfish is highest in the eastern Gulf of Alaska: the Yakutat area consistently has by far the greatest RPN and RPW values for shortraker rockfish.

#### *Biomass Estimates from Bottom Trawl Surveys*

Bottom trawl surveys were conducted on a triennial basis in the Gulf of Alaska in 1984 through 1999, and these surveys became biennial starting in 2001 (Table 11-5). The surveys provide much information on shortraker rockfish, including estimates of absolute abundance (biomass) and population length compositions. The trawl surveys have covered all areas of the GOA out to a depth of 500 m (in some surveys to 1,000 m), but the 2001 survey did not sample the eastern GOA. To compensate for this lack of sampling in 2001, survey biomass for this area was estimated by averaging the eastern GOA biomass estimates in the three previous trawl surveys (for details, see Heifetz et al. 2001). Also, the 1984 and 1987 survey results should be treated with some caution. A different, non-standard survey design was used in the eastern Gulf of Alaska in 1984; furthermore, much of the survey effort in the western and central Gulf of Alaska in 1984 and 1987 was by Japanese vessels that used a very different net design

than what has been the standard used by U.S. vessels throughout the surveys. To deal with this latter problem, fishing power comparisons of rockfish catches have been done for the various vessels used in the surveys (for a discussion see Heifetz et al. 1994). Results of these comparisons have been incorporated into the biomass estimates discussed here, and the estimates are believed to be the best available. Even so, the reader should be aware that an element of uncertainty exists as to the standardization of the 1984 and 1987 surveys.

Gulfwide biomass estimates for shortraker rockfish have sometimes shown rather large fluctuations between surveys; for example, biomass was 42,851 mt in 1987 and then decreased to 12,681 mt in 1990. However, the confidence intervals have usually overlapped and differences in the estimates do not appear significant, with three exceptions: the 2003, 2005, and 2009 estimates (42,023, 42,568, and 44,185 mt, respectively) appear to be significantly greater than the 1990 estimate (12,681 mt) (Table 11-5 and Figure 11-1). There has been a general upward trend in the biomass estimates since 1990, with the 2011 biomass of 64,835 mt being much larger than any of the previous years. Spatial distribution of catches of shortraker rockfish in the last three GOA trawl surveys indicate the fish are rather evenly spread along an offshore band along the continental slope, with only a few large catches and virtually no catch near shore (Figure 11-2). Much of the large increase in biomass in 2011 is due to a very large catch of over 1,600 kg in a single haul in the Chirikof area. This is much larger than any previous shortraker catch in this area and is one of the largest catches of shortraker ever in the survey. This unusually large catch is likely responsible at least in part for the very wide confidence bounds of the 2011 biomass estimate and the relatively high coefficient of variation (CV) of 33.1%. Compared with many other species of *Sebastes*, the biomass estimates for shortraker rockfish show relatively moderate confidence intervals and low CVs (compare CVs for shortraker in Table 11-5 vs. those for sharpchin, redstripe, harelequin, and silvergray rockfish in the “Other Rockfish” chapter of this SAFE report). The low CVs are an indication of the generally even distribution of shortraker rockfish that was noted in the introduction of this chapter.

Despite the relative precision of the biomass estimates, however, it is uncertain whether the trawl surveys are accurately assessing abundance of shortraker rockfish. Nearly all the catch of these fish is found on the upper continental slope at depths of 300-500 m. Much of this area in the GOA is not trawlable by the survey's gear because of the area's steep and rocky bottom, except for gully entrances where the bottom is more gradual. Consequently, biomass estimates for shortraker rockfish are mostly based on the relatively few hauls in gully entrances, and they may not be showing a true picture of abundance or abundance trends. An example of one possible problem in the trawl survey results can be seen when RPWs by statistical area for shortraker rockfish in longline surveys are compared with corresponding biomass estimates in the trawl surveys (see Table 11-4 vs. Table 11-5). The longline surveys consistently indicate that shortraker rockfish are most abundant in the Yakutat area, and that this area often comprises >50% of the Gulfwide RPW for this species. In contrast, the trawl survey results by area are much more variable, and the Yakutat area does not stand out as a particular area of abundance. In this case, the longline survey may be providing a better index of abundance by area, as the longline gear can be fished nearly anywhere in the steep 300-500 m slope environment inhabited by shortraker rockfish.

### *Trawl Survey Size Compositions*

Size compositions for shortraker rockfish from the 1990-2007 and 2011 trawl surveys were all unimodal, with almost no fish <35 cm in length (Figure 11-3). However, results from the 2009 trawl survey were different because there was a modest catch of small fish that ranged in sized between 10 and 35 cm long. The reason these small fish occurred in 2009, and not in the other surveys, is unknown. The size compositions indicate that mean length of the shortraker rockfish population in the GOA trawl surveys progressively declined from 61.0 cm in 1990 to 53.9 cm in 2003, followed by increases in 2005, 2007, and 2011, with a mean for the latter year of 60.8 cm. The relatively low mean length in 2009 of 54.3 cm is largely attributable to the fish < 35 cm that were caught that year. The 2001 results may be biased by the fact that they do not include fish from the eastern GOA because this area was not sampled that year. Gulfwide trawl surveys (e.g., Martin and Clausen 1995; Martin 1997; von Szalay et al. 2008 and 2010) have shown shortraker rockfish to be larger in the eastern Gulf of Alaska, and the 2001 survey seems to be missing many fish >70 cm in length compared to the other surveys.

### *Trawl Survey Age Compositions*

Shortraker rockfish have long been considered among the most difficult rockfish species to age. The usual method for determining rockfish ages, i.e., counting annular growth zones on otoliths, did not appear to work because the growth pattern of shortraker otoliths is so unclear. However, Hutchinson (2004) developed a new aging method for this species based on using thin sections of otoliths and on applying an innovative set of aging criteria to determine which growth bands correspond to an annulus. A comparison between his results and those of a previous radiometric study of shortraker rockfish age (Kastelle et al. 2000) indicated general agreement and provided a limited degree of validation. This new aging methodology was used to determine the age compositions of shortraker rockfish in the 1996, 2003, and 2005 GOA trawl surveys (Figure 11-4). Ages ranged from 5 to 146 years, and the results indicate the shortraker rockfish population in the GOA is quite old (mean age varied between 32 and 44 years, depending on the survey). To provide direct validation of the new aging method, in 2008 a validation study was conducted based on carbon-14 levels in shortraker rockfish otoliths from nuclear bomb testing in the 1960s. Results were unsuccessful, however, because carbon-14 could not found in sufficient quantity in the otoliths<sup>3</sup>. Thus, alternative validation techniques will be necessary to verify the aging methodology. One possibility is to conduct an updated and more detailed radiometric study than the previously mentioned Kastelle et al. 2000 study, which was done before the shortraker aging technique had been developed and was somewhat problematic because it was based on using length of the fish as a proxy for age.

Because of the lack of direct validation for the aging method, and the consequent uncertainty about the ages, production aging for shortraker rockfish has now been put on hold. Due to this uncertainty, use of an age-structured model to assess Gulf of Alaska shortraker rockfish is not recommended at present. Although we hope to move to an age-structured assessment at some time in the future, better validation of the shortraker rockfish aging methodology is needed before we do so.

---

<sup>3</sup> C. Hutchinson, National Marine Fisheries Service, Alaska Fisheries Science Center, REFM Division, 7600 Sand Point Way NE, Seattle WA 98115. Pers. commun. Jan. 2009.

## ASSESSMENT PARAMETERS

### Mortality, Maximum Age, Female Age- and Size-at-50% Maturity

Estimates of mortality, maximum age, and female age- and size-at-50% maturity for shortraker rockfish are listed as follows:

Mortality rate <sup>a</sup>	Mortality rate method	Maximum age	Age at Maturity	Size at Maturity	Area	References
-	-	120	-	-	BC	1
0.027-0.042	GSI	-	21.4	44.9	WC,GOA,AL,EBS	2,3
-	-	157	-	-	GOA	4
-	-	146	-	-	GOA	5

Area indicates location of study: British Columbia (BC), West Coast of U.S. (WC), Gulf of Alaska (GOA), Aleutians (AL), and eastern Bering Sea (EBS).

GSI: gonad somatic index (Gunderson and Dygert (1988).

References: 1) Chilton and Beamish 1982; 2) McDermott 1994; 3) Hutchinson 2004; 4) Munk 2001; 5) this report.

The two values for maximum age of shortraker rockfish in the GOA (146 and 157), if true, would make this species one of the longest-lived of all fishes. McDermott (1994) determined that size-at-50% maturity for female shortraker rockfish was 44.9 cm based on samples collected in several regions of the northeast Pacific, including the Gulf of Alaska. Hutchinson's (2004) experimental aging study of shortraker rockfish computed von Bertalanffy growth parameters for females, and he used these parameters to convert McDermott's size-of-maturity to an age-of-50% maturity of 21.4 years. Because it was based on experimental aging, however, and was also determined indirectly, the estimate needs to be confirmed by additional study.

### Length- and Weight-at-Age

Length-weight coefficients and von Bertalanffy parameters for shortraker rockfish are listed below.

Length-weight coefficients are from the formula  $W = aL^b$  where  $W$  = weight in kg and  $L$  = length in cm (based on data from the 1996 GOA trawl survey in Martin 1997):

Sex	a	b	# sampled
combined	$9.85 \times 10^{-6}$	3.13	620
males	$1.26 \times 10^{-5}$	3.07	302
females	$1.02 \times 10^{-5}$	3.12	318

Von Bertalanffy parameters for shortraker rockfish ( GOA = Gulf of Alaska; AI = Aleutian Islands; EBS = Eastern Bering Sea):

Area	Sex	$t_0$	k	$L_{inf}$ (cm)
GOA/AI/EBS	female	-3.62	0.030	84.60

The von Bertalanffy parameters are based on the previously discussed Hutchinson (2004) study which has been only partially validated, so they should be used with caution. Although the analysis combined samples from the the GOA, Aleutian Islands, and eastern Bering Sea, most were from the GOA.

## **ANALYTIC APPROACH AND ABC RECOMMENDATIONS**

Due to the lack of biological information for shortraker rockfish (especially an absence of validated age data), past assessments have all used a biomass-based approach based on trawl survey data to calculate ABCs. We continue to use this approach in the present assessment. As previously mentioned, we anticipate moving to an age-structured assessment for shortraker rockfish at some time in the future if the aging methodology can be successfully validated.

### **Determination of Current Exploitable Biomass**

In all the past SAFE reports, exploitable biomass in the GOA for shortraker/rougheye rockfish or shortraker rockfish as its own management category has been determined based on the average Gulfwide biomass for the three most recent trawl surveys (Clausen 2009). Before the 2007 assessment (Clausen 2007), exploitable biomass computations did not include the biomass in the 1-100 m depth stratum. This was a holdover from a period in the late 1980s when shortraker rockfish was part of a much larger management group that included all slope rockfish, such as Pacific ocean perch and northern rockfish. Pacific ocean perch in the 1-100 m stratum were thought to be mostly small juveniles and therefore not exploitable. However, in the 2007 assessment for shortraker rockfish, an analysis indicated that excluding the 1-100 m stratum in the exploitable biomass calculations was unnecessary because catches of shortraker rockfish in this stratum are negligible in the surveys (Clausen 2007). Since 2007, the exploitable biomass determinations for shortraker rockfish have included all the strata covered by the trawl surveys.

Therefore, current exploitable biomass for shortraker rockfish is calculated based on the average Gulfwide biomass estimates (including the 1-100 m stratum) for the three most recent trawl surveys in 2007, 2009, and 2011 (Table 11-6). The Gulfwide biomasses for these surveys are 35,125 mt, 44,185 mt, and 64,835 mt, respectively, yielding a current exploitable biomass of 48,048 mt for shortraker rockfish in the GOA. This is an increase of 18% over the exploitable biomass of 40,626 mt in the previous assessment (Clausen 2009) and is a reflection of the high biomass found in the 2011 trawl survey.

### **ABC Recommendations for Shortraker Rockfish**

When the shortraker/rougheye category was created in 1991, there was no estimate at that time of  $M$  or  $Z$  for shortraker rockfish. Therefore, the SSC suggested the following computation for a proxy estimate of  $M$ : use the ratio of maximum age of rougheye to shortraker (140/120) from British Columbia and then multiply this value by the mid-point of the range of  $Z$  for rougheye rockfish in British Columbia (mid-point = 0.025) to yield an  $M$  of 0.03 for shortraker rockfish. In a later study,  $M$  for shortraker rockfish was estimated to range between 0.027 and 0.042 (McDermott 1994), so the original estimate of 0.03 for  $M$  seems reasonable.

In previous assessments, shortraker rockfish were always classified as “tier 5” in the NPFMC definitions for ABC and Overfishing Level (OFL) based on Amendment 56 to the Gulf of Alaska FMP. The population dynamics information available for tier 5 species consists of reliable estimates of biomass and natural mortality  $M$ , and the definitions state that for these species, the fishing rate that determines ABC (i.e.,  $F_{ABC}$ ) is  $\leq 0.75M$ . Now that that age and maturity data are available for shortraker rockfish, theoretically this species could be moved into tier 4, where  $F_{ABC} \leq F_{40\%}$ . However, because of the uncertainty of the present aging method and the lack of age validation, we recommend keeping shortraker

rockfish in tier 5 for the present. Thus, the recommended  $F_{ABC}$  for shortraker rockfish is 0.0225 (i.e.,  $0.75 \times M$ , where  $M = 0.03$ ). Applying this  $F_{ABC}$  to the estimate of current exploitable biomass of 48,048 mt for shortraker rockfish results in a Gulfwide ABC of 1,081 mt for 2012. This is an increase of 18% compared to the 2010 and 2011 ABCs of 914 mt. The increase is due to the large biomass in the 2011 trawl survey that now goes into the calculation of current exploitable biomass.

### **Area Allocation of ABC**

Since 1991, the Gulfwide ABC for shortraker/rougheye rockfish or shortraker rockfish alone has been allocated amongst the Western, Central, and Eastern GOA regulatory areas based on the geographic distribution of the species' exploitable biomass in the trawl surveys. Beginning with the 1996 SAFE report, this distribution has been computed as a weighted average of the percent exploitable biomass distribution for each area in the three most recent trawl surveys. In the computations, each successive survey is given a progressively heavier weighting using factors of 4, 6, and 9, respectively. This 4:6:9 weighting scheme was originally recommended by the GOA Groundfish Plan Team, and had already been used for Pacific ocean perch in the 1996 fishery. The Plan Team believed that for consistency among the rockfish assessments, the same weighting should be applied to shortraker/rougheye rockfish. The Plan Team's scheme was adopted for the 1997 fishery, and the scheme has continued to be used in the years since. Therefore, based on a 4:6:9 weighting of the 2007, 2009, and 2011 trawl surveys, the percent distribution of exploitable biomass for shortraker rockfish biomass in the GOA is: Western area, 9.59%; Central area, 41.82%, and Eastern area, 48.59% (Table 11-6). Applying these percentages to the recommended Gulfwide ABC of 1,081 mt yields the following apportionments for the GOA in 2012: Western area, 104 mt; Central area, 452 mt; and Eastern area, 525 mt.

### **Overfishing Level for Shortraker Rockfish**

Based on Amendment 56 in the Gulf of Alaska FMP, overfishing for a tier 5 species such as shortraker rockfish is defined to occur at a harvest rate of  $F=M$ . Therefore, applying the estimate of  $M$  for shortraker rockfish (0.03) to the estimate of current exploitable biomass (48,048 mt) yields an overfishing catch limit of 1,441 mt for 2012.

### **Summary**

A summary of tier, current exploitable biomass, values of  $F$ , and recommended ABC and OFL for shortraker rockfish is listed below for 2012 (biomass and yield are in mt):

Tier	Exploit.	<u>ABC</u>	Yield	<u>Overfishing</u>	Yield
	biomass	$F$		$F$	
5	48,048	$F = 0.75M = 0.0225$	1,081	$F = M = 0.030$	1,441

## ECOSYSTEM CONSIDERATIONS

In general, a determination of ecosystem considerations for shortraker rockfish is hampered by the lack of biological and habitat information. A summary of the ecosystem considerations presented in this section is listed in Table 11-7.

### Ecosystem Effects on the Stock

*Prey availability/abundance trends:* similar to other rockfish species, stock condition of shortraker rockfish is probably influenced by periodic abundant year classes. Availability of suitable zooplankton prey items in sufficient quantity for larval or post-larval rockfish may be an important determining factor of year-class strength. Unfortunately, there is no information on the food habits of larval or post-larval rockfish to help determine possible relationships between prey availability and year-class strength. Moreover, visual identification to the species level for field-collected larval or post-larval rockfish is generally not reliable, although genetic techniques allow identification for larvae/post-larvae of many rockfish, including shortraker (Gharrett et. al 2001; Kondzela et al. 2007). Very few juvenile shortraker rockfish have ever been caught in Alaska, and therefore there is no information on their food items. Adult shortraker rockfish are apparently opportunistic feeders that in Alaska prey on shrimp, deepwater fish such as myctophids, and squid (Yang and Nelson 2000; Yang 2003; Yang et al. 2006). Little if anything is known about abundance trends of these rockfish prey items.

*Predator population trends:* Rockfish are preyed on by a variety of other fish at all life stages, and to some extent by marine mammals during late juvenile and adult stages. Whether the impact of any particular predator is significant or dominant is unknown. Predator effects would likely be more important on larval, post-larval, and small juvenile shortraker rockfish, but information on these life stages and their predators is nil. Due to their large size, older shortraker rockfish likely have few potential predators other than very large animals such as sleeper sharks or sperm whales.

*Changes in physical environment:* Strong year classes corresponding to the period around 1976-77 have been reported for many species of groundfish in the GOA, including Pacific ocean perch, northern rockfish, sablefish, and Pacific cod. Therefore, it appears that environmental conditions may have changed during this period in such a way that survival of young-of-the-year fish increased for many groundfish species, including slope rockfish. The environmental mechanism for this increased survival remains unknown. Changes in water temperature and currents could have an effect on prey item abundance and success of transition of rockfish from the pelagic to demersal stage. Rockfish in early juvenile stage have been found in floating kelp patches which would be subject to ocean currents.

Changes in bottom habitat due to natural or anthropogenic causes could affect survival rates by altering available shelter, prey, or other functions. Associations of juvenile rockfish with biotic and abiotic structure have been noted by Carlson and Straty (1981), Pearcy et al. (1989), Love et al. (1991), and Freese and Wing (2003). A study in the GOA based on observations from a manned submersible found that adult "large" rockfish had a strong association with *Primnoa* spp. coral growing on boulders: less than 1 percent of the observed boulders had coral, but 85 percent of the "large" rockfish were next to boulders with coral (Krieger and Wing 2002). Although the "large" rockfish could not be positively



identified, it is likely based on location and depth that many were shortraker rockfish. The Essential Fish Habitat Environmental Impact Statement (EFH EIS) for groundfish in Alaska (NMFS 2005) concluded that the effects of commercial fishing on the habitat of groundfish is minimal or temporary based largely on the criterion that stocks were above the Minimum Stock Size Threshold (MSST). However, a review of the EFH EIS suggested that this criterion was inadequate to make such a conclusion (Drinkwater 2004). The trend in shortraker abundance suggests that any adverse effect has not prevented the stock from increasing since 1990.

### **Fishery Effects on the Ecosystem**

Because there is only a small amount of targeted fishing on shortraker rockfish in the GOA that is the result of “topping off” by trawlers (see subsection “Description of the Fishery”), most of the catch in the GOA is taken incidentally in longline fisheries for sablefish and Pacific halibut or in the rockfish trawl fishery for Pacific ocean perch. Thus, the reader is referred to the discussions on “Fishery Effects” in the sablefish and Pacific ocean perch chapters in this SAFE report.

*Fishery-specific contribution to bycatch of HAPC biota:* In the GOA, bottom trawl fisheries for shortraker and roughey rockfish accounted for very little bycatch of HAPC biota (Table 11-8). This low bycatch is likely explained by the fact that little targeted fishing occurs for these fish.

*Fishery-specific concentration of target catch in space and time relative to predator needs in space and time (if known) and relative to spawning components:* Unknown.

*Fishery-specific effects on amount of large size target fish:* Unknown.

*Fishery contribution to discards and offal production:* Annual fishery discard rates during 2009-2011 have been 21-31 % for shortraker rockfish. The discard amount of species other than shortraker rockfish in hauls targeting shortraker rockfish is unknown.

*Fishery-specific effects on age-at-maturity and fecundity of the target fishery:* Unknown.

*Fishery-specific effects on EFH non-living substrate:* unknown, but the heavy-duty “rockhopper” trawl gear commonly used in the rockfish fishery can move around rocks and boulders on the bottom.

## **DATA GAPS AND RESEARCH PRIORITIES**

Currently, validation of aging methods for shortraker rockfish is the most important research priority so that an age-structured model can be used for assessment. Also, much additional research is needed on other aspects of shortraker rockfish biology and assessment. There is little to no information on larval, post-larval, or early stage juveniles of shortraker rockfish. In particular, information is lacking on juvenile shortraker rockfish, which are very seldom caught in any sampling gear. Habitat requirements for larval, post-larval, and early stages are mostly unknown. Habitat requirements for later stage juvenile and adult fish are mostly anecdotal or conjectural. Research needs to be done on the bottom habitat of the fishing grounds, on what HAPC biota are found on these grounds, and on what impact bottom trawling has on the grounds. Investigation is needed on the distribution and abundance of shortraker rockfish in areas of rough bottom that cannot be sampled by trawl surveys. Further analyses of the longline survey should be completed to help determine if longline data can be used to assess stock condition of shortraker rockfish.

## REFERENCES

- Ackley, D. R. and J. Heifetz. 2001. Fishing practices under maximum retainable bycatch rates in Alaska's groundfish fisheries. *Alaska Fish. Res. Bull.* 8: 22-44.
- Carlson, H. R., and R. R. Straty. 1981. Habitat and nursery grounds of Pacific rockfish, *Sebastes* spp., in rocky coastal areas of Southeastern Alaska. *Mar. Fish. Rev.* 43: 13-19
- Chilton, D. E. and R. J. Beamish. 1982. Age determination methods for fishes studied by the groundfish program at the Pacific Biological Station. *Can. Spec. Pub. Fish. Aquat. Sci.* 60.
- Clausen, D. M. 2004. Alternative ABCs for shortraker/rougheye rockfish in the Gulf of Alaska. In Stock assessment and fishery evaluation report for the groundfish resources of the Gulf of Alaska, Appendix 9A, p. 416–428. North Pacific Fishery Management Council, 605 W 4<sup>th</sup> Ave, Suite 306, Anchorage AK 99501.
- Clausen, D. M. 2007. Shortraker and other slope rockfish. In Stock assessment and fishery evaluation report for the groundfish resources of the Gulf of Alaska, p. 735–780. North Pacific Fishery Management Council, 605 W 4<sup>th</sup> Ave, Suite 306, Anchorage AK 99501. Available on-line: <http://www.afsc.noaa.gov/refm/docs/2007/GOAshortraker.pdf>
- Clausen, D. M. 2009. Assessment of shortraker and other slope rockfish. In Stock assessment and fishery evaluation report for the groundfish resources of the Gulf of Alaska, p. 875–924. North Pacific Fishery Management Council, 605 W 4<sup>th</sup> Ave, Suite 306, Anchorage AK 99501. Available on-line: <http://www.afsc.noaa.gov/refm/docs/2009/GOAshortraker.pdf>
- Clausen, D. M., and J. T. Fujioka. 2007. Variability in trawl survey catches of Pacific ocean perch, shortraker rockfish, and rougheye rockfish in the Gulf of Alaska. In J. Heifetz, J. Dicosimo, A. J. Gharrett, M. S. Love, V. M. O'Connell, and R. D. Stanley (editors), *Biology, assessment, and management of North Pacific rockfishes*, p. 411-428. Alaska Sea Grant, Univ. of Alaska Fairbanks.
- Clausen, D. M. and J. Heifetz. 1989. Slope rockfish. In T.K. Wilderbuer (editor), *Condition of groundfish resources of the Gulf of Alaska in 1988*, p. 99-149. U.S. Dept. Commer., NOAA Tech. Memo. NMFS F/NWC-165.
- Drinkwater, K. 2004. Summary report: review on evaluation of fishing activities that may adversely affect Essential Fish Habitat (EFH) in Alaska. Center of Independent Experts Review (CIE) June 2004, Alaska Fisheries Science Center, Seattle, Washington.
- Freese, J. L., and B. L. Wing. 2003. Juvenile red rockfish, *Sebastes* sp., associations with sponges in the Gulf of Alaska. *Mar. Fish. Rev.* 65(3): 38-42.
- Gharrett, A. J., A. K. Gray, and J. Heifetz. 2001. Identification of rockfish (*Sebastes* spp.) from restriction site analysis of the mitochondrial NM-3/ND-4 and 12S/16S rRNA gene regions. *Fish. Bull.* 99: 49-62.
- Gharrett, A. J., E. L. Peterson, A. K. Gray, Z. Li, and J. Heifetz. 2003. Population structure of Alaska shortraker rockfish, *Sebastes borealis*, inferred from mitochondrial DNA variation. Fisheries Division, School of Fisheries and Ocean Sciences, Univ. of Alaska Fairbanks, Juneau AK 99801 Unpublished contract report. 21 p.

- Gunderson, D. R., and P. H. Dygert. 1988. Reproductive effort as a predictor of natural mortality rate. *J. Cons. Int. Explor. Mer.* 44: 200-209.
- Heifetz, J., D. M. Clausen, and J. N. Ianelli. 1994. Slope rockfish. *In* Stock assessment and fishery evaluation report for the 1995 Gulf of Alaska groundfish fishery, p. 5-1 - 5-24. North Pacific Fishery Management Council, 605 W 4<sup>th</sup> Ave, Suite 306 Anchorage, AK 99501.
- Heifetz, J., J. N. Ianelli, D. M. Clausen, D. L. Courtney, and J. T. Fujioka. 2001. Slope rockfish. *In* Stock assessment and fishery evaluation report for the groundfish resources of the Gulf of Alaska, p. 6-1 – 6-72. North Pacific Fishery Management Council, 605 W 4<sup>th</sup> Ave, Suite 306 Anchorage, AK 99501.
- Hutchinson, C. E. 2004. Using radioisotopes in the age determination of shorttraker (*Sebastes borealis*) and canary (*Sebastes pinniger*) rockfish. Masters Thesis. Univ. Washington, Seattle. 84 p.
- Ito, D. H. 1999. Assessing shorttraker and rougheye rockfishes in the Gulf of Alaska: addressing a problem of habitat specificity and sampling capability. Ph. D. Thesis. Univ. Washington, Seattle. 204 p.
- Kastelle, C. R., D. K. Kimura, and S. R. Jay. 2000. Using <sup>210</sup>Pb/<sup>226</sup>Ra disequilibrium to validate conventional ages in Scorpaenids (genera *Sebastes* and *Sebastolobus*). *Fish. Res.* 46: 299-312.
- Kondzela, C. M., A. W. Kendall, Z. Li, D. M. Clausen, and A. J. Gharrett. 2007. Preliminary identification of pelagic juvenile rockfishes collected in the Gulf of Alaska. *In* J. Heifetz, J. DiCosimo, A.J. Gharrett, M.S. Love, V.M. O'Connell, and R.D. Stanley (editors), *Biology, assessment, and management of North Pacific rockfishes*, p. 153-166. Alaska Sea Grant, Univ. of Alaska Fairbanks.
- Krieger, K. J., and D. H. Ito. 1999. Distribution and abundance of shorttraker rougheye, *Sebastes borealis*, and rougheye rockfish, *S. aleutianus*, determined from a manned submersible. *Fish. Bull.* 97: 264-272.
- Krieger, K.J., and B.L. Wing. 2002. Megafauna associations with deepwater corals (*Primnoa* spp.) in the Gulf of Alaska. *Hydrobiologia* 471: 83-90.
- Love, M .S., M. H. Carr, and L. J. Haldorson. 1991. The ecology of substrate-associated juveniles of the genus *Sebastes*. *Environmental Biology of Fishes* 30:225-243.
- Martin, M. H. 1997. Data report: 1996 Gulf of Alaska bottom trawl survey. U.S Dept. Commer. NOAA Tech. Memo. NMFS-AFSC-82. 235 p.
- Martin, M. H., and D. M. Clausen. 1995. Data report: 1993 Gulf of Alaska bottom trawl survey. U.S Dept. Commer. NOAA Tech. Memo. NMFS-AFSC-59. 217 p.
- Matala, A. P., A. K. Gray, J. Heifetz, and A. J. Gharrett. 2004. Population structure of Alaska shorttraker rockfish, *Sebastes borealis*, inferred from microsatellite variation. *Environ. Biol. Fishes.* 69: 201-210.
- McDermott, S.F. 1994. Reproductive biology of rougheye and shorttraker rockfish, *Sebastes aleutianus* and *Sebastes borealis*. Masters Thesis. Univ. Washington, Seattle. 76 p.

- Mecklenburg, C. W., T. A. Mecklenburg, and L. K. Thorsteinson. 2002. Fishes of Alaska. American Fisheries Society, Bethesda, Maryland. 1,037 p.
- Munk, K. M. 2001. Maximum ages of groundfishes in waters off Alaska and British Columbia and considerations of age determination. Alaska Fish. Res. Bull. 8(1): 12-21.
- National Marine Fisheries Service. 2005. Final environmental impact statement for essential fish habitat identification and conservation in Alaska. Available on-line:  
<http://www.fakr.noaa.gov/habitat/seis/efheis.htm>.
- North Pacific Fishery Management Council. 2008. Gulf of Alaska rockfish pilot program review. Unpubl. report, 35 p. North Pacific Fishery Management Council, 605 W 4<sup>th</sup> Ave, Suite 306 Anchorage, AK 99501. Available on-line:  
[http://www.fakr.noaa.gov/npfmc/current\\_issues/groundfish/RPPreview508.pdf](http://www.fakr.noaa.gov/npfmc/current_issues/groundfish/RPPreview508.pdf)
- Orlov, A. M. 2001. Ocean current patterns and aspects of life history of some northwestern Pacific scorpaenids. In: G. H. Kruse, N. Bez, A. Booth, M. W. Dorn, A. Hills, R. N. Lipcius, D. Pelletier, C. Roy, S. J. Smith, and D. Witherell (editors), Spatial processes and management of marine populations. Pub. No. AK-SG-01-02. Univ. Alaska Sea Grant College Program, Fairbanks AK.
- Pearcy, W. G., D. L. Stein, M. A. Hixon, E. K. Pikitch, W. H. Barss, and R. M. Starr. 1989. Submersible observations of deep-reef fishes of Heceta Bank, Oregon. Fish. Bull. 87: 955-965.
- Rodgveller, C. J., C. R. Lunsford, and J. T. Fujioka. 2008. Evidence of hook competition in longline surveys. Fish. Bull. 106: 364-374.
- Rodgveller, C. J., M. F. Sigler, D. H. Hanselman, and D. H. Ito. 2011. Sampling efficiency of longlines for shortraker and roughey rockfish using observations from a manned submersible. Mar. Coast. Fish: Dynamics, Management, and Ecosystem Sci. 3: 1-9.
- Sasaki, T., and K. Teshima. 1988. Data report of abundance indices of flatfishes, rockfishes, and shortspine thornyhead and grenadiers based on results from Japan-U.S. joint longline surveys, 1979-1987. Unpubl. manuscr., 5 p. (Document submitted to the annual meeting of the International North Pacific Fisheries Commission, Tokyo, Japan, October 1988.) Fisheries Agency of Japan, Far Seas Fisheries Research Laboratory, 5-7-1 Orido, Shimizu, Japan 424.
- von Szalay, P. G., M. E. Wilkins, and M. M. Martin. 2008. Data report: 2007 Gulf of Alaska bottom trawl survey. U.S Dept. Commer. NOAA Tech. Memo. NMFS-AFSC-189. 247 p.
- von Szalay, P. G., N. W. Raring, F. R. Shaw, M. E. Wilkins, and M. M. Martin. 2010. Data report: 2009 Gulf of Alaska bottom trawl survey. U.S Dept. Commer. NOAA Tech. Memo. NMFS-AFSC-208. 245 p.
- Westrheim, S.J. 1975. Reproduction, maturation, and identification of larvae of some *Sebastes* (Scorpaenidae) species in the northeast Pacific Ocean. J. Fish. Res. Board Can. 32:2399-2411.
- Yang, M-S., and M. W. Nelson. 2000. Food habits of the commercially important groundfishes in the Gulf of Alaska in 1990, 1993, and 1996. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-AFSC-112, 174 p.

Yang, M-S. 2003. Food habits of the important groundfishes in the Aleutian Islands in 1994 and 1999. AFSC Proc. Rep 2003-07. 233 p. (Available from National Marine Fisheries Service, Alaska Fisheries Science Center, 7600 Sand Point Way NE, Seattle WA 98115).

Yang, M-S., K. Dodd, R. Hibpshman, and A. Whitehouse. 2006. Food habits of groundfishes in the Gulf of Alaska in 1999 and 2001. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-AFSC-164, 199 p.

Table 11-1.--Commercial catch (mt) of fish in the shortraker/rougheye rockfish and shortraker rockfish management categories in the Gulf of Alaska, with Gulfwide values of acceptable biological catch (ABC) and total allowable catch (TAC), 1991-2011. Updated through October 3, 2011.

Year	<u>Area of Gulf</u>			Gulfwide total	Gulfwide ABC	Gulfwide TAC
	Western	Central	Eastern			
<u>Shortraker/Rougheye Rockfish</u>						
1991	123	408	171	702	2,000	2,000
1992	115	1,367	683	2,165	1,960	1,960
1993	85	1,197	650	1,932	1,960	1,764
1994	114	996	722	1,832	1,960	1,960
1995	216	1,222	812	2,250	1,910	1,910
1996	127	941	593	1,661	1,910	1,910
1997	137	931	541	1,609	1,590	1,590
1998	129	870	735	1,734	1,590	1,590
1999	194	580	537	1,311	1,590	1,590
2000	137	887	721	1,745	1,730	1,730
2001	126	998	852	1,976	1,730	1,730
2002	263	631	429	1,323	1,620	1,620
2003	225	856	321	1,402	1,620	1,620
2004	277	337	383	997	1,318	1,318
<u>Shortraker Rockfish</u>						
2005	70	223	205	498	753	753
2006	91	303	270	664	843	843
2007	194	164	250	608	843	843
2008	133	244	221	598	898	898
2009	155	209	223	588	898	898
2010	64	136	257	457	914	914
2011	78	228	216	522	914	914

Sources: Catch: 1991-2008: National Marine Fisheries Service, Alaska Region, Sustainable Fisheries Division, P.O. Box 21668, Juneau, AK 99802. Updated through October 3, 2009. 2009-2011: National Marine Fisheries Service, Alaska Region, Catch Accounting System, accessed via the Alaska Fishery Information Network (AKFIN). Updated through October 3, 2011. ABC and TAC: 1991-2007, Clausen (2007); 2008- 2011, North Pacific Fishery Management Council website (<http://www.fakr.noaa.gov/npfmc/Council0910specs.pdf>).

Table 11-2.--Estimated commercial catch (mt) of shortraker rockfish in the Gulf of Alaska, 1993-2003, based on data from the NMFS Alaska Observer Program database and from the NMFS Alaska Regional Office. See Clausen (2004) for an explanation of how these numbers were estimated.

Year	Catch
1993	1,348
1994	1,254
1995	1,545
1996	1,102
1997	1,065
1998	1,069
1999	992
2000	1,214
2001	1,385
2002	1,051
2003	1,010

Table 11-3.--Estimated catch (%) of shorttraker rockfish in the Gulf of Alaska by target fishery, 2005-2011.

Year	Target Fishery					Total
	Rockfish	Sablefish	Halibut	Pollock	Other	
2005	50	38	6	3	4	100
2006	38	25	22	10	5	100
2007	45	32	12	8	3	100
2008	36	29	22	9	4	100
2009	42	25	23	3	7	100
2010	29	49	17	2	3	100
2011	44	25	11	4	16	100

Source: National Marine Fisheries Service, Alaska Region, Catch Accounting System, accessed via the Alaska Fishery Information Network (AKFIN). Updated through October 17, 2011.



Table 11-4.--Relative population number (RPN) and relative population weight (RPW) for Gulf of Alaska shortraker rockfish in the Alaska Fishery Science Center longline survey, 1988-2011. Data are for the upper continental slope only, 201-1,000 m. depth (gullies are not included).

	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
<u>Shortraker RPN:</u>												
Shumagin	4,492	3,272	3,015	3,074	1,660	1,523	2,549	5,765	4,098	2,888	4,630	5,011
Chirikof	1,290	858	773	776	572	229	613	531	646	918	973	823
Kodiak	2,332	2,691	3,476	2,412	1,374	1,067	1,040	1,325	2,231	2,200	2,498	3,078
Yakutat	5,830	6,492	9,281	10,575	9,130	7,121	5,222	7,992	8,409	12,408	15,295	13,394
Southeastern	1,420	1,972	1,403	2,247	1,479	2,199	1,862	2,427	1,967	2,459	3,258	3,167
Total	15,364	15,285	17,948	19,085	14,214	12,139	11,286	18,039	17,352	20,873	26,654	25,473
<u>Shortraker RPW:</u>												
Shumagin	4,869	4,301	5,004	5,953	2,078	2,192	3,956	7,940	5,946	4,468	6,716	6,954
Chirikof	2,591	1,449	1,216	1,384	914	293	1,174	812	1,007	1,471	1,422	1,165
Kodiak	5,043	5,833	6,787	4,874	2,802	1,912	2,649	2,554	4,657	4,273	5,201	5,562
Yakutat	13,320	13,335	19,093	20,585	17,033	14,411	11,046	15,248	17,352	26,830	30,685	26,500
Southeastern	2,474	3,384	2,214	3,546	2,053	4,124	3,102	4,034	3,377	3,970	5,818	4,569
Total	28,297	28,302	34,313	36,343	24,880	22,932	21,927	30,588	32,338	41,013	49,842	44,750
	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
<u>Shortraker RPN:</u>												
Shumagin	9,481	5,150	3,386	3,576	6,477	2,041	3,901	3,566	3,349	4,633	4,529	8,188
Chirikof	1,298	1,031	951	809	474	274	931	714	813	482	804	1,331
Kodiak	2,904	3,703	1,982	1,510	1,409	1,807	3,080	4,200	3,748	5,967	2,346	3,928
Yakutat	13,995	14,177	9,942	7,312	7,519	6,963	7,970	13,169	12,517	10,124	6,244	7,703
Southeastern	4,025	2,646	3,098	3,951	2,874	1,905	2,106	2,876	2,536	2,292	1,837	2,227
Total	31,703	26,706	19,358	17,158	18,754	12,990	17,989	24,524	22,964	23,498	15,759	23,377
<u>Shortraker RPW:</u>												
Shumagin	15,050	7,314	4,978	5,874	9,678	3,458	5,830	4,944	4,827	6,390	6,375	11,708
Chirikof	1,607	1,682	1,324	1,420	624	378	969	1,067	1,129	659	1,423	1,975
Kodiak	5,553	7,413	3,305	2,908	2,496	3,144	6,086	8,003	6,120	11,487	3,622	7,101
Yakutat	28,754	28,382	18,314	14,583	14,292	12,751	14,056	22,684	21,605	17,340	10,724	12,747
Southeastern	7,099	4,574	5,598	7,455	5,045	2,946	3,203	4,914	4,140	3,541	3,004	4,167
Total	58,063	49,365	33,518	32,240	32,134	22,677	30,144	41,612	37,821	39,416	25,147	37,698

Source: 1988-2009: C. Lunsford, National Marine Fisheries Service, Alaska Fisheries Science Center, Auke Bay Laboratories, 17109 Pt. Lena Loop Rd., Juneau AK 99801. Pers. commun. October 15, 2009. 2010-2011: AFSC longline survey database accessed via the Alaska Fishery Information Network (AKFIN).

Table 11-5.--Biomass estimates (mt) for shortraker rockfish in the Gulf of Alaska, by statistical area, based on bottom trawl surveys conducted between 1984 and 2011. Gulfwide 95% confidence bounds, variance, and coefficient of variation (CV) are also shown for each year.

Year	Statistical areas					Gulfwide Total	Gulfwide 95% Conf. bounds		Biomass variance	Biomass CV (%)
	Shumagin	Chirikof	Kodiak	Yakutat	South-eastern		Lower	Upper		
	<u>Shortraker Rockfish</u>									
1984	4,874	659	4,685	6,288	2,051	18,557	4,600	32,515	34,829,252	31.8
1987	3,232	13,182	18,950	4,408	3,079	42,851	13,392	72,311	196,602,336	32.7
1990	284	1,729	3,027	6,037	1,604	12,681	6,412	18,951	9,085,499	23.8
1993	2,775	2,320	4,735	7,740	1,903	19,472	11,290	27,654	15,474,771	20.2
1996	1,905	2,406	7,726	4,523	3,699	20,258	10,652	29,865	20,532,868	22.4
1999	2,208	3,931	8,459	9,831	3,845	28,275	16,841	39,709	30,393,883	19.5
2001*	4,313	1,589	11,513	7,350	3,149	27,914	18,819	37,008	21,530,717	16.6
2003	11,166	2,996	14,292	11,936	1,633	42,023	23,572	60,474	81,168,454	21.4
2005	5,946	6,342	10,741	16,866	2,673	42,568	25,603	59,532	69,018,739	19.5
2007	2,492	1,911	8,275	8,197	14,250	35,125	17,296	52,954	66,950,870	23.3
2009	8,810	3,209	13,541	12,518	6,109	44,185	25,332	63,039	79,840,212	20.2
2011	2,464	23,382	9,113	22,561	7,316	64,835	18,028	111,643	461,441,570	33.1

\*The 2001 survey did not sample the eastern Gulf of Alaska (Yakutat and Southeastern areas). Substitute estimates of biomass for these areas in 2001 were obtained by averaging the Yakutat and Southeastern biomass in the 1993, 1996, and 1999 surveys. These eastern Gulf of Alaska estimates have been included in the 2001 biomass estimates, confidence bounds, biomass variances, and biomass CVs listed in this table.

Table 11-6.--Biomass estimates (mt) and the percent of biomass for shortraker rockfish in the Gulf of Alaska, by NPFMC regulatory area, for the three most recent trawl surveys in 2007, 2009, and 2011. Weighted averages use weights of 4:6:9 for the 2007, 2009, and 2011 surveys, respectively.

Year	Regulatory area			Total
	Western	Central	Eastern	
2007	2,492	10,186	22,447	35,125
2009	8,810	16,749	18,626	44,185
2011	2,464	32,495	29,877	64,835
average				48,048
	<u>Percent of biomass by area</u>			
2007	7.10%	29.00%	63.91%	100.00%
2009	19.94%	37.91%	42.16%	100.00%
2011	3.80%	50.12%	46.08%	100.00%
	<u>4:6:9 weighted average</u>			
	9.59%	41.82%	48.59%	100.00%

Table 11-7.-- Analysis of ecosystem considerations for shortraker rockfish.

<i>Indicator</i>	<i>Observation</i>	<i>Interpretation</i>	<i>Evaluation</i>
<b><i>ECOSYSTEM EFFECTS ON STOCK</i></b>			
<b><i>Prey availability or abundance trends</i></b>	important for larval and post-larval survival, but no information known	may help to determine year class strength	possible concern if some information available
<b><i>Predator population trends</i></b>	unknown		little concern for adults
<b><i>Changes in habitat quality</i></b>	variable	variable recruitment	possible concern
<b><i>FISHERY EFFECTS ON ECOSYSTEM</i></b>			
<b><i>Fishery contribution to bycatch</i></b>			
Prohibited species	unknown		
Forage (including herring, Atka mackerel, cod, and pollock)	unknown		
HAPC biota (seapens/whips, corals, sponges, anemones)	fishery disturbing hard-bottom biota, i.e., corals, sponges	could harm the ecosystem by reducing shelter for some species	concern
Marine mammals and birds	probably few taken		little concern
Sensitive non-target species	unknown		
<b><i>Fishery concentration in space and time</i></b>	little overlap between fishery and reproductive activities	fishery does not hinder reproduction	little concern
<b><i>Fishery effects on amount of large size target fish</i></b>	unknown		
<b><i>Fishery contribution to discards and offal production</i></b>	discard rates moderate	some unnatural input of food into the ecosystem	some concern
<b><i>Fishery effects on age-at-maturity and fecundity</i></b>	unknown		

Table 11-8. Average bycatch (kg) and bycatch rates during 1997 - 99 of living substrates in the Gulf of Alaska; POT - pot gear; BTR - bottom trawl; HAL - Hook and line (source - Draft Programmatic SEIS).

Target fishery	Gear	Bycatch (kg)				Target catch (mt)	Bycatch rate (kg/mt target)			
		Coral	Anemone	Sea whips	Sponge		Coral	Anemone	Sea whips	Sponge
Arrowtooth flounder	POT	0	0	0	0	4	0.0000	0.0000	0.0000	0.0000
Arrowtooth flounder	BTR	58	99	13	24	2,097	0.0276	0.0474	0.0060	0.0112
Deep water flatfish	BTR	1,626	481	5	733	2,001	0.8124	0.2404	0.0024	0.3663
Rex sole	BTR	321	306	11	317	2,157	0.1488	0.1417	0.0053	0.1468
Shallow water flatfish	POT	0	0	0	0	5	0.0000	0.0000	0.0000	0.0000
Shallow water flatfish	BTR	53	4,741	115	403	2,024	0.0261	2.3420	0.0567	0.1993
Flathead sole	BTR	3	267	1	136	484	0.0071	0.5522	0.0019	0.2806
Pacific cod	HAL	28	4,419	961	33	10,765	0.0026	0.4105	0.0893	0.0030
Pacific cod	POT	0	14	0	1,724	12,863	0.0000	0.0011	0.0000	0.1340
Pacific cod	BTR	34	5,767	895	788	37,926	0.0009	0.1521	0.0236	0.0208
Pollock	BTR	1,153	55	0	23	2,465	0.4676	0.0222	0.0000	0.0092
Pollock	PTR	41	110	0	0	97,171	0.0004	0.0011	0.0000	0.0000
Demersal shelf rockfish	HAL	0	0	0	141	226	0.0000	0.0000	0.0000	0.6241
Northern rockfish	BTR	25	90	0	103	1,938	0.0127	0.0464	0.0000	0.0532
Other slope rockfish	HAL	0	0	0	0	14	0.0000	0.0000	0.0000	0.0000
Other slope rockfish	BTR	0	0	0	0	193	0.0000	0.0000	0.0000	0.0000
Pelagic shelf rockfish	HAL	0	0	0	0	203	0.0000	0.0000	0.0000	0.0000
Pelagic shelf rockfish	BTR	324	176	3	245	1,812	0.1788	0.0969	0.0017	0.1353
Pacific ocean perch	BTR	549	90	5	1,968	6,564	0.0837	0.0136	0.0007	0.2999
Pacific ocean perch	PTR	7	0	0	55	1,320	0.0052	0.0000	0.0000	0.0416
Shortraker/rougheye	HAL	6	0	0	0	19	0.3055	0.0000	0.0000	0.0000
Shortraker/rougheye	BTR	0	18	0	0	21	0.0000	0.8642	0.0000	0.0000
Sablefish	HAL	156	154	68	27	11,143	0.0140	0.0138	0.0061	0.0025
Sablefish	BTR	0	0	0	0	27	0.0000	0.0000	0.0000	0.0000
Shortspine thornyhead	HAL	0	0	0	0	2	0.0000	0.0000	0.0000	0.0000
Shortspine thornyhead	BTR	0	9	0	1	2	0.0000	4.8175	0.0000	0.4069




Figure 11-1.--Estimated biomass of shortraker rockfish in the Gulf of Alaska based on results of bottom trawl surveys from 1984 through 2011. The vertical bars show the 95% confidence limits associated with each estimate. The eastern Gulf of Alaska was not sampled in the 2001 survey, but substitute estimates of biomass and confidence limits for this region in 2001 were calculated and included in the above graph.




Figure 11-2.--Spatial distribution of shorttraker rockfish catches in the Gulf of Alaska during the 2007, 2009, and 2011 NMFS bottom trawl surveys.




Figure 11-3.--Size composition of the estimated population of shortraker rockfish in the Gulf of Alaska based on trawl surveys conducted between 1990 and 2011. (Figure continued on next page.)






Figure 11-3.--Continued.




Figure 11-4.--Age composition of the estimated population of shorttraker rockfish in the 1996, 2003, and 2005 Gulf of Alaska trawl surveys.

## Appendix 11A – Supplemental Catch Data

Research catches of shortraker rockfish for the years 1977-2009 are listed in Table 11A-1. Although data are not available for a complete accounting of all research catches, the values in the table indicate that generally these catches have been modest. The one exception is 1999, when a total of almost 110 mt was taken, mostly by research trawling. However, because commercial catches for shortraker/rougheye rockfish this year were below ABC by nearly 280 mt (see Table 11-1 in the main document), this relatively large catch was not a conservation concern.

In order to comply with the Annual Catch Limit (ACL) requirements, two new datasets have been generated to help estimate total catch and removals from NMFS stocks in Alaska. The first dataset, non-commercial removals, estimates total removals that do not occur during directed groundfish fishing activities (Table 11A-2). This includes removals incurred during research, subsistence, personal use, recreational, and exempted fishing permit activities, but does not include removals taken in fisheries other than those managed under the groundfish FMP. These estimates represent additional sources of removals to the existing Catch Accounting System estimates and are only available for 2010. The second dataset, Halibut Fishery Incidental Catch Estimation (HFICE), is an estimate of the incidental catch of groundfish in the halibut IFQ longline fishery in Alaska, which at present is largely unobserved (Table 11A-3). To estimate removals in the halibut fishery, methods were developed by the HFICE working group and approved by the Gulf of Alaska and Bering Sea/Aleutian Islands Groundfish Plan Teams and the Scientific and Statistical Committee of the North Pacific Fishery Management Council. A detailed description of the methods is available in Tribuzio et al. (2011). The HFICE estimates should be considered preliminary estimates for what is caught in the IFQ halibut fishery. Improved estimates of groundfish catch in the halibut fishery may become available following restructuring of the Observer Program in 2013.

The non-commercial removals show that a total of almost 15 mt of shortraker rockfish was taken in 2010 during research cruises and in sport fisheries (Table 11A-2). Nearly all was taken in longline surveys by either the International Pacific Halibut Commission or the NMFS Alaska Fishery Science Center. This total is only about 3% of the reported commercial catch of 457 mt for shortraker rockfish in 2010 (see table 11-1 in the main document). Therefore, this presents no risk to the stock especially because commercial catches in recent years have been much less than ABCs.

Estimated catches of shortraker rockfish in the Pacific halibut longline fishery have been much higher than the non-commercial removals and range from 27 mt in 2003 to 166 mt in 2009 (Table 11A-3). Although a significant portion of the reported catches of shortraker rockfish in the NMFS Alaska Region Catch Accounting System come from the halibut longline fishery (Table 11-3 in main document), the data in Table 11A-3 suggest that the actual amount taken in this fishery may be even higher. For example, Table 11-3 indicates that 17% of the shortraker catch in 2010 (78 mt) was taken in the halibut fishery, whereas Table 11A-3 estimates 166 mt were taken in this fishery. This level of unaccounted catch, although relatively high compared to ABCs, does not appear to have put stocks of shortraker rockfish at risk in recent years because the annual catch of this species in the GOA has consistently been much less than ABC.

### Reference:

Tribuzio, C. A, S. Gaichas, J. Gasper, H. Gilroy, T. Kong, O. Ormseth, J. Cahalan, J. DiCosimo, M. Furuness, H. Shen, and K. Green. 2011. Methods for the estimation of non-target species catch in the unobserved halibut IFQ fleet. August 2011 Groundfish Plan Team document. Presented to the Joint Groundfish Plan Teams of the North Pacific Fishery Management Council.

Table 11A-1.--Catch (mt) of shortraker rockfish taken during NMFS research cruises in the Gulf of Alaska, 1977-2009. Longline data refers only to catches in the AFSC longline survey and does not include the International Pacific Halibut Commission longline survey. (n.a.=not available; tr=trace).

Year	Gear		Total
	Trawl	Longline	
1977	0.1	0.0	0.1
1978	0.6	n.a.	0.6
1979	0.5	n.a.	0.5
1980	1.0	n.a.	1.0
1981	6.2	n.a.	6.2
1982	2.4	n.a.	2.4
1983	0.2	n.a.	0.2
1984	6.8	n.a.	6.8
1985	3.5	n.a.	3.5
1986	0.9	n.a.	0.9
1987	15.5	n.a.	15.5
1988	0.0	n.a.	0.0
1989	0.1	n.a.	0.1
1990	2.4	n.a.	2.4
1991	tr	n.a.	tr
1992	0.1	n.a.	0.1
1993	3.0	n.a.	3.0
1994	0.1	n.a.	0.1
1995	tr	n.a.	tr
1996	4.3	5.9	10.2
1997	0.0	11.1	11.1
1998	20.7	9.7	30.4
1999	101.5	8.1	109.6
2000	0.0	10.0	10.0
2001	1.0	7.1	8.1
2002	0.5	6.1	6.6
2003	4.3	5.5	9.8
2004	0.0	4.7	4.7
2005	4.1	4.5	8.6
2006	0.0	6.0	6.0
2007	4.7	7.9	12.6
2008	0.0	8.4	8.4
2009	8.3	6.7	15.0

Table 11A-2.--Estimated research and sport catches (mt) of shortraker rockfish in the Gulf of Alaska in 2010, based on data provided by the NMFS Alaska Regional Office (AK R.O.). IPHC longline = International Pacific Halibut Commission longline survey; AFSC longline = NMFS Alaska Fishery Science Center longline survey.

Source	Research trawl	IPHC longline	AFSC longline	Sport	Total
AK R.O.	0.1	3.6	10.5	0.5	14.7

Table 11A-3.--Estimated catch (mt) of shortraker rockfish in the Gulf of Alaska halibut fishery, 2001-2010, from the Halibut Fishery Incidental Catch Estimation working group.

Year	Catch
2001	57
2002	86
2003	27
2004	57
2005	83
2006	102
2007	104
2008	120
2009	166
2010	89
average	89

*(This page intentionally left blank)*