17. Bering Sea and Aleutian Islands Skates

Olav Ormseth, Beth Matta, and Jerry Hoff
NMFS Alaska Fisheries Science Center, Seattle, WA

Executive Summary

Summary of Major Changes

In the 2007 assessment, we introduced a new age-structured model for the Alaska skate (*Bathyraja parmifera*). This model was not adopted by the SSC due to several concerns, primarily regarding model fits to survey and length-at-age data. Many of these problems have largely been resolved, although the model continues to underestimate length-at-age of older skates. The base model has also changed from the version presented in September 2008. An alternative model is presented and used to explore lack of fit to the length-at-age data.

Changes in the input data:

- Total catch (t) for the BSAI skate assemblage is updated with 2007 and partial 2008 data.
- Biomass estimates from the 2008 EBS shelf and slope surveys are incorporated for all species.
- Catch and survey length composition data were updated, and 4-cm length bins were used instead of 5-cm length bins.
- Length-at-age data from the 2007 EBS shelf survey are included.
- Sample size (N) for length composition data was based on the number of hauls sampled, rather than the total number of length measurements.

Changes in assessment methodology:

Last year, we recommended splitting the BSAI skates assemblage into two categories, 'Alaska skate' (*Bathyraja parmifera*) and 'Other Skates'. The goal of these separate management categories is to provide increased protection to rare and endemic skate species in the Other Skates group. In the current assessment we provide management recommendations based on Tier 3 criteria and an age-structured model for Alaska skates. As alternatives we present Tier 5 calculations for the Alaska skate using two different values of M: the accepted aggregate skate M for the last several years (0.10) and an estimate specific to the Alaska skate (0.13). We continue to apply the Tier 5 approach using the aggregate M of 0.10 to generate recommendations for the Other Skates group.

The following specific changes were made to the age-structured model for Alaska skates:

- The steepness of the Beverton-Holt stock-recruit relationship (h) was fixed at 1.0.
- The standard deviation of log recruitment (σ_R) was fixed at 0.4.
- Selectivity at age was modeled as a logistic function.
- Independent estimates of survey selectivity have been incorporated into the Alaska skate model.
- Survey catchability (Q) was fixed at 1.0 and a logistic function for survey length selectivity was fixed so that the selectivity matched the results of an independent analysis of skate capture probability.

Summary of results

We recommend applying Tier 3 criteria to the Alaska skate and Tier 5 criteria to the Other Skates complex, resulting in the following specifications for each management group. Tier 5 specifications for the Alaska skate are also provided for comparison.

	Recommendations		Alternatives	
	Alaska skate	**Other Skates**	Alaska skate	Alaska skate
M	**0.13**	**0.10**	0.10	0.13
Tier	**3**	**5**	5	5
B100%	**186,142 t**	**N/A**	N/A	N/A
B35%	**65,149 t**	**N/A**	N/A	N/A
B40%	**74,457 t**	**N/A**	N/A	N/A
2009				
proj. age 0+/avg. biomass	**551,439**	**82,206**	406,492	406,492
proj. spawning biomass	**108,093**	**N/A**	N/A	N/A
F_{OFL}	**0.10**	**0.10**	0.10	0.13
Max F_{ABC}	**0.087**	**0.075**	0.075	0.0975
recommended F_{ABC}	**0.087**	**0.075**	0.075	0.0975
OFL	**30,077**	**8,221**	40,649	52,844
Max ABC	**25,854**	**6,165**	30,487	39,633
recommended ABC	**25,854**	**6,165**	30,487	39,633
2010				
proj. age 0+/avg. biomass	**549,000**	**82,206**	406,492	406,492
proj. spawning biomass	**108,089**	**N/A**	N/A	N/A
F_{OFL}	**0.10**	**0.10**	0.10	0.13
Max F_{ABC}	**0.087**	**0.075**	0.075	0.0975
recommended F_{ABC}	**0.087**	**0.075**	0.075	0.0975
OFL	**30,009**	**8,221**	40,649	52,844
Max ABC	**25,796**	**6,165**	30,487	39,633
recommended ABC	**25,796**	**6,165**	30,487	39,633

In the event the SSC chooses Tier 5 criteria for the Alaska skate, we strongly recommend using the more conservative estimate of *M*=0.10 to estimate ABC and OFL. Increasing *M* from 0.10 to 0.13 results in a substantial increase (33%, or approximately 10,000 t) from last year's ABC under Tier 5 criteria. Because skates are managed with the rest of the Other Species complex under a single TAC, such an increase would reduce protection to all the species within this complex, and could encourage the development of directed fisheries.

Responses to SSC Comments

SSC comments specific to the BSAI Skates assessment:

The SSC had several major comments and some minor ones at the October 2008 meeting. The minor comments have all been dealt with, and major comments are addressed below and in the section on Tier 3 assessment.

1) The fit to the size-at-age data has improved but remains biased: the LVB model tends to consistently overestimate length-at-age of younger fish and underestimate length-at-age of older fish (Fig. 35), probably due to limitations of the assumed growth model. This bias appears to result in an overestimation of the number of skates in intermediate size classes and an underestimation of the number of skates in larger size classes (Fig. A13). Because skates mature at relatively large sizes (Fig. A10), underestimating the abundance of large skates may greatly underestimate spawning biomass. It is our understanding that the new version of SS2 can accommodate more flexible growth models and we encourage the authors to fit one of these more flexible models to improve the fit to size-at-age data. For some elasmobranchs, growth rate shifts at or near size of maturity, and models (e.g., two-stage von Bertalanffy) have been developed to handle such situations. In addition, we encourage the author to explore and document the sensitivity of the model to the assumption that L_1 is fixed at 22 cm, given the large uncertainty (CV) of this parameter (Table A6).

Response: A concerted effort was made to transition to the new version of Stock Synthesis (SS3). SS3 contains an advanced growth function based on the generalized 4-parameter Schnute model. However, we discovered that the use of this growth model in SS3 is currently limited in that the Richards coefficient must have a positive value (R. Methot, pers. comm.). Independent analysis of skate growth suggested that models using either a negative coefficient or a coefficient of 0 (which approximates the Gompertz growth model) provided the best fits to skate growth. Additional capabilities are planned for future versions of SS3 and we hope to use these when they become available. For 2008, we continue to use SS2. The addition of length-at-age data from 2007 and changes to model specification provided better fits to the length-at-age data, but the model does continue to underestimate length-at-age for older skates. A discussion of this issue, including an alternative model and possible explanations for the lack of a better fit, are presented in the model evaluation section. Briefly, we feel that underestimation of skate growth, while not desirable, is at least precautionary in that it possibly underestimates spawning biomass.

2) The authors present output from a single model that was based on a number of assumptions that are difficult to evaluate. In particular, the authors make a strong assumption about the limited level of recruitment variability (fixed at $\sigma_R = 0.3$). The authors argue that skate recruitment should display low variability because skates are equilibrium strategists. However, recruitment is effectively estimated at age-4 by the model and variability in egg deposition and in the survival between egg deposition and emergence could easily lead to considerable variability in age-4 recruitment. The authors chose $\sigma_R = 0.3$ "…on the basis of improved model fits", but differences in model fits were not presented (last year's model assumed $\sigma_R = 0.1$). Therefore, the SSC recommends that the authors document the sensitivity of the model to the specification of σ_R or provide a stronger rationale for their choice. For example, alternative models with different levels of σ_R or a likelihood profile for σ_R could be presented.

Response: A likelihood-profile analysis of σ_R was performed and is discussed in the model evaluation section. Based on this analysis, σ_R was fixed at 0.4.

3) The authors assume that egg case development takes 3.6 years based on a study by Hoff (2006). The SSC requests that the authors include a brief description of the available evidence for this determination, including some discussion of the reliability of skate aging data and of the methods used to determine development times and age determinations.

Response: The existing sections regarding embryo development have been expanded to provide more complete information and two figures from Hoff (2007) have been included.

4) There should be some discussion on the sensitivity of model results to the assumptions that were made regarding selectivity parameters. The SSC notes that many of these parameters were arbitrarily bounded and parameter estimates were often near their specified bounds (e.g., p1 for longline length selectivity; and, p3, p4 and p6 for trawl length selectivities, Table A6).

Response: The bounds used for the selectivity parameters were, with the exception of p1, the default bounds suggested for the parameters of the double-normal function in SS2. It is our understanding that these bounds cover the entire range of possible values for the double-normal parameters. Parameters for p1 were taken from an SS2 assessment of longnose skates in the Pacific Northwest. In the model presented here, all estimated selectivity parameters were well within the specified bounds. The text has been expanded and clarified where appropriate to address these issues.

SSC comments on assessments in general:
There were no SSC comments on assessments in general.

General Introduction

Description, scientific names, and general distribution
Skates (family Rajidae) are cartilaginous fishes which are related to sharks. They are dorso-ventrally depressed animals with large pectoral "wings" attached to the sides of the head, and long, narrow whiplike tails (Fig. 1). At least 15 species of skates in three genera, *Raja*, *Bathyraja*, and *Amblyraja,* are distributed throughout the eastern North Pacific and are common from shallow inshore waters to very deep benthic habitats (Eschmeyer et al. 1983, Stevenson et al. 2006). Table 1 lists the species found in Alaskan waters, with their depth distributions and selected life history characteristics (which are outlined in more detail below).

The species within the skate assemblage occupy different habitats and regions within the BSAI FMP area (Fig. 2). In this assessment, we distinguish three habitat areas: the EBS shelf (< 200 m depth), the EBS slope (> 200 m depth), and the Aleutian Islands (AI) region (all depths) (Fig. 3). Within the Eastern Bering Sea (EBS), the skate species composition varies by depth, and species diversity is generally greatest on the upper continental slope at 250 to 500 m depth (Fig. 4; Stevenson et al. 2006). The EBS shelf skate complex is dominated by a single species, the Alaska skate (*Bathyraja parmifera*) (Table 2 & Fig. 3). The Alaska skate is distributed throughout the EBS shelf habitat area (Fig. 5), most commonly at depths of 50 to 200 m (Stevenson 2004), and has accounted for between 91% and 97% of aggregate skate biomass estimates since species identification became reliable in 1999. The Bering or sandpaper skate (*B. interrupta*) is the next most common species on the EBS shelf, and is distributed on the outer continental shelf (Fig. 6).

While skate biomass is much higher on the EBS shelf than on the slope, skate diversity is substantially greater on the EBS slope (Fig. 3). The dominant species on the EBS slope is the Aleutian skate (*B. aleutica*) (Table 2 & Fig. 3). A number of other species are found on the EBS slope in significant

numbers, including the Alaska skate, Commander skate (*B. lindbergi*), whiteblotched skate (*B. maculata*), whitebrow skate (*B. minispinosa*), roughtail skate (*B. trachura*), and mud skate (*B. taranetzi*) (Table 2). Two rare species, the deepsea skate (*B. abyssicola*) and roughshoulder skate (*Amblyraja badia*), have only recently been reported from EBS slope bottom trawl surveys (Stevenson and Orr 2005). The Okhotsk skate (*B. violacea*) is also occasionally found on the EBS slope.

The skate complex in the AI is quite distinct from the EBS shelf and slope complexes, with different species dominating the biomass, as well as at least one endemic species, the recently described butterfly skate, *Bathyraja mariposa* (Stevenson et al. 2004). In the AI, the most abundant species is the whiteblotched skate, *B. maculata* (Table 2 & Fig. 3). The whiteblotched skate is found primarily in the eastern and far western Aleutian Islands (Fig. 7). Aleutian and Alaska skates are also common in the AI. The mud skate (*B. taranetzi*) is relatively common in the AI but represents a lower proportion of total biomass because of its smaller body size. We note that the common species formerly known as the Alaska skate in the western Aleutians looks very different from the Alaska skate found on the EBS shelf (Fig. 8). The Aleutian Islands type or "leopard skate" (*Bathyraja* sp. cf. *parmifera*) has been confirmed to be a separate species (J. Orr pers. comm.).

Management units

In the North Pacific, skate species are part of the "Other species" management category within the Bering Sea Aleutian Islands (BSAI) Fishery Management Plan (FMP). Skate catch is reported as "Other" in aggregate with the catch of sharks, sculpins, and octopus. Because catch is officially reported within the Other species complex, estimates of skate catch must be made independently (see Bycatch and discards, below).

In the BSAI, catch of Other species is limited by a Total Allowable Catch (TAC) which is based on an Allowable Biological Catch (ABC) estimated by the NPFMC Scientific and Statistical Committee (SSC). Currently skates are taken only as bycatch in fisheries directed at target species in the BSAI, so future catches of skates are more dependent on the distribution and limitations placed on target fisheries than on any harvest level established for this category. An FMP amendment was initiated by the NPFMC in 1999 to remove both skates and sharks from the Other species category to increase the level of management attention and control for these potentially vulnerable species groups; this action is still in the process of revision and review. In response to a developing fishery in the GOA, the GOA FMP was amended to remove skates from the Other species category. FMP amendments have been proposed to split the Other species category into component groups in both the BSAI and GOA, and this assessment is written as a stand-alone skate assessment in support of this effort to improve Other species management.

Life history and stock structure (general)

Skate life cycles are similar to sharks, with relatively low fecundity, slow growth to large body sizes, and dependence of population stability on high survival rates of a few well developed offspring (Moyle and Cech 1996). Sharks and skates in general have been classified as "equilibrium" life history strategists (Winemiller and Rose 1992), with very low intrinsic rates of population increase implying that sustainable harvest is possible only at very low to moderate fishing mortality rates (King and McFarlane 2003). Within this general equilibrium life history strategy, there can still be considerable variability between skate species in terms of life history parameters (Walker and Hislop 1998). While smaller sized species have been observed to be somewhat more productive, large skate species with late maturation (11+ years) are most vulnerable to heavy fishing pressure (Walker and Hislop 1998; Frisk et al. 2001; Frisk et al. 2002). The most extreme cases of overexploitation have been reported in the North Atlantic, where the "common" skate *Dipturus batis* has been extirpated from the Irish Sea (Brander 1981) and much of the North Sea (Walker and Hislop 1998), and the barndoor skate *Dipturus laevis* disappeared from much of its range off New England (Casey and Myers 1998). The relative difference in life history

traits between smaller and larger skate species has led to apparent population stability for the aggregated "skate" group in many areas where fisheries occur, and this combined with the common practice of managing skate species within aggregate complexes has masked the decline of individual skate species in European fisheries (Dulvy et al. 2000). A similar situation has occurred off the northeast coast of the United States, where skates are managed as a complex and are the subject of skate wing and lobster bait target fisheries; skates are also taken incidentally in other fisheries (NEFSC 2007). Aggregate skate biomass was relatively stable in the 1970s, but has fluctuated since the early 1980s, with apparent shifts in the relative abundance of individual species (NEFSC 2007). Declines in barndoor skate abundance were concurrent with an increase in the biomass of skates as a group (Sosebee 1998). While barndoor skate biomass is now above minimum threshold levels, winter skates (*Leucoraja ocellata*) and thorny skates (*Amblyraja radiata*) have become overfished, and smooth skates (*Malacoraja senta*) and little skates (*Leucoraja erinacea*) are in danger of becoming overfished according to the New England Fishery Management Council's definitions, requiring immediate action to reduce mortality and initiate rebuilding of overfished stocks (NEFSC 2007 and http://www.nefmc.org/skates/index.html).

Several recent studies have explored the effects of fishing on a variety of skate species in order to determine which life history traits might indicate the most effective management measures for each species. While full age-structured modeling is difficult for many relatively information-poor species, Leslie matrix models parameterized with fecundity, age/size at maturity, and longevity have been applied to identify the life stages most important to population stability. Major life stages include the egg stage, the juvenile stage, and the adult stage (summarized here based on Frisk et al. 2002). All skate species are oviparous (egg-laying), investing considerably more energy per large, well-protected embryo than most commercially exploited teleost groundfish. The large, leathery egg cases incubate for extended periods (several months to over a year) in benthic habitats, exposed to some level of predation and physical damage, until the fully formed juveniles hatch. The juvenile stage lasts from hatching through maturity, several years to over a decade depending on the species. The reproductive adult stage may last several more years to decades depending on the species.

Age and size at maturity and adult size/longevity appear to be more important predictors of resilience to fishing pressure than fecundity or egg survival in the skate populations studied to date. Frisk et al. (2002) estimated that although annual fecundity per female may be on the order of less than 50 eggs per year (extremely low compared with teleost groundfish), there is relatively high survival of eggs due to the high parental investment, and therefore egg survival did not appear to be the most important life history stage contributing to population stability under fishing pressure. Juvenile survival appears to be most important to population stability for most North Sea species studied (Walker and Hislop 1998) and for the small and intermediate sized skates from New England (Frisk et al. 2002). For the large and long-lived barndoor skate, adult survival was the most important contributor to population stability (Frisk et al. 2002). Comparisons of length frequencies for surveyed North Sea skates from the mid and late 1900s led Walker and Hislop (1998, p. 399) to the conclusion that after years of very heavy exploitation "all the breeding females, and a large majority of the juveniles, of *Dipturus batis*, *Leucoraja fullonica* and *R. clavata* have disappeared, whilst the other species have lost only the very largest individuals." Although juvenile and adult survival may have different importance by skate species, all studies found that one metric, adult size, reflected overall sensitivity to fishing. After modeling several New England skate populations, Frisk et al. (2002, p. 582) found "a significant negative, nonlinear association between species total allowable mortality, and species maximum size." This may be an oversimplification of the potential response of skate populations to fishing; in reality it is the interaction of natural mortality, age at maturity, and the selectivity of fisheries which determines a given species' sensitivity to fishing and therefore the total allowable mortality (ABC).

Life history and stock structure (Alaska-specific)

Known life history parameters of Alaskan skate species are presented in Table 1. Zeiner and Wolf (1993) determined age at maturity and maximum age for big skates (*Raja binoculata*) and longnose skates (*R. rhina*) from Monterey Bay, CA. The maximum age of CA big skates was 11-12 years, with maturity occurring at 8-11 years; estimates of maximum age for CA longnose skates were 12-13 years, with maturity occurring at 6-9 years. McFarlane and King (2006) recently completed a study of age, growth, and maturation of big and longnose skates in the waters off British Columbia (BC), finding maximum ages of 26 years for both species, much older than the estimates of Zeiner and Wolf. Age at 50% maturity occurs at 6-8 years in BC big skates, and at 7-10 years in BC longnose skates. However, these parameter values may not apply to Alaskan stocks. The AFSC Age and Growth Program has recently reported a maximum observed age of 25 years for the longnose skate in the GOA, significantly higher than that found by Zeiner and Wolf but close to that observed by McFarlane and King (Gburski et al. 2007). In the same study, the maximum observed age for GOA big skates was 15 years, closer to Zeiner and Wolf's results for California big skates. The life histories of these two species are reported in more detail in the GOA skate SAFE (Ormseth and Matta 2007).

Considerable research has been directed at skates in the Bering Sea within the past few years. Graduate students at the University of Washington and California State University (Moss Landing Marine Laboratories) have begun or completed projects detailing aspects of life history and population dynamics of several Bering Sea species. A comprehensive study on the age, growth, and reproductive biology of the Alaska skate, the most common skate species on the eastern Bering Sea shelf, was recently completed (Matta 2006). Life history aspects examined in this study include estimates of maximum age, instantaneous rate of natural mortality (M), length and age at maturity, growth parameters, annual fecundity, and seasonal reproductive timing. Age and size at 50% maturity were 9 years and 92 cm TL for males and 10 years and 93 cm TL for females (Table 1). Von Bertalanffy growth parameters were estimated for males (L_∞ = 126.29 cm TL, k = 0.120 year^{-1}, t_0 = -1.39 year) and females (L_∞ = 144.62 cm TL, k = 0.087 year^{-1}, t_0 = -1.75 year), although length-at-age data were fit slightly better by a Gompertz growth function for both sexes. Based on seasonal reproductive data, including ova diameter, gonadosomatic index (GSI), and the presence of egg cases, the Alaska skate appears to be reproductively active throughout the year. A reproductive resting phase (e.g. 'spent' gonads) was never observed in either large males or females, and females containing egg cases were encountered during each month of collection. Annual fecundity was estimated to average 21 to 37 eggs per year, based on the relationship between annual reproductive effort and natural mortality (Gunderson 1997). While the fecundity estimate needs to be validated using direct methods, fecundity is still likely to be low for the Alaska skate, as is typical for most elasmobranchs.

Hoff (2007) recently completed a dissertation examining skate reproduction and skate nursery habitat of the Alaska skate and the Aleutian skate from the eastern Bering Sea. The relationships between successful skate reproduction and selected nursery grounds were examined. Vulnerability sources, reproductive cycles, habitat selection criteria, and physical factors controlling reproduction were addressed. To date, six nursery sites for three different skate species have been described in the eastern Bering Sea (Fig. 9), and there is ample evidence that additional nursery areas exist. All sites are located along the shelf-slope interface in approximately 140-360 m of water. Two sites, those of the Alaska and Aleutian skates, have been studied in detail through seasonal monitoring. An index location at each nursery site was re-sampled approximately once every 60 days from June 2004 through July 2005 for a total of eight sampling periods. During each sampling period data on mortality, reproductive cycles, embryo developmental, species utilization and adult reproductive states were examined.

The Alaska skate nursery in Bering Canyon is located in 149 meters of water near the shelf-slope interface in a highly productive area of the eastern Bering Sea. The nursery is small in area (< 2 nautical

miles), persistent, and highly productive. Density estimates from trawling showed the most active part of the nursery contained >100,000 eggs/km^2. Two peak reproductive periods during summer and winter were evident in the Alaska skate nursery. During each active period the nursery showed high densities of mature reproductive adults and high numbers of newly deposited egg cases. Although there are peak reproductive periods at any single sampling time, the nursery contained embryos in all stages of development, and specific cohorts were easily discernable from frequency stage monitoring. Cohort analysis based on embryo lengths measured at an Alaska skate nursery site in the EBS suggested that the Alaska skate has an eggcase development time of over 3 years, possibly due to the cold ocean temperatures in the EBS (Fig. 10; Hoff 2007). Captive studies are currently underway at the Alaska Sealife Center (Seward, AK) to validate this finding, but the field observations are consistent with development times observed in other skate species (Fig. 11; Hoff 2007). For example, thorny skate (*Raja radiata*) embryos spend approximately 2.5 years in the eggcase development stage at warmer temperatures than those found in the EBS (Berestovskii 1994 in Hoff 2007).

The Oregon triton *Fusitriton oregonensis* was the most likely predator on newly deposited egg cases and mortality rate was estimated at 3.64% (Hoff 2007). After hatching, young skates were vulnerable to predation by Pacific cod, *Gadus macrocephalus* and Pacific halibut, *Hippoglossus stenolepis*. Predation by these two large fish species peaked during the summer and winter periods and was highly correlated with hatching events. The Alaska skate nursery site was occupied by mature male and female skates throughout the year, with juvenile and newly hatched individuals extremely rare. Evidence suggests that newly hatched skates quickly move out of the nursery site and immature skates are infrequent visitors to nursery sites. The nursery is located in a highly fished area and is vulnerable to disturbances due to continuous use of the nursery grounds by skates throughout the year. Some degree of intra-species habitat partitioning is evident and is being examined for the Alaska skate throughout the eastern Bering Sea shelf environment.

Researchers at the Pacific Shark Research Center (PSRC), Moss Landing Marine Laboratories (MLML) are currently conducting investigations into aspects of the age, growth, reproduction, demography, and diet of several Alaskan skates. In cooperation with the Alaska Department of Fish and Game and the AFSC, they have examined more than 5,000 specimens comprising 13 species, including Aleutian skate, Commander skate, whiteblotched skate, whitebrow skate, Alaska skate, roughtail skate, Bering skate, and mud skate (Ebert, 2005). Currently, four graduate students are working towards their Masters degrees with thesis projects on Alaskan skate species. In addition, two other students, Chante Davis (2006) and Heather Robinson (2006), have recently completed their respective thesis research on two skate species (roughtail skate and longnose skate) that occur in Alaskan waters. Although their studies were conducted outside of Alaskan waters, their findings represent new and original information on the life history of these two skate species.

Age determination and validation studies are currently ongoing at the PSRC to obtain essential information on the age at maturity, growth rates and longevity of seven Alaskan skate species: Aleutian skate, Commander skate, whiteblotched skate, whitebrow skate, roughtail skate, Bering skate, and mud skate. Theoretical longevity and indirect estimates of natural morality will be calculated from the resulting growth parameters. Additionally, the suitability of caudal thorns as an alternative ageing structure is being investigated, potentially providing a valuable, non-lethal ageing technique for this group. Preliminary estimates of maximum ages for Aleutian and Bering skates are 17 and 13 years, respectively (Ebert et al. 2007). Age validation remains to be completed for these species (D. Ebert, PSRC, pers. comm.). Additional age and growth studies are currently being conducted by Jasmine Fry (mud skate), and Shaara Ainsley (whitebrow skate) for their thesis research.

Reproductive studies are also currently ongoing at the PSRC to obtain information on the size at maturity, seasonality, and fecundity of several Alaskan skate species. The reproductive biology of the Aleutian

skate, Bering skate, big skate, and longnose skate has been investigated as part of a NPRB funded study to assess life history characteristics of Alaskan skate species (Ebert et al. 2007). Median length at maturity (cm TL) was estimated to be 124.4 for the Aleutian skate, 70.2 for the Bering skate, 148.6 for the big skate, and 113.1 for the longnose skate (Ebert et al. 2007). Reproductive studies are also being conducted on mud and whitebrow skates by graduate students affiliated with the PSRC.

The PSRC has also conducted demographic analyses to improve understanding of the population dynamics and vulnerability of these species to fisheries exploitation. Preliminary estimates of annual population growth rates are 25% for the Aleutian skate, 36% for the Bering skate, 33% for the big skate, and 20% for the longnose skate (Ebert et al. 2007). Other demographic parameters have also been estimated for these species (Ebert et al. 2007). Information generated from this project will be incorporated into a life history data matrix (LHDM) developed by the PSRC for eastern North Pacific chondrichthyans; the most recent version of the LHDM is currently available via the worldwide web (http://psrc.mlml.calstate.edu/).

Fishery

Directed fishery

In the BSAI, there is no directed fishery for skates at present; however, skates support directed fisheries in other parts of the world (Agnew et al. 1999, NE stock assessment 1999, Martin and Zorzi 1993). A directed skate fishery developed in the Gulf of Alaska in 2003 (Gaichas et al. 2003). There has been interest in developing markets for skates in Alaska (J. Bang and S. Bolton, Alaska Fishworks Inc., 11 March 2002 personal communication), and the resource was economically valuable to the GOA participants in 2003, although the price apparently dropped in 2004. Nevertheless, we should expect continued interest in skates as a potential future target fishery in the BSAI as well as in the GOA.

Bycatch and discards

Skate catch in the BSAI is officially reported as “Other” in aggregate with the catch of sharks, sculpins, and octopus, and thus estimates of skate catch must be made independently for each year using observer data, shoreside processor landings data, and processor weekly production report data. In 2003 the Alaska Regional Office (AKRO) converted to the Catch Accounting System (CAS), an improvement over the previous “Blend” system. However, at present the CAS is only capable of reporting aggregate skate catch in the BSAI; species composition of the catch can only be inferred from the observed portion of the catch or from survey species composition (see Data section below). The CAS data are continuously updated and checked for errors by AKRO; the CAS estimates reported here represent the best and most accurate data available.

Skates constitute the bulk of the Other species FMP category catches, accounting for between 51% and 75% of the estimated totals in 1992-2008 (Table 3). While skates are caught in almost all fisheries and areas of the Bering Sea shelf, most of the skate bycatch is in the hook and line fishery for Pacific cod, with trawl fisheries for pollock, rock sole, flathead sole, and yellowfin sole also catch significant amounts (Tables 4 & 5). In this assessment, "bycatch" is interpreted as incidental or unintentional catch regardless of the disposition of catch – it can be either retained or discarded. We do not use the Magnuson Act definition of "bycatch," which always implies discard. When caught as bycatch, skates may be discarded (and may survive depending upon catch handling practices) although skates caught incidentally are sometimes retained and processed. Due to incomplete observer coverage, it is difficult to determine how many skates are actually retained. However, between 24% and 39% of the total observed skate catch was retained during the years 2003-2006 (Table 6). More skates were retained in the EBS than the AI, and it appears that species that grow to a larger maximum size (>100 cm TL) are more likely to be retained than smaller-bodied species. For example, while the Aleutian skate, a large-bodied species, made up a relatively small portion of the observed skate catch in 2005 (approximately 2%), 31% of the Aleutian

skates caught were retained. However, Bering skates (a small-bodied species less than 100 cm TL) were retained less frequently (10% in 2005). Larger percentages of Alaska skates and *Raja* species (big and longnose skates) are also retained; all three are relatively large-bodied skates.

Historically, skates were almost always recorded as "skate unidentified", with very few exceptions between 1990 and 2002. However, due to improvements in species identification by fishery observers initiated by Dr. Duane Stevenson (AFSC) within the Observer program in 2003, we can estimate the species composition of observed skate catches 2004-2006 (Fig. 12). Recent observer data indicates that only about 50% of skate catch is not identified to the species level. This is largely because most skates are caught in longline fisheries, and if the animal drops off the longline as unretained incidental catch, it cannot be identified to species by the observer (approximately 80% of longline-caught skates are unidentified, and longline catch accounts for the majority of observed skate catch).

In 2005, observers were encouraged to identify skates dropped off longlines to genus, which can be done without retaining the skate; hence in 2005 more than half of the unidentified skates were at least assigned to the genus *Bathyraja*. Of the identified skates, the majority (90%) were Alaska skates, as would be expected by their dominance in terms of overall skate biomass in the BSAI. The next most commonly identified species BSAI-wide was Aleutian skate, at 6.6% of identified catch, followed by Bering skates at 4.3 %, big skates at 3.6%, and whiteblotched at approximately 1.3% across the BSAI. It should be noted that the observed skate catch composition may not reflect the true catch composition, possibly due to selective retention of larger species or to a higher likelihood of identifying distinctive species. However, when viewed by area (EBS vs. AI), it is clear that the majority of identified Aleutian and whiteblotched skates are caught in AI fisheries, and that the species composition of the observed catch in the AI is very different from the EBS (Fig. 12).

Reporting areas encompassing the EBS outer shelf and upper continental slope experienced high catch rates during 2003-2006 (Fig. 13). Longline fisheries targeting Pacific cod take much of the incidental skate catch, and they tend to operate on the outer EBS shelf and slope where skate species diversity is high and where Aleutian skates are more prevalent than Alaska skates. Therefore it is possible that the species composition of the catch is not in proportion to the overall species composition (from survey data) across the BSAI. However, depth analysis of the observed catch demonstrates that most of the skate catch occurs <200m (98%). More work is needed to determine the actual species composition of the catch.

ALASKA SKATE – Tier 3 assessment

Overview

The model presented here begins in 1992. In the 2007 assessment, we included an alternative model starting in 1958 that included historical catch data and survey biomass estimates from 1982. The alternative model was eliminated from the 2008 assessment for several reasons:

1) Catch data for skates prior to 1992 are highly uncertain. Skate catches were not reported and it was necessary to assume that catch of all skates was accurately recorded as part of an aggregate non-target species catch. This also necessitated various assumptions about the proportion of skates and *B. parmifera* in the non-target category (resulting in the low- and high-catch scenarios used in the 2007 assessment).
2) The nature of the fisheries that might have caught skates prior to 1992 is also highly uncertain and required assumptions regarding the gears employed in various target fisheries. It is unlikely that the fishery selectivity estimated in the model (which is based on one year- 2007- of length composition data) is applicable to the foreign and joint-venture fisheries that were conducted prior to the 1990s.
3) The EBS shelf survey biomass estimates for skates exhibit a dramatic increase in the 1980s. It is unclear at this time whether this increase was due to enhanced recruitment and survival of skates, a shift in skate distribution, or recovery from previous intense exploitation. In addition, it is unclear what proportion of this biomass increase was due to *B. parmifera*. Modeling the dynamics of the Alaska skate population using this information has the potential to result in serious errors in model specification.
4) Survey length composition data for *B. parmifera* are only available from 2000. To model the dynamics of the population more than 30 years prior to this would likely result in fitting the model to noise.

For these reasons, the population is modeled during the "modern era" for skates in the BSAI, where the biomass has remained relatively stable and available data are substantially more complete and reliable.

This assessment model resembles teleost groundfish models in many ways, but we made some changes to incorporate life history features unique to elasmobranchs. As previously discussed, all skate species have an extended embryonic period during which they develop within protective eggcases on the seafloor. Alaska skates do not appear to form visible annual growth marks in their vertebrae during embryonic development. However, cohort analysis based on embryo lengths measured at an Alaska skate nursery site in the EBS suggested that the Alaska skate has an eggcase development time of approximately 3.6 years, possibly due to the cold ocean temperatures in the EBS (Hoff 2007; Fig. 10). Captive studies are currently underway at the Alaska Sealife Center (Seward, AK) to validate this finding, but the field observations are consistent with development times observed in other skate species (Fig. 11; Hoff 2007). For example, thorny skate (*Raja radiata*) embryos spend approximately 2.5 years in the eggcase development stage at warmer temperatures than those found in the EBS (Berestovskii 1994 in Hoff 2007). Incorporating this information in the model is complicated by the possibility that embryo development times may be temperature-dependent (G. Hoff, pers. comm.). In addition, the timing of *B. parmifera* reproduction is uncertain. While most females appear to deposit eggcases during the summer, with emergence of young skates occurring during the winter, some level of skate reproduction seems to occur year-round. We assigned the first three age classes of Alaska skates (0-2) to an embryonic period where growth differed from older age classes and individuals were not available to either the fishery or survey. Thus, free-swimming skates in their first year were considered to be 3½ years old. In addition, we adjusted parameters of the length model and age selectivity to accommodate the developmental delay and the uncertainty in its duration. This approach allowed us to more accurately model skate population

dynamics and ensured that characteristics of the spawning population would correspond to the appropriate year class. In addition, we considered the equilibrium life history strategy in specifying recruitment parameters and evaluating our model results.

We present a base model that we determined to provide the best description of Alaska skate population dynamics given the data and the limitations of the modeling software. We also present an alternative model to explore problems with the model fit to length-at-age data. Aspects of this model are described in the section on model evaluation.

Data

Survey biomass

Three bottom trawl surveys are conducted in the BSAI region: EBS shelf, EBS slope, and the Aleutian Islands. Because the Alaska skate population is concentrated on the EBS shelf, and the EBS shelf survey provides yearly estimates of biomass, we used biomass data from only the EBS shelf survey in this assessment. Recent (1999-2008) survey information on species composition was used to describe the relative proportion (0.95) of the Alaska skate to all other skate species ("Other Skates") within the EBS shelf area (Table 7 & Fig. 14). Biomass estimates from 1992 through 2008 were utilized in the Alaska skate stock assessment base model. For each survey prior to 1999, total skate biomass estimates were partitioned into Alaska skate and Other Skates based on the average proportion of each group in the 1999-2008 surveys (Table 7). The model employs the standard deviation (s) associated with each estimate, which as calculated using the equation: $\ln(1 + CV)$, where CV is the standard error of the observation divided by the value of the observation (Methot 2007). For the estimates prior to 1999, a value of s was chosen that was intermediate to recent values and a high s observed in 1999 (Table 7).

Survey length composition

Total length (TL) data from the EBS shelf survey were available from 2000-2008 (Table 8). The survey takes length measurements for every skate in each haul. Each haul where skate lengths were taken was treated as an independent sample.

Binning: Discussions with staff from the Resource Assessment and Conservation Engineering (RACE) division at the AFSC during summer 2008 indicated that there may be a slight bias in the length measurements of skates in the EBS shelf survey towards odd-numbered sizes. This is likely due to the design of the length measuring boards, which display the odd sizes along the edge closest to the biologists, and the general difficulty of measuring a disc-shaped animal like a skate. This bias might be important when 5-cm length bins are used, as the bins contain different proportions of odd and even sizes. To ameliorate this problem, for 2008 the length composition bins were changed to a 4-cm width that includes equal numbers of odd and even sizes.

Total catch

Commercial catches of BSAI skates are reported FMP area-wide in aggregate with sculpins, octopus, and squid. Independent estimates of BSAI skate catch from 1992-2008 were made by the Blend system and AKRO CAS as described in the 2007 BSAI skate assessment. For the base model, catches were broken down by habitat area (EBS shelf, EBS slope, and AI) and by fishery gear type from 1992-2008 (Table 9). Total skate catch estimates for the EBS and AI are available since 1997; the average proportion of the skate catch in both of these areas (94% EBS and 6% AI) was assumed to remain constant prior to 1997 in order to reconstruct the area-specific catch. Catch is not estimated separately for the EBS shelf and EBS slope habitat areas by Blend or CAS; therefore a proxy based on fishery observer depth data was developed. The observed total skate catch from 2003-2007 in the EBS was partitioned by depth in order to approximate the proportion of the catch occurring in each of the two EBS habitat areas; catches less than 200 m were considered to occur on the EBS shelf (about 98%) and catches deeper than 200 m were considered to occur on the EBS slope (about 2%).

The average area-specific species compositions from the 1999-2008 bottom trawl surveys (Fig. 14) were utilized to further partition the catch into Alaska skates and Other Skates. Two major fishery gear types with different size selectivities for skates operate in the BSAI management area: trawlers and longliners. (Pot gear also accounts for a minor portion of the skate catch (<0.1%) and was considered negligible for the purposes of this assessment.) The proportion of the catch by each fishery gear type differs by habitat area; for years without gear type data, the average proportion of each gear type from 2003 to 2005 was applied. The results were then totaled to obtain the total Alaska skate catch for each fishery across the entire BSAI management area, which was incorporated into the model (Table 9 and Fig. 15).

Catch length composition

Length data for the Alaska skate were collected as a special project by fishery observers aboard trawl and longline vessels operating in the EBS in 2007. Observers were requested to randomly sample up to 20 skates in one set per week during the study period. Length data were aggregated into 4-cm bins for incorporation into the stock assessment model (Table 10). When these data were updated for the 2008 assessment, it was discovered that the use of incomplete length data in 2007 had produced a bias in the description of the commercial catch. In both fisheries, catches in the third quarter contained a substantially greater proportion of large skates than catches during the first quarter (Fig. 16). As a result, the length compositions used in the 2008 assessments differ markedly from those included in the 2007 assessment.

Length at age

Mean length at age data were obtained from Matta (2006) and from production ageing at the AFSC. Age was determined through examination of annual growth rings which begin to form in vertebral thin sections following hatching from the eggcase. Skate age determination is inherently difficult due to the typically faint appearance of growth zones, and CVs associated with many skate ageing studies tend to be high. However, Matta (2006) was able to corroborate ages generated from two different ageing structures in the Alaska skate, vertebrae and caudal thorns, as well as to verify the annual periodicity of vertebral growth ring formation through marginal increment analysis. Three sample sets were included in the model; one from the 2003 EBS shelf survey (n=182; Fig. 17), one from the 2005 longline fishery (n=208; Fig. 18), and one from the 2007 EBS shelf survey (n=243).

Weight at length

Parameters from the allometric length-weight relationship ($W = aTL^b$, where W is weight in kg and TL is total length in cm) were obtained from Matta (2006) for the Alaska skate. For sexes combined, a was estimated as $4.01*10^{-6}$ and b was estimated as 3.149 (n = 526; Fig. 19).

Analytical approach

Model structure

The Stock Synthesis 2 (SS2) assessment program[1] (Methot 2005, 2007) was used to develop an age-structured population model of Alaska skates. SS2 allows the flexibility to incorporate both age-and size-structured information in the model. In the model described here, natural mortality is the only parameter that is explicitly age-based; selectivity, maturity, and mean body weight are length-based parameters. Length-at-age data and estimates of ageing error are used by SS2 to convert the size-based information into age-specific values that can be used to model the population through time.

[1] NOAA Fisheries Toolbox Version 2.10, 2006. Stock Synthesis 2, Version 2.00g, Richard Methot, Northwest Fisheries Science Center, Seattle, WA. [Internet address: http://nft/nefsc.noaa.gov]

SS2 is comprised of three submodels. A population submodel captures the dynamics of an age-structured population and an observation model specifies likelihood components for comparing model predictions to observed data. A statistical model incorporates those components and others into an objective function that SS2 uses to maximize the overall likelihood by altering the parameters that govern the population dynamics model. SS2 also contains a forecasting routine that specifies fishery management targets and projects the population into the future, but we used an alternative projection model that was designed exclusively for use in Alaska fisheries by Jim Ianelli (AFSC, NMFS). The structure of SS2 is explained in detail elsewhere (Methot 1990, 2005, 2007), and we offer here only a limited explanation of the model structure.

The population dynamics model is depicted schematically in Fig. 20. Briefly, unfished recruitment and *M* determine the age structure of an unfished population. The unfished age structure is then modified by *M* and equilibrium catch to produce an initial age structure. For each subsequent year in the model, individuals are added through recruitment and subtracted through *M* and catch. The expected level of recruitment in each year results from estimates of spawning biomass in the previous year and the parameters of the Beverton-Holt stock-recruit curve. Model estimates of recruitment deviate from the expected level according to the standard deviation of log recruitment (σ_R), which can be fixed or estimated within the model. In all cases, catch is modified by fishery age and length selectivity. For Alaska skates, the observation submodel includes three likelihood components based on model fits to observed data: EBS shelf survey biomass, length compositions from the shelf survey and each of the fisheries, and mean length at age. An additional likelihood component compares the deviations in recruitment to the value of σ_R. The objective function combines these four components to calculate overall likelihood. All likelihood components were weighted equally in the model.

This assessment model included a number of simplifications and assumptions. The entire BSAI was treated as one homogenous area. Because growth and maturity patterns are similar for males and females, we specified only one sex. Spawning was assumed to occur at the midpoint of the year. No informative priors were used. We also assumed that parameters did not vary with season or year and were not influenced by environmental conditions. All parameters used in the base model are listed in Table 11 and described in more detail below.

<u>Parameters estimated independently:</u>
Natural mortality (*M*)
In earlier runs of the model presented to the Plan Team and SSC in September 2008, a fixed value of *M* of 0.12 was used. After adding 2008 survey data and re-specifying parts of the model, an *M* of 0.13 (the same value used in 2007) provided the best fit (see likelihood-profile analysis in the model evaluation section). In 2007, a conservative value of 0.13 was chosen from a set of *M* values estimated using different life history parameters (Matta 2006; Table 12): growth parameters (Alverson and Carney 1975, Pauly 1980, Charnov 1993), longevity (Hoenig 1983), reproductive potential (Rikhter and Efanov 1976, Roff 1986), von Bertlanffy k (Jensen 1996, Gunderson 2003), and age at maturity (Jensen 1996).

Length at maturity
SS2 incorporates female maturity parameters into the model using the following equation:

$$\text{Proportion Mature} = \frac{1}{1+e^{b(L-L_{50})}},$$

where L_{50} is the length at 50% maturity and *b* is a slope parameter. Maturity parameters were obtained from Matta (2006), where b = -0.548 and L_{50} = 93.28 cm TL (Table 11 & Fig. 21). Maturity was

estimated directly from paired length and maturity stage data; maturity stage was easily assessed through macroscopic examination of the reproductive organs.

Ageing error
Each vertebra was aged three independent times by a primary age reader without knowledge of the specimen's biological information. For each true age, the standard deviation of the estimated age was calculated from the three reads of each vertebra and incorporated into the model to account for variability in age determination.

Survey catchability
For November 2008, a major change was made to model specifications of survey catchability and selectivity. Empirical evidence suggests that the capture probability of a combined *Bathyraja* species group in the shelf bottom survey is highly length-dependent with a maximum value of 0.846 for the largest skates (Kotwicki and Weinberg 2005; Fig. 22). In previous runs of the model, we set the catchability equal to a maximum capture probability of 0.836 for the Alaska skate (maximum length = 125 cm). We then adjusted selectivity parameters so that the combination of catchability and selectivity approximated the empirical estimates of size-based capture probability. For November 2008, we took the opposite approach: we assumed a catchability of 1.0 and fixed the survey length selectivity parameters according to parameters of the logistic equation given in Kotwicki and Weinberg (2005). We also chose not to adjust catchability for the segments of the Alaska skate population (AI and EBS slope) that are not observed by the EBS shelf survey. Over 96% of the Alaska skate population is on the shelf, surveys from the other areas are infrequent, and the AI survey has not been conducted since 2006. We felt it was a precautionary measure not to account for the small amount of Alaska skate biomass on the slope and in the AI.

Length selectivity
A logistic selectivity pattern was specified for the EBS shelf survey. Parameters of the logistic function given in Kotwicki and Weinberg (2005) were adapted for the form of the function used in SS2, and both parameters were fixed (Table 11). Fishery length selectivity was governed by a double-normal function defined by six parameters for each fishery or survey, where p1 was the peak or ascending inflection size, p2 was the width of the plateau, p3 was the ascending width, p4 was the descending width, p5 was the selectivity at the first length bin, and p6 was the selectivity at the last length bin. Selectivity parameters are summarized in Table 11. For each fishery, p6 was fixed so that selectivity was asymptotic and all other parameters were estimated within the model. With the exception of p1, all bounds were the default values specified in the SS2 documentation. Bounds for p1 were taken from an SS2 model for longnose skates in the Pacific Northwest (Gertseva et al. 2007).

Age selectivity
The uncertainty surrounding the embryonic development period for the Alaska skate posed some problems in this assessment, and age selectivity was used to partially offset these problems. The best estimate of embryo development times is approximately 3.6 years (Hoff 2007), and the majority of young skates appear to emerge during the winter. Therefore, surveys conducted during the following summer would be catching age-4 skates. A logistic age selectivity function was used for the survey and both fisheries. In all cases, the age at 50% selection was fixed and the width of the selectivity curve was estimated within the model. Age at 50% selectivity was set at age 3.5 for the trawl fishery and age 6 for the longline fishery (based on the lack of earlier ages in the length-at-age data available for the longline fishery). An age of 4 was specified for the trawl survey; a likelihood-profile analysis of this specification is discussed in the model evaluation section.

Parameters estimated conditionally:

Growth parameters

The form of the von Bertalanffy growth equation (LVB) used in SS2 is:

$$L_A = L_\infty + (L_1 - L_\infty)e^{-k(A-A_1)},$$

where L_A is the mean length at age A, A_1 is a reference age near the youngest age well represented in the data, L_1 is the mean length at age A_1, k is the von Bertalanffy growth coefficient, and L_∞ is the mean asymptotic length, calculated from the equation:

$$L_\infty = L_1 + \frac{L_2 - L_1}{1 - e^{-k(A_2 - A_1)}},$$

where A_2 is a reference age near the oldest age well represented in the data, and L_2 is the mean length at age A_2. The reference ages A_1 and A_2 were set to 3.5 and 20 years, respectively, because these ages were frequently observed and represented nearly the entire age range of the Alaska skate. L_1, L_2 and k were estimated within the model.

Spawner-recruit parameters

A Beverton-Holt function was used to describe the spawner-recruit relationship of the Alaska skate (Table 11). The steepness of this function was fixed at 1.0, which has the effect of producing an average level of recruitment unaffected by the level of spawning biomass (SSB). This value was chosen because there is very little contrast in SSB for the modeling time period and the data are thus uninformative regarding steepness. The unfished level of recruitment (R_0) was freely estimated within the model. Recruitment deviations were included in the model, and the standard deviation of log recruitment (σ^R) was fixed at 0.4. A likelihood-profile analysis and discussion of this specification are discussed in the model evaluation section.

Initial fishing mortality

Initial fishing mortality was estimated within the model for each of the two fisheries.

Model Evaluation

Two models are provided for evaluation, a base model and an alternative model:

- Base model: The base model employs the analytical approach and parameters described above. Major changes from the preliminary model presented in September 2008 are the specification of survey catchability and selectivity, and changes to fixed values of M and σ_R.
- Alternative model: An alternative model is presented to explore the lack of fit to the length-at-age data. The only difference in the alternative model relative to the base model regards length selectivity. A double-normal function was specified for both fisheries and the survey. Starting values for the fisheries were identical to the base model; survey starting values were calculated by approximating the survey size composition data using an Excel function supplied with the SS2 software. All length selectivity parameters were estimated within the model.

Model evaluation criteria

Likelihood values are given in Table 13. We evaluated the base model based on the following criteria:

1) Model fit to survey biomass estimates.
2) Model fit to length compositions.
3) Model fit to length-at-age data.
4) Reasonable estimates of fishery length selectivity parameters.

5) Reasonable estimates of unfished recruitment and recruitment variability.
6) Likelihood profile analysis of assumed values for M, age selectivity, and σ^R.

The alternative model is presented to contrast assumptions regarding selectivity and to explore problems in the model fit to length-at-age data, and a full model evaluation was not performed.

Evaluation of the base model

1) The expected survey biomass produced by the model provided a good fit to the observed biomass (Fig. 23). The expected survey biomass is within the confidence interval of all but 3 of the observed biomass estimates. While the biomass estimate dropped considerably in 2008, the model fit reflects the general increase in Alaska skate biomass since 2003.

2) The model provided good fits to the length composition data from the EBS shelf survey (Fig. 24) and both fisheries (Fig. 25). The model is unable to capture the spikes in large skates observed in the 2003 and 2004 surveys, but it does fit the two modes observed in the survey length composition data. The fit to the fishery data is very good.

3) The model fit the observed length-at-age data from the survey (Fig. 26) and longline fishery (Fig. 27) reasonably well, except for older skates. These fits are improved relative to the 2007 assessment. Fig. 28 shows the fit of the population growth estimate (in contrast to the observed estimate) to the three length-at-age datasets.

   The model continues to underestimate length for skates older than 13 years. The lack of a better fit may be partially due to the limitations of the von Bertalanffy growth model employed in SS2. An attempt was made to model Alaska skates using a new version of Stock Synthesis (SS3), which has a more flexible growth model (the generalized Schnute 4-parameter model). However, the SS3 software is still under development and the capabilities necessary for modeling skate growth (specifying the Richards coefficient as either zero or a negative number, as determined through an independent analysis) are not yet available (R. Methot, pers. comm.).

   An alternative explanation is that the length-at-age data do not accurately reflect skate growth. In the survey length compositions, a single length bin (96-99 cm) consistently has the highest proportion of skates throughout the 9-year time series (marked in red in Figs. 24 &25). The magnitude of this length bin proportion declined from 2002 onwards, presumably as members of that size class were removed by natural mortality. The observation that the length-bin position of this size class did not move suggests that for most skates, growth stops when they reach approximately 100 cm in length (approximately 13 years of age). This is approximately the age of maturation, and a cessation of growth after reaching maturity has been hypothesized for some elasmobranch species. A possible contradiction is that the mean length of older skates is higher than 100 cm. However, these may be exceptional cases. If the growth of most skates ceases, growth in the vertebrae that are used for aging likely ceases as well. Thus, there may be 23-year-old skates 100 cm in length may be misidentified as younger skates. Because the collection of skate vertebrae is length-stratified, it may be that skates with extended growth are preferentially selected. This is supported by the observation that sample size for length-at-age drops considerably after 14-15 years of age.

   Additional discussion of the length-at-age data is included in the evaluation of the alternative model. Although the model may underestimate spawning biomass due to the lack of fit of the length-at-age data, the error is in a precautionary direction.

4) Estimates of selectivity parameters (Table 11) and selectivity at length (Fig. 29) for the longline and trawl fisheries were reasonable. Longline fisheries displayed high selectivity for larger skates, which is consistent with the length composition data. This selectivity may be due in part to the emergence of large skates from the nursery grounds during the third quarter of the year, when the longline catch of large skates is particularly high (Fig. 16). The estimate of trawl selectivity also seems reasonable, as the increased selectivity on smaller skates (relative to longline) is likely due to the concentration of trawl fisheries in areas where small skates are less abundant. All estimated values were well within the SS2 default bounds specified in the model.

5) The base model estimate of unfished recruitment was consistent with the amount of spawning biomass and our limited knowledge of skate fecundity. Evaluating recruitment variability is difficult because little is known about recruitment of equilibrium strategists. The estimated levels of recruitment variability (Figs. 30 & 31) were higher than expected but still seem reasonable for this population. See below for a discussion of likelihood-profile analysis of σ_R.

6) To evaluate the estimates of *M*, age at 50% selectivity, and σ_R we created likelihood profiles by varying the fixed values while monitoring the overall likelihood of the model (Fig. A9).

   M: The value of *M* fixed in the model (0.13) had the lowest negative log likelihood (Fig. 32, upper panel).

   Age at 50% selectivity in the survey: A value of 4.5 years provided the best model fit (Fig. 32, lower panel). However, there were two problems with this value: 1) although the model fit to length-at-age data improved using an age at 50% selectivity of 4.5 years, the fit to the survey degraded and resulted in a predicted increase in survey biomass (Fig. 33) which was not reflected in the observed 2008 biomass estimate; 2) a fixed value of 4.5 years for the age at 50% selectivity meant that no skates younger than 4 years were selected by the survey. Because variability in growth and embryo deposition are expected to result in selection of younger as well as older skates, this is an unrealistic description of survey selectivity. Therefore, an age at 50% selectivity of 4 years (which had the second-lowest likelihood value) was specified in the model.

   Standard deviation of log recruitment (σ_R): Because recruitment deviations are included in the overall likelihood and have larger values with increasing σ_R, for this analysis the likelihood components for the survey, length composition, and length-at-age model fits were considered separately (Fig. 34). All of these components decline with increasing σ_R. This is likely because increased σ_R provides greater flexibility for the model to fit the data. However, an informal analysis of other BSAI groundfish (pollock and Pacific cod) with a σ_R of approximately 0.6 and a comparison of observed year-class variation suggest that a value of σ_R less than 0.6 would be appropriate for Alaska skates. The decline in likelihood values for the survey component plateaus at a σ_R of 0.4, and the decline in the length composition values shows the steepest decline from 0.1 to 0.4. Therefore, a σ_R of 0.4 was selected as the most appropriate value for Alaska skates.

Evaluation of the alternative model

The alternative model was not fully evaluated with the above criteria because there were several aspects of the model that were inappropriate for use in Alaska skates. It is included here to explore how a different model specification might address the problem of the lack of fit to the length-at-age data. Throughout the entire process of model specification, this was the only model configuration that substantially improved the fit to the length-at-age data.

1) The alternative model provides better fits to the length composition data and the length-at-age data than the base model (examples given in Fig. 35). The overall likelihood of the alternative model is 118.6, substantially lower than the base model value of 246.8.

2) Freeing all selectivity patterns resulted in dome-shaped selectivity for both fisheries and the survey, with extreme selectivity for skates in the 90-105 cm range for the longline fishery and survey (Fig. 36). Given that we have an independent analysis of skate capture probability in the survey that indicates increasing selectivity for larger skates (Kotwicki and Weinberg 2005), these curves seem unrealistic. There is a possibility that some large skates are in nursery areas during the summer and are unavailable to the survey. However, some level of reproduction occurs year-round, and our understanding of skate movements is too limited to properly assess this phenomenon. In addition, the longline fishery appears to catch large skates in the fall fishery. If a high number of larger skates does exist, we would expect to see these skates captured in the fishery. A tagging study currently underway may provide more insight into seasonal skate movements.

3) The alternative model also predicts a growth curve with a very high L_∞ (Fig. 37), which is inconsistent with the existing data regarding skate growth.

4) The alternative model estimates the 2008 spawning biomass at 463,022 t, a 4-fold increase over the base model estimate.

The results of the alternative model contradict our existing knowledge regarding survey selectivity and skate growth. In addition, the spawning biomass estimate would result in an ABC far in excess of the OFL recommended using the base model or Tier 5 approaches. Given the uncertainty and inconsistency in the alternative model results, using the alternative model would be inappropriate for Alaska skate assessment and management. This is especially so if the pattern of skate growth suggested in the base model evaluation section is true. The base model represents a precautionary approach with results that are consistent with observations of survey selectivity and growth. Therefore, we recommend the base model specification.

Results

Results presented below are from the base model.

Definitions

Biomass is shown as total (age 0+) biomass (metric tons; t) of all Alaska skates in the population, and as female spawning biomass (t). Recruitment is reported as the number (in thousands) of Alaska skates at age 0. As described above, this corresponds to the number of viable embryos deposited in egg cases.

Biomass time series

Time series of total biomass and spawning biomass estimates from 1992-2007 are reported in Table 14 and in Fig. 38, respectively. These estimates suggest that while total skate biomass has been increasing slightly since 2000, spawning biomass has decreased somewhat during the same period.

Recruitment

Time series of age 0 recruitment are reported in Table 14 and Fig. 30, and the relationship between spawning biomass and recruitment is shown in Fig. 31. The model suggests that recruitment has been relatively low in recent years after being above average during the late 1990s and early 2000s.

Exploitation rate

A time series of exploitation (catch/total biomass) is given in Table 15. The exploitation rate appears to have declined slightly over the last several years. This is likely due to a slight decrease in catches and the increase in skate biomass.

Projections and Harvest Alternatives

Reference points and tier assignment

This assessment using the base model provides us with reliable estimates of B_0, $B_{40\%}$, and the fishing mortality rates corresponding to $F_{40\%}$ and $F_{35\%}$. Therefore, management recommendations are made under Tier 3 of the BSAI Groundfish Fishery Management Plan. Using Tier 3, ABC and OFL are set according to the following criteria:

3a) Stock status: $B/B_{40\%} > 1$

$F_{OFL} = F_{35\%}$

$F_{ABC} \leq F_{40\%}$

3b) Stock status: $0.05 < B/B_{40\%} < 1$

$F_{OFL} = F_{35\%}\ H\ (B/B_{40\%} - 0.05) \times 1/0.95$

$F_{ABC} < F_{40\%}\ H\ (B/B_{40\%} - 0.05) \times 1/0.95$

3c) Stock status: $B/B40\% < 0.05$

$F_{OFL} = 0$

$F_{ABC} = 0$

Specification of OFL and maximum allowable ABC

Values for this section, including estimates of equilibrium catch, spawning biomass, and fishing mortality are given in Table 17-16. The 2009 estimate of spawning biomass for BSAI Alaska skates is 108,093 t. The estimate of B40% is 74,457 t, so B/B40% is 1.45 and 2009 Alaska skate harvest levels can be assigned according to subtier 3a. Therefore, $F_{OFL}= F_{35\%}$ is 0.10 and maximum $F_{ABC}= F_{40\%}$ is 0.087. The corresponding 2009 OFL is 30,077 t and maximum allowable ABC is 25,854 t. Specifications for 2010 are given in Table 16.

Tier 5 estimates of ABC for BSAI Alaska skates are described in further detail below, are 30,487 t using *M*=0.1 and 39,633 t using *M*=0.13. The Tier 3 estimates represent a 15% lower value relative to Tier 5 estimates using *M*=0.10, and a 34.7% lower value relative to Tier 5 estimates using *M*=0.13.

Recommended ABC for 2009

This assessment uses the best available data from current research and provides a more realistic analysis of the Alaska skate population in the BSAI than a Tier 5 approach. In particular, the model captures the late maturation (and thus reduced spawning biomass) of Alaska skates, while the Tier 5 approach does not. For these reasons, we recommend that Tier 3a criteria be applied and recommend a 2009 ABC of 25,854 t.

ALASKA SKATE and OTHER SKATES – Tier 5 assessment

Data

Survey biomass

The biomass of the skate assemblage as a whole has increased since the early 1980s (Table 17, Fig. 39). Because skates as a group are contiguous and found in nearly all habitats, the uncertainty (measured as the coefficient of variation, CV) in aggregate skate biomass estimates is rather low, but the uncertainty for individual species is greater (Table 2). Survey species identification are considered reliable after 1998. Unfortunately, due to taxonomic uncertainty, we cannot evaluate individual species trends within the complex for surveys prior to 1999. Recent surveys demonstrate the variable species composition of the skate complex within each of the three habitat areas, the EBS shelf, the EBS slope, and the Aleutian Islands (Figure 3). The Alaska skate (*B. parmifera*) is dominant and highly abundant on the EBS shelf, while in each of the other two habitat areas, the skate species composition is far more diverse, especially on the EBS slope (Table 2). To generate harvest recommendations, we used the average biomass for each area during 2000-2008. This approach allowed the use of four surveys in the AI, nine surveys from the EBS shelf, and two surveys from the EBS slope. The 2002 biomass estimate from the slope was excluded because it is much higher than the estimate from the other two years and was affected by extremely high catch of skates in a single tow.

Analytic Approach and Results

Parameters estimated independently: *M*

As in previous years, *M* was estimated based on other life history parameters. Several methods were employed based on correlations of *M* with life history parameters including growth parameters (Alverson and Carney 1975, Pauly 1980, Charnov 1993), longevity (Hoenig 1983), and reproductive potential (Rikhter and Efanov 1976, Roff 1986). Natural mortality was estimated using life history parameters from California big skate (*Raja binoculata*) and longnose skate (*R. rhina*) (Zeiner and Wolf 1993), which are found in the GOA but are rare in the BSAI. We also estimated *M* for big and longnose skates from British Columbia and the Gulf of Alaska based on two life history studies (McFarlane and King 2006, Gburski et al. 2007). These estimates of *M* are close to the estimate of *M*=0.10 derived from CA big and longnose skates, which has been accepted by the Plan Team and the SSC as a reasonable approximation of "aggregate skate" *M* for the Other Skates group. Considering the uncertainty inherent in applying this method to the multi-species Other Skates group, we elected to use the lowest estimate of *M* (*M*=0.10, Table 18), which results in conservative estimates of ABC and OFL under Tier 5 criteria. Until better information is available on the productivity of individual skate species in the BSAI Other Skates group, we recommend this strategy in the interim in order to promote skate conservation while still allowing for historical levels of incidental catch in target groundfish fisheries.

Alaska skate natural mortality was derived using the methods described above in conjunction with available species-specific life history parameters (Table 12). These methods are described in more detail in the Tier 3 Alaska skate assessment section. The lowest Alaska skate mortality rate of *M*=0.10 was applied to obtain estimates of ABC and OFL using Tier 5 methodology. The age-structured assessment estimate of *M*=0.13 was also applied to Alaska skate biomass estimates using Tier 5 for comparison.

Projections and Harvest Alternatives

Acceptable Biological Catch and Overfishing Limit

We recommend that a Tier 5 approach be applied to the Other Skate species complex if the catch remains incidental and no target fishery develops. Tier 5 is recommended because reliable estimates of biomass

exists, and *M* =0.10 is considered a reasonable approximation of "aggregate skate" *M* by the Plan Team and SSC. We note that though the proxy *M* was applied to all species, it was based on relatively sensitive skate species. Therefore it is likely an underestimate of *M* for more productive species, which results in conservative specifications.

Tier 5 specifications for the Alaska skate are also shown here for comparison with the specifications generated from the single-species population model described earlier in this assessment. In addition to the default multi-species *M*=0.10 from last year's assessment, a new estimate of *M*=0.13 has been used to generate the Tier 5 ABC and OFL. This new estimate of *M*, based on Alaska skate life history parameters, has been reviewed and accepted by the Plan Team and the SSC as a reasonable approximation of natural mortality for this species. Biomass estimates for both the Alaska skate and the Other Skates group were used from years when research survey species identification is considered most reliable (1999-2008).

Tier 6 is not recommended because the catch history for skates is not considered reliable (reported as "Other species"), and average catch for untargeted species is likely to constrain target fisheries if used to specify harvest limits. For the Tier 5 estimate, we recommend using a 9-year average of skate biomass so that we may include multiple estimates from each of the trawl surveys, while capturing recent biomass levels.

	Alaska skate			other skates		
	EBS shelf	AI	EBS slope	EBS shelf	AI	EBS slope
2000	300,954	9,801		24,338	19,518	
2001	402,909			17,405		
2002	347,873	10,662	35,932	18,441	23,752	33,344
2003	354,244			32,095		
2004	402,354	12,727	4,248	14,205	40,344	28,908
2005	461,067			20,127		
2006	424,511	13,484		18,045	40,726	
2007	457,941			17,083		
2008	362,127		4,516	19,617		33,033
average	**390,442**	**11,668**	**4,382**	**20,151**	**31,085**	**30,970**

Tier 5 Alaska skate ABC

Applying the default multi-species *M* estimate of 0.10 to the 9-year average of bottom trawl survey biomass estimates, we calculate an ABC of 0.75 * 0.10 * (total BSAI biomass of 406,492) = 0.075 * 406,492 t = **30,487 t**.

Alternatively, applying the Alaska skate-specific *M* estimate of 0.13 to the 9 year average of bottom trawl survey biomass estimates, we calculate an ABC of 0.75 * 0.13 * (total BSAI biomass of 406,492) = 0.0975 * 406,492 t = **39,633 t**.

Tier 5 Alaska skate OFL

Applying the default multi-species *M* estimate of 0.10 to the 9-year average of bottom trawl 406,492 t = **40,649 t**.

Applying the Alaska skate-specific *M* estimate of 0.13 to the 9-year average of bottom trawl survey biomass estimates, we calculate an OFL of 0.13 * (total BSAI biomass of 406,492) = 0.13 * 406,492 t = **52,844 t**.

Tier 5 Recommended Other Skates ABC

Applying the *M* estimate of 0.10 to the 9-year average of bottom trawl survey biomass estimates, we calculate an ABC of 0.75 * 0.10 * (total BSAI biomass of 82,206 t) = 0.075 * 82,206 t = **6,165 t**.

Tier 5 Recommended Other Skates OFL

Applying the *M* estimate of 0.10 to the 9 year average of bottom trawl survey biomass estimates, we calculate an OFL of 0.10 * (total BSAI biomass of 82,206 t) = 0.1 * 82,206 t = **8,221 t**.

In the event the SSC chooses Tier 5 criteria for the Alaska skate, we strongly recommend using the default estimate of *M*=0.10 to generate specifications. Increasing *M* from 0.10 to 0.13 results in a substantial increase (32%, or approximately 10,000 t) from last year's ABC. Because skates are managed with the rest of the Other species complex under a single TAC, such an increase would reduce protection to all the species within this complex, and could encourage the development of directed fisheries. Because we do not yet fully understand skate population dynamics or the effects of directed fishing on species within the Other species complex, we feel it is prudent to use the more conservative estimate of *M* until such information becomes available.

Assemblage analysis and recommendations

Because skates represent a potentially valuable fishery resource as well as a potentially sensitive species group, we recommend that they be managed separately from the BSAI Other species complex. There is a reliable biomass time series for the skate assemblage as a whole in both the EBS and AI, and recently (since 1999) there are also reliable estimates of biomass for each species within the assemblage.

We further recommend splitting the Alaska skate (*Bathyraja parmifera*) from the BSAI skate assemblage to form two management groups: Alaska skate and "Other Skates". The purpose of separate recommendations is to provide increased protection to rare or endemic species in the EBS slope and AI habitat areas, since the Alaska skate constitutes the bulk of the skate biomass in the EBS shelf habitat area. We have shown that the distribution of species differs greatly by habitat areas within the BSAI, and that overall catch is not necessarily in proportion to BSAI-wide biomass due to the distribution of fishing effort. Because it would be difficult to manage skates by habitat area, managing Alaska skates and the Other Skates complex separately represents a reasonable compromise which increases protection to the species within each ecosystem but maintains a level of management simplicity appropriate to nontarget species complexes. In the event that target fisheries develop for individual skate species in the Other Skates complex, we would recommend that target skate species be further separated from the complex and managed individually. Furthermore, directed fishing for skates in the BSAI should only be allowed when sufficient life history information becomes available to make reasonable species-specific estimates of productivity.

Ecosystem Considerations

This section focuses on the Alaska skate in both the EBS and AI, with all other species found in each area summarized within in the group "Other Skates." We also include supplemental information on the other biomass dominant species in the AI, the Aleutian and whiteblotched skates. This level of aggregation is necessary due to current data constraints, but improved species-specific information will be incorporated as it becomes available.

Skates are predators in the BSAI FMP area. Some species are piscivorous while others specialize in benthic invertebrates; additionally, at least three species, deepsea skate, roughtail skate, and longnose skate, are benthophagic during the juvenile stage but become piscivorous as they grow larger (Ebert 2003, Robinson 2006) (Table 1). Each skate species would occupy a slightly different position in EBS and AI food webs based upon its feeding habits, but in general skates as a group are predators at a relatively high trophic level. For simplicity, we show the food webs for all skate species combined in each system (Figure 40; EBS in upper panel, AI in lower panel). In the EBS food web, the skate biomass and therefore the general skate food web position is dominated by the Alaska skate, which eats primarily pollock (as do most other piscivorous animals in the EBS). The food web indicates that aside from sperm whales, most of the "predators" of EBS skates are fisheries, and that cod and halibut are both predators and prey of skates. The AI food web shows skates with different predators and prey than in the EBS, but still at the same moderately high trophic level. Relative to EBS skates, AI skates display more diet diversity (because the species complex is more diverse than in the Alaska skate-dominated EBS), and have more non-fishery predators including sharks and sea lions. These food webs were derived from mass balance ecosystem models assembling information on the food habits, biomass, productivity and consumption for all major living components in each system (Aydin et al. in review).

The density and mortality patterns for skates also differ greatly between the EBS and AI ecosystems. The biomass density of Alaska skates is much higher in the EBS than in the AI (Fig. 41 upper left panel) and we now know they are likely separate species between the areas as well. The density of Alaska skates in the EBS also far exceeds that of all other *Bathyraja* species in any area (Fig. 41 upper right panel), but the density of other *Bathyraja* skates is highest in the AI. One simple way to evaluate ecosystem (predation) effects relative to fishing effects is to measure the proportions of overall mortality attributable to each source. The lower panels of Fig. 41 distinguish predation from fishing mortality, and further distinguish these measured sources of mortality from sources that are not explained within the ecosystem models, which are based on early 1990s fishing and food habits information. While there are many uncertainties in estimating these mortality rates, the results suggest that (early 1990s) fishing mortality exceeded predation mortality for Alaska skates and for Other Skates in the EBS and AI (and for Other Skates in the GOA as well). Furthermore, predation mortality appeared to be higher for AI skates than for EBS skates, both for Alaska and Other Skate species in the early 1990s, suggesting that skates experience higher overall mortality in the AI relative to the EBS. One source of uncertainty in these results is that all skate species in all areas were assumed to have the same total mortality rate, which is an oversimplification, but one which is consistent with the assumptions regarding natural mortality rate (the same for all skate species) in this stock assessment. We expect to improve on these default assumptions as data on productivity and catch for the skate species in each area continue to improve.

In terms of annual tons removed, it is instructive to compare fishery catches with predator consumption of skates. We estimate that fisheries were annually removing about 13,000 and 1,000 tons of skates from the EBS and AI, respectively on average during the early 1990s (Fritz 1996, 1997). While estimates of predator consumption of skates are perhaps more uncertain than catch estimates, the ecosystem models incorporate uncertainty in partitioning estimated consumption of skates between their major predators in each system. The predators with the highest overall consumption of Alaska skates in the EBS are sperm whales, which account for less than 2% of total skate mortality and consumed between 500 and 2,500 tons of skates annually in the early 1990s. Consumption of EBS Alaska skates by Pacific halibut and cod are too small to be reliably estimated (Fig. 42, left panels). Similarly, sperm whales account for less than 2% of Other Skate mortality in the EBS, but are still the primary predator of Other Skates there, consuming an estimated 50 to 400 tons annually. Pacific halibut consume very small amounts of Other Skates in the EBS, according to early 1990s information integrated in ecosystem models (Fig. 42, right panels). The predators with the highest consumption of Alaska skates in the AI are also sperm whales, which account for less than 2% of total skate mortality and consumed between 20 and 120 tons of skates

annually in the early 1990s. Pinnipeds (Steller sea lions) and sharks also contributed to Alaska skate mortality in the AI, averaging less than 50 tons annually (Fig. 43, left panels). Similarly, sperm whales account for less than 2% of Other Skate mortality in the AI, but are still the primary predator of Other Skates there, consuming an estimated 20 to 150 tons annually. Pinnipeds and sharks consume very small amounts of Other Skates in the AI, according to early 1990s information (Fig. 43, right panels). Gerald Hoff's research on skate nursery areas suggests that gastropod predation on skate egg cases may account for a significant portion of mortality during the embryonic stage, and Pacific cod and Pacific halibut consume substantial numbers of newly hatched juvenile skates within nursery areas. These sources of mortality may be included in future stock assessments.

Diets of skates are derived from food habits collections taken in conjunction with EBS and AI trawl surveys. Skate food habits information is more complete for the EBS than for the AI, but we present the best available data for both systems here. Over 40% of EBS Alaska skate diet measured in the early 1990s was adult pollock, and another 15% of the diet was fishery offal, suggesting that Alaska skates are opportunistic piscivores (Fig. 44, upper left panel). Eelpouts, rock soles, sandlance, arrowtooth flounder, salmon, and sculpins made up another 25-30% of Alaska skates' diet, and invertebrate prey made up the remainder of their diet. This diet composition combined with estimated consumption rates and the high biomass of Alaska skates in the EBS results in an annual consumption estimate of 200,000 to 350,000 tons of pollock annually (Fig. 44, lower left panel). EBS Other Skates also consume pollock (45% of combined diets), but their lower biomass results in consumption estimates ranging from 20,000 to 70,000 tons of pollock annually (Fig. 44, right panels). Other Skates tend to consume more invertebrates than Alaska skates in the EBS, so estimates of benthic epifaunal consumption due to Other Skates range up to 50,000 tons annually, higher than those for Alaska skates despite the disparity in biomass between the groups (Fig. 44, lower panels).

Because Alaska skates and all Other Skates are distributed differently in the EBS, with Alaska skates dominating the shallow shelf areas and the more diverse species complex located on the outer shelf and slope, we might expect different ecosystem relationships for skates in these habitats based on differences in food habits among the species. Similarly, in the AI the unique skate complex has different diet compositions and consumption estimates from those estimated for EBS skates. The skate in the AI formerly known as the Alaska skate is opportunistically piscivorous like its EBS relative, feeding on the common commercial forage fish, Atka mackerel (65% of diet) and pollock (14% of diet), as well as fishery offal (7% of diet; Fig. 45 upper left panel). Diets of Other Skates in the AI are more dominated by benthic invertebrates, especially shrimp (pandalid and non-pandalid total 42% of diet), but include more pelagic prey such as juvenile pollock, adult Atka mackerel, adult pollock and squids (totaling 45% of diet; Fig. 45 upper right panel). Estimated annual consumption of Atka mackerel by AI (former) Alaska skates in the early 1990s ranged from 7,000 to 15,000 tons, while pollock consumption was below 5,000 tons (Fig. 45 lower left panel). Shrimp consumption by AI Other Skates was estimated to range from 4,000 to 15,000 tons annually in the early 1990s, and consumption of pollock ranged from 2,000 to 10,000 tons (Fig. 45 lower right panel). Atka mackerel consumption by AI Other Skates was estimated to be below 5,000 tons annually. The diet composition estimated for AI Other Skates is likely dominated by the biomass dominant species in that system, whiteblotched skate and Aleutian skate. The diet compositions of both Aleutian and whiteblotched skates in the AI appear to be fairly diverse (Fig. 46), and are described in further detail in Yang (2007) along with the diets of big skate, Bering skate, Alaska skate, roughtail skate, and mud skate in the AI. In the future, we hope to use diet compositions to make separate consumption estimates for whiteblotched and Aleutian skates along with (former) Alaska skates in the AI.

Examining the trophic relationships of EBS and AI skates provides a context for assessing fishery interactions beyond the direct effect of bycatch mortality. In both areas, the biomass-dominant species of skates feed on commercially important fish species, so it is important for fisheries management to

maintain the health of pollock and Atka mackerel stocks in particular to maintain the forage base for skates (as well as for other predators and for human commercial interests).

Data gaps and research priorities

Aggregate skate and Alaska skate catches have been estimated using several different methods each with a number of inherent assumptions. We used species composition from recent surveys to partition the Alaska skate catch; however there are two caveats involved with this approach: 1) we assume species composition has remained constant prior to 1999, and 2) we assume that survey species composition is representative of the catch species composition. Also, aggregate skate catch records can mask shifts in species composition, and fishing gear may be more selective for larger-bodied species. Species identification by fishery observers has vastly improved in recent years; however it is still difficult to make accurate identifications in the longline fishery, as many skates are dropped off the line without being brought on board. Mounted video camera systems may be a cost-effective way to determine the species composition of the catch in the future.

In the Alaska skate model, we assumed a catch rate with 100% mortality. In reality, skate mortality is dependent upon the time spent out of water, the type of gear, and handling practices after capture. From fishery observer data, approximately 30% of skates are retained; however we currently have no information regarding the survival of skates that are discarded at sea.

Very few biomass indices are available from the Bering Sea slope survey. The Bering Sea slope habitat area has very high skate species diversity, yet there are only two years of survey data from this area where species identification can be considered reliable (2002 and 2004). Continuation of the Bering Sea slope survey, at least in alternate years, would help to identify overall trends in skate abundance as well as potential shifts in the relative species composition there.

We have initiated a tagging program to gather information regarding movement, distribution patterns, and growth of the Alaska skate. In 2008, approximately 1,200 skates were tagged and released during the shelf and slope surveys. The vast majority of these releases occurred during the shelf survey, and releases were distributed over the entire shelf survey area. As of October 2008, two of these tags had already been recovered through the commercial longline fishery. We expect to deploy additional tags during future trawl surveys and other research cruises.

Fecundity is a very difficult quantity to measure in skates, as individuals of some species may reproduce throughout the year and thus the number of mature or maturing eggs present in the ovary may represent only a fraction of the annual reproductive output. Matta (2006) estimated the average fecundity of the Alaska skate to range between 21 and 37 eggs per female per year, based on the assumed relationship between reproductive potential and *M* (Gunderson 1997). However, due to the uncertainty involved with this parameter, fecundity estimates were not included in the stock assessment model. Fecundity estimates for other skate species range from 48 to 150 young per year (Holden et al. 1971; Holden 1975; Luer & Gilbert 1985; Ellis & Shackley 1995), and it is conceivable that the Alaska skate also has very low annual fecundity. Additional work, such as laboratory rearing experiments, is needed to validate these estimates.

Skate habitat is only beginning to be described in detail. Adults appear capable of significant mobility in response to general habitat changes, but any effects on the small scale nursery habitats crucial to reproduction could have disproportionate population effects. Eggs are mostly limited to isolated nursery grounds, and juveniles use different habitats than adults. Changes in these habitats have not been monitored historically, so assessments of habitat quality and its trends are not currently available. We recommend continued study of skate nursery areas to evaluate their importance to population production.

Because skates are at a relatively high trophic level in the EBS and AI, predation mortality is less significant than fishing mortality for adult skates. Therefore, the assessment of skate population dynamics and response to fishing should be continued and improved as fishing represents the largest explained source of mortality in the EBS and AI (especially since this mortality is not from targeted fishing, but from incidental catch). Highest priority research should continue to focus on direct fishing effects on skate populations. The most important component of this research is to fully evaluate the productive capacity of skate populations, including information on age and growth, maturity, fecundity, and habitat associations. This research has been initiated for major skate species in the EBS and AI, and some results have already become available. Such research should be fully funded to completion.

Juvenile skates and skate egg cases are likely to be much more vulnerable to predation and disturbance than adults. Gerald Hoff's (AFSC) work on skate nursery areas, described in the life history section of this assessment, suggests that the egg case and neonate life history stages are susceptible to predation by snails and some groundfish. Differences between life history stages in terms of predation and effects of trawling on nursery areas have not been examined in population or ecosystem models.

The PSRC (MLML) has recently received funding from the North Pacific Research Board (NPRB) to examine the feeding habits of Aleutian, Bering, big, and longnose skates. Simon Brown, a graduate student, is currently working on this project. Specific objectives are to: 1) determine the diets of Alaskan skate species through analysis of stomach contents, 2) examine temporal, ontogenetic, and intergender differences in diet for each species, 3) investigate aspects of foraging habitat and trophic relationships for each species, and 4) compare interspecific diets of these Alaskan skate species to determine degree of dietary overlap. The results of this study will provide basic biological information on skates for inclusion in multi-species and predator/prey models.

We do not see any conflict at present between commercial fishing and skate foraging on pollock or Atka mackerel, but we do recommend continued monitoring of skate populations and food habits at appropriate spatial scales to ensure that these trophic relationships remain intact as fishing for these commercial forage species continues and evolves.

Ecosystem Effects on Stock and Fishery Effects on the Ecosystem: Summary

In the following tables, we summarize ecosystem considerations for BSAI skates and the entire groundfish fishery where they are caught incidentally. Because there is no "skate fishery" in the EBS or AI at present, we attempt to evaluate the ecosystem effects of skate bycatch from the combined groundfish fisheries operating in these areas in the second portion of the summary table. The observation column represents the best attempt to summarize the past, present, and foreseeable future trends. The interpretation column provides details on how ecosystem trends might affect the stock (ecosystem effects on the stock) or how the fishery trend affects the ecosystem (fishery effects on the ecosystem). The evaluation column indicates whether the trend is of *no concern, probably no concern, possible concern, definite concern,* or *unknown.*

Ecosystem effects on BSAI Skates *(evaluating level of concern for skate populations)*

Indicator	Observation	Interpretation	Evaluation
Prey availability or abundance trends			
Pollock	Currently declining from high biomass levels	Probably still adequate forage available for piscivorous skates	Probably no concern
Atka mackerel	Cyclically varying population with slight upward trend overall 1977-2005	Adequate forage available for piscivorous skates	No concern
Shrimp/ Benthic invertebrates	Trends are not currently measured directly, only short time series of food habits data exist for potential retrospective measurement	Unknown	Unknown
Predator population trends			
Sperm whales	Populations recovering from whaling?	Possibly higher mortality on skates? But still a very small proportion of mortality	No concern
Steller sea lions	Declined from 1960s, low but level recently	Lower mortality on skates?	No concern
Sharks	Population trends unknown	Unknown	Unknown
Changes in habitat quality			
Benthic ranging from shallow shelf to deep slope, isolated nursery areas in specific locations	Skate habitat is only beginning to be described in detail. Adults appear adaptable and mobile in response to habitat changes. Eggs are limited to isolated nursery grounds and juveniles use different habitats than adults. Changes in these habitats have not been monitored historically, so assessments of habitat quality and its trends are not currently available.	Continue study on small nursery areas to evaluate importance to population production	Possible concern if nursery grounds are disturbed or degraded.

Groundfish fishery effects on ecosystem via skate bycatch *(evaluating level of concern for ecosystem)*

Indicator	Observation	Interpretation	Evaluation
Fishery contribution to bycatch			
Skate catch	Has varied from 12,226 t to 22,982 t from 1992-2007	Largest portion of total mortality for skates	Possible concern
Forage availability	Skates have few predators, and skates are small proportion of diets for their predators	Fishery removal of skates has a small effect on predators	Probably no concern
Fishery concentration in space and time			
	Skate bycatch is spread throughout FMP areas, although higher proportion of skate bycatch occurs on outer continental shelf and upper slope	Potential impact to skate populations if fishery disturbs nursery or other important habitat, but small effect on skate predators	Possible concern for skates, probably no concern for skate predators
Fishery effects on amount of large size target fish			
	Survey length compositions (2000-2007) suggest that large size classes of Alaska skates appear to be stable	Fishery removals do not appear to have an effect on size structure	Probably no concern
Fishery contribution to discards and offal production			
	Skate discard is a relatively high proportion of skate catch, some incidentally caught skates are retained and processed	Unclear whether discard of skates has ecosystem effect	Unknown
Fishery effects on age-at-maturity and fecundity			
	Skate age at maturity and fecundity are just now being described; fishery effects on them difficult to determine due to lack of unfished population to compare with	Unknown	Unknown

Summary

Recommendations	Alaska skate	Alaska skate	Alaska skate	Other Skates
M	0.13	0.10	0.13	0.10
Tier	3	5	5	5
2009				
proj./avg. biomass	551,439	406,492	406,492	82,206
F_{OFL}	0.10	0.10	0.13	0.10
Max F_{ABC}	0.087	0.075	0.0975	0.075
recommended F_{ABC}	0.087	0.075	0.0975	0.075
OFL	30,077	40,649	52,844	8,221
Max ABC	25,854	30,487	39,633	6,165
recommended ABC	25,854	30,487	39,633	6,165
2010				
proj./avg. biomass	549,000	406,492	406,492	82,206
F_{OFL}	0.10	0.10	0.13	0.10
Max F_{ABC}	0.087	0.075	0.0975	0.075
recommended F_{ABC}	0.087	0.075	0.0975	0.075
OFL	30,009	40,649	52,844	8,221
Max ABC	25,796	30,487	39,633	6,165
recommended ABC	25,796	30,487	39,633	6,165

Acknowledgements

Many thanks to the following for their valuable contributions to this document: Bob Lauth, Mark Wilkins, and others in the AFSC RACE program for providing survey biomass estimates, the AFSC's Age and Growth Program for providing skate ages, the AFSC's Fishery Monitoring and Analysis program for their hard work in the field and compiling data, and the Alaska Regional Office for making nontarget species catch estimates available. Jim Ianelli provided the projection model. Rick Methot, Grant Thompson, Anne Hollowed, Vladlena Gertseva, and Martin Dorn provided valuable advice regarding the age-structured model.

Literature Cited

Agnew, D.J., C.P. Nolan, J.R. Beddington, and R. Baranowski. 2000. Approaches to the assessment and management of multispecies skate and ray fisheries using the Falkland Islands fishery as an example. Can. J. Fish. Aquat. Sci. 57: 429-440.

Allen, M.J., and G.B. Smith. 1988. Atlas and zoogeography of common fishes in the Bering Sea and northeastern Pacific. NOAA Technical Report NMFS 66, 151 pp.

Alverson, D.L., and W.T. Pereyra. 1969. Demersal fish explorations in the northeastern Pacific Ocean: An evaluation of exploratory fishing methods and analytical approaches to stock size and yield forecasts. J. Fish. Res. Bd. Canada 26: 1985-2001.

Alverson, D.L., and M.J. Carney. 1975. A graphic review of the growth and decay of population cohorts. J. Cons. Int. Explor. Mer 36:133-143.

Aydin, K., S. Gaichas, I. Ortiz, D. Kinzey, and N. Friday. In review. A comparison of the Bering Sea, Gulf of Alaska, and Aleutian Islands large marine ecosystems through food web modeling. NOAA Tech Memo.

Brander, K. 1981. Disappearance of common skate *Raja batis* from Irish Sea. Nature 290: 48-49.

Casey, J.M. and R.A. Myers. 1998. Near extinction of a large, widely distributed fish. Science 281(5377):690-692.

Charnov, E.L. 1993. Life history invariants some explorations of symmetry in evolutionary ecology. Oxford University Press Inc., New York. 167p.

Davis, C.D. 2006. Age, growth, and reproduction of the roughtail skate, *Bathyraja trachura* (Gilbert, 1892). M.S. thesis, Moss Landing Marine Laboratories, CSU Monterey Bay.

Dulvy, N.K., J.D. Metcalfe, J. Glanville, M.G. Pawson, and J.D. Reynolds. 2000. Fishery stability, local extinctions, and shifts in community structure in skates. Conservation Biology 14(1): 283-293.

Ebert, D.A. 2003. Sharks, rays, and chimeras of California. University of California Press, Berkeley, CA, 285 pp.

Ebert, D.A. 2005. Reproductive biology of skates, *Bathyraja* (Ishiyama), along the eastern Bering Sea continental slope. J. Fish. Biol. 66: 618-649.

Ebert, D.A., Smith, W.D., Haas, D.L., 1, Ainsley, S.M., Cailliet, G.M. 2007. Life history and population dynamics of Alaskan skates: providing essential biological information for effective management of bycatch and target species. Final Report to the North Pacific Research Board, Project 510.

Eschmeyer, W.N., E.S. Herald, and H. Hammann. 1983. A field guide to Pacific coast fishes of North America. Houghton Mifflin Co., Boston: 336 pp.

Frisk, M.G., T. J. Miller, and M. J. Fogarty. 2001. Estimation and analysis of biological parameters in elasmobranch fishes: a comparative life history study. Can. J. Fish. Aquat. Sci. 58: 969-981.

Frisk, M. G. , T. J. Miller, and M. J. Fogarty. 2002. The population dynamics of little skate *Leucoraja erinacea*, winter skate *Leucoraja ocellata*, and barndoor skate *Dipturus leavis*: predicting exploitation limits using matrix analysis. ICES J. Mar. Sci. 59: 576-586.

Fritz, L. W. 1996. Squid and other species. Chapter 13 In Stock Assessment and Fishery Evaluation Report for the Groundfish Resources of the Bering Sea/Aleutian Islands Region. North Pacific Fishery Management Council, 605 W. 4th Ave., Suite 306, Anchorage, AK 99501.

Fritz, L. W. 1997. Squid and other species. Pp. 463-484 In Stock Assessment and Fishery Evaluation Report for the Groundfish Resources of the Bering Sea/Aleutian Islands Region. North Pacific Fishery Management Council, 605 W. 4th Ave., Suite 306, Anchorage, AK 99501.

Fritz LW, Greig A, Reuter RF. 1998. Catch-per-unit-effort, length, and depth distributions of major groundfish and bycatch species in the Bering Sea, Aleutian Islands, and Gulf of Alaska regions based on groundfish fishery observer data, 179 p. NTIS No. PB98-139298

Gaichas, S., J. Ianelli, and L. Fritz. 1999. Other species considerations for the Gulf of Alaska.

Gaichas, S., M. Ruccio, D. Stevenson, and R. Swanson. 2003. Stock assessment and fishery evaluation for skate species (Rajidae) in the Gulf of Alaska.

Gburski, C.M., S.K. Gaichas, and D.K. Kimura. 2007. Age and growth of big skate (*Raja binoculata*) and longnose skate (*R. rhina*) and implications to the skate fisheries in the Gulf of Alaska. Env. Bio. Fishes 80: 337-349.

Gunderson, D.R. 1997. Trade-off between reproductive effort and adult survival in oviparous and viviparous fishes. Can. J. Fish. Aquat. Sci. 54: 990-998.

Gunderson, D.R., Zimmerman, M., Nichol, D.G., and Pearson, K. 2003. Indirect estimates of natural mortality rate for arrowtooth flounder (*Atheresthes stomias*) and darkblotched rockfish (*Sebastes crameri*). Fishery Bulletin 101: 175 – 182.

Hoenig, J.M. 1983. Empirical use of longevity data to estimate mortality rates. Fish. Bull. 82(1): 898-902.

Hoff, G.R. 2006. Investigations of a skate nursery area in the eastern Bering Sea. Progress report to the NPRB. March 7, 2006.

Hoff, G.R. 2007. Reproduction of the Alaska skate (*Bathyraja parmifera*) with regard to nursery sites, embryo development and predation. PhD dissertation, University of Washington, Seattle.

Ishihara, H. and R. Ishiyama. 1985. Two new North Pacific skates (Rajidae) and a revised key to *Bathyraja* in the area. Jpn. J. Ichthyol. 32(2): 143-179.

Jensen, A.L. 1996. Beverton and Holt life history invariants result from optimal trade-off of reproduction and survival. Can. J. Aquat. Fish. Sci. 53: 820-822.

King, J.R., and G.A. McFarlane. 2003. Marine fish life history strategies: applications to fishery management. Fish. Man. and Ecology, 10: 249-264.

Kotwicki, S., and Weinberg, K.L. 2005. Estimating capture probability of a survey bottom trawl for Bering Sea skates (*Bathyraja spp.*) and other fish. Alaska Fishery Research Bulletin 11(2): 135-145.

Martin, L. and G.D. Zorzi. 1993. Status and review of the California skate fishery. In Conservation biology of elasmobranchs (S. Branstetter, ed.), p. 39-52. NOAA Technical Report NMFS 115.

Matta, M.E. 2006. Aspects of the life history of the Alaska skate, *Bathyraja parmifera*, in the eastern Bering Sea. M.S. thesis, University of Washington, Seattle.

McEachran, J.D., and K.A. Dunn. 1998. Phylogenetic analysis of skates, a morphologically conservative clade of elasmobranchs (Chondrichthyes: Rajidae). Copeia, 1998(2), 271-290.

McEachran, J.D. and T. Miyake. 1990a. Phylogenetic relationships of skates: a working hypothesis (Chondrichthyes, Rajoidei). In Elasmobranchs as living resources: advances in the biology, ecology, systematics, and the status of the fisheries (H.L. Pratt, Jr., S.R. Gruber, and T. Taniuchi, eds.), p. 285-304. NOAA Technical Report NMFS 90.

McFarlane, G.A. and J.R. King. 2006. Age and growth of big skate (*Raja binoculata*) and longnose skate (*Raja rhina*) in British Columbia waters. Fish Res. 78: 169-178.

Mecklenberg, C.W., T.A. Mecklenberg, and L.K. Thorsteinson. 2002. Fishes of Alaska. American Fisheries Society, 1037 pp.

Methot RD. 1990. Synthesis model: an adaptable framework for analysis of diverse stock assessment data. International North Pacific Fisheries Commission Bulletin 50:259-277

Methot RD. 2005. Technical description of the Stock Synthesis II assessment program. NOAA Fisheries, Seattle, WA.

Methot, R. 2007. User manual for the integrated analysis program Stock Synthesis 2 (SS2). Model version 2.00b. Northwest Fisheries Service, NOAA Fisheries, Seattle, WA.

Moyle, P.B., and J.J. Cech, Jr. 1996. Fishes, an introduction to ichthyology (Third edition). Prentice Hall: New Jersey, 590 pp.

Murray, J.D. 1989. Mathematical Biology. Springer-Verlag: New York. 767 pp.

Musick, J.A., S.A. Berkeley, G.M. Cailliet, M. Camhi, G. Huntsman, M. Nammack, and M.L. Warren, Jr. 2000. Protection of marine fish stocks at risk of extinction. Fisheries 25(3):6-8.

Myers RA, Bowen KG, Barrowman NJ. 1999. Maximum reproductive rate of fish at low population sizes. Canadian Journal of Fisheries and Aquatic Sciences 56:2404-2419

New England Fishery Management Council (NEFMC). Skate Fishery Management Plan. http://www.nefmc.org/skates/index.html

Nelson, J. S. 1994. Fishes of the world, Third edition. John Wiley and Sons, Inc., New York: 600 pp.

NMFS 2000. Skate complex. In Draft 30th Northeast Regional Stock Assessment Workshop (30th SAW), Stock assessment review committee (SARC) consensus summary of assessments, p. 7-173.

NMFS PSEIS 2001. Draft Programmatic Environmental Impact Statement.

Northeast Fisheries Science Center (NEFSC). 2007. 44th Northeast Regional Stock Assessment Workshop (44th SAW). Section B. Skate Complex: Assessment Summary for 2006. *In:* 44th SAW assessment summary report. US Dep Commer, Northeast Fish Sci Cent Ref Doc. 07-03; 58 p.

Orlov, A.M. 1998. The diets and feeding habits of some deep-water benthic skates (Rajidae) in the Pacific waters off the northern Kuril Islands and southeastern Kamchatka. Alaska Fishery Research Bulletin 5(1): 1-17.

Orlov, A.M. 1999. Trophic relationships of commercial fishes in the Pacific waters off southeastern Kamchatka and the northern Kuril Islands. p. 231-263 in Ecosystem Approaches for Fishery Management, AK Sea Grant College Program AK-SG-99-01, U. of AK Fairbanks, 756 pp.

Ormseth, O.A. and B. Matta. 2007. Gulf of Alaska skates. In: Stock Assessment and Fishery Evaluation Report for the Groundfish Resources of the Gulf of Alaska Region. North Pacific Fishery Management Council, 605 W. 4th Ave., Suite 306, Anchorage, AK 99501.

Pauly, D. 1980. On the interrelationships between natural mortality, growth parameters, and mean environmental temperature in 175 fish stocks. J. Cons. Int. Explor. Mer 39(2):175-192.

Rikhter, V.A., and V.N. Efanov. 1976. On one of the approaches to estimation of natural mortality of fish populations. ICNAF Res. Doc. 76/VI/8. Serial N. 3777. 13p.

Robinson, H.J. 2006. Dietary analysis of the longnose skate, *Raja rhina* (Jordan and Gilbert, 1880), in California waters. M.S. thesis, Moss Landing Marine Laboratories, CSU Monterey Bay.

Roff, D.A. 1986. The evolution of life history parameters in teleosts. Can. J. Fish. Aquat. Sci. 41:989-1000.

Sosebee, K. 1998. Skates. In Status of Fishery Resources off the Northeastern United States for 1998 (Stephen H. Clark, ed.), p. 114-115. NOAA Technical Memorandum NMFS-NE-115.

Stevenson, D. 2004. Identification of skates, sculpins, and smelts by observers in north Pacific groundfish fisheries (2002-2003), U.S. Department of Commerce Technical Memorandum NMFS-AFSC-142. 67 p.

Stevenson, D.E. and J.W. Orr. 2005. New records of two deepwater skate species from the eastern Bering Sea. Northwestern Naturalist 86: 71-81.

Stevenson, D.E., J.W. Orr, G.R. Hoff, and J.D. McEachran. 2004. *Bathyraja mariposa*: a new species of skate (Rajidae: Arhynchobatinae) from the Aleutian Islands. Copeia 2004(2):305-314.

Stevenson, D.E., J.W. Orr, G.R. Hoff, and J.D. McEachran. 2006. The skates of Alaska: distribution, abundance, and taxonomic progress. Marine Science in Alaska 2006 Symposium, Anchorage, AK, Jan 2006, poster.

Stevenson, D. E., Orr, J. W., Hoff, G. R., and McEachran, J. D. 2007. Field guide to sharks, skates, and ratfish of Alaska. Alaska Sea Grant.

Thompson, G.G. 1993. A proposal for a threshold stock size and maximum fishing mortality rate. Pages 303-320 in Risk evaluation and biological reference points for fisheries management (S.J. Smith, J.J. Hunt, and D. Rivard, eds.). Can. Spec. Publ. Fish. Aquat. Sci. 120, 440 pp.

Wakefield, W.W. 1984. Feeding relationships within assemblages of nearshore and mid-continental shelf benthic fishes off Oregon. M.S. Thesis, OSU.

Walker, P.A., and R. G. Hislop. 1998. Sensitive skates or resilient rays? Spatial and temporal shifts in ray species composition in the central and north-western North Sea between 1930 and the present day. ICES J. Mar Sci., 55: 392-402.

Winemiller, K.O., and K.A. Rose. 1992. Patterns of life history diversification in North American fishes: implications for population regulation. Can. J. Fish. Aquat. Sci. 49: 2196-2218.

Yang, M-S. 2007. Food habits and diet overlap of seven skate species in the Aleutian Islands. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-AFSC-177, 46 p.

Zeiner, S.J. and P. Wolf. 1993. Growth characteristics and estimates of age at maturity of two species of skates (*Raja binoculata* and *Raja rhina*) from Monterey Bay, California. In Conservation biology of elasmobranchs (S. Branstetter, ed.), p. 39-52. NOAA Technical Report NMFS 115.

Tables

Table 1. Life history and depth distribution information available for BSAI and GOA skate species, from Stevenson (2004) unless otherwise noted.

Species	Common name	Max obs. length (TL cm)	Max obs. age	Age, length Mature (50%)	Feeding mode [2]	N embryos/ egg case [1]	Depth range (m) [9]
Bathyraja abyssicola	deepsea skate	135 (M) [10] 157 (F) [11]	?	110 cm (M) [11] 145 cm (F) [13]	benthophagic; predatory [11]	1 [13]	362-2904
Bathyraja aleutica	Aleutian skate	150 (M) 154 (F) [12]	14 [6]	121 cm (M) 133 cm (F) [12]	predatory	1	15-1602
Bathyraja interrupta	Bering skate (complex?)	83 (M) 82 (F) [12]	19 [6]	67 cm (M) 70 cm (F) [12]	benthophagic	1	26-1050
Bathyraja lindbergi	Commander skate	97 (M) 97 (F) [12]	?	78 cm (M) 85 cm (F) [12]	?	1	126-1193
Bathyraja maculata	whiteblotched skate	120	?	94 cm (M) 99 cm (F) [12]	predatory	1	73-1193
Bathyraja mariposa [3]	butterfly skate	76	?	?	?	1	90-448
Bathyraja minispinosa	whitebrow skate	83[10]	?	70 cm (M) 66 cm (F) [12]	benthophagic	1	150-1420
Bathyraja parmifera	Alaska skate	118 (M) 119 (F) [4]	15 (M) 17 (F) [4]	9 yrs, 92cm (M) 10 yrs, 93cm(F) [4]	predatory	1	17-392
Bathyraja sp. cf. *parmifera*	"Leopard" parmifera	133 (M) 139 (F)	?	?	predatory	?	48-396
Bathyraja taranetzi	mud skate	67 (M) 77 (F) [12]	?	56 cm (M) 63 cm (F) [12]	predatory [13]	1	58-1054
Bathyraja trachura	roughtail skate	91 (M) [14] 89 (F) [11]	20 (M) 17 (F) [14]	13 yrs, 76 cm (M) 14 yrs, 74 cm (F)[14, 12]	benthophagic; predatory [11]	1	213-2550
Bathyraja violacea	Okhotsk skate	73	?	?	benthophagic	1	124-510
Amblyraja badia	roughshoulder skate	95 (M) 99 (F) [11]	?	93 cm (M) [11]	predatory [11]	1 [13]	1061-2322
Raja binoculata	big skate	244	15 [5]	6-8 yrs, 72-90 cm [7]	predatory [8]	1-7	16-402
Raja rhina	longnose skate	180	25 [5]	7-10 yrs, 65-83 cm [7]	benthophagic; predatory [15]	1	9-1069

[1] Eschemeyer 1983. [2] Orlov 1998 & 1999 (Benthophagic eats mainly amphipods, worms. Predatory diet primarily fish, cephalopods). [3] Stevenson et al. 2004. [4] Matta 2006. [5] Gburski et al. 2007. [6] Gburski unpub data. [7] McFarlane & King 2006. [8] Wakefield 1984. [9] Stevenson et al. 2006. [10] Mecklenberg et al. 2002. [11] Ebert 2003. [12] Ebert 2005. [13] Ebert unpub data. [14] Davis 2006. [15] Robinson 2006.

Table 2. Species composition of the EBS and AI skate complexes from the most recent AFSC bottom trawl surveys.

Skate species	Common name	2008 EBS shelf bio (t)	cv	2008 EBS slope bio (t)	cv	2006 Aleutians bio (t)	cv
Bathyraja abyssicola	deepsea			165	0.62	0	
Bathyraja aleutica	Aleutian	6,278	0.57	17,160	0.15	6,684	0.23
Bathyraja interrupta	Bering	9,943	0.16	2,520	0.16	186	0.55
Bathyraja lindbergi	Commander			3,437	0.15	0	
Bathyraja maculata	whiteblotched	238	1.00	4,574	0.17	29,712	0.19
Bathyraja minispinosa	whitebrow			1,934	0.17	0	
Bathyraja parmifera	Alaska	362,127	0.06	4,516	0.32	13,484	0.19
Bathyraja taranetzi	mud	125	1.00	1,018	0.22	2,970	0.28
Bathyraja trachura	roughtail			2,213	0.14	0	
Bathyraja violacea	Okhotsk			0		0	
Raja binoculata	big	2,870	0.63	0		568	0.72
Raja rhina	longnose	162	1.00	12	1.00	0	
Rajidae unid	Unidentified skate species					605	0.41
Total skate complex		**381,744**		**37,548**		**54,210**	

Table 3. Time series of BSAI Other Species ABC, TAC, OFL and catch (t), with skate catch proportion.

Year	Other species ABC	Other species TAC	Other species OFL	Other species catch	BSAI skate catch	Skate % of Other species catch
1991	28,700	15,000		17,199		
1992	27,200	20,000	27,200	33,075	16,962	51%
1993		22,610		23,851	12,226	51%
1994	27,500	26,390	141,000	24,555	14,223	58%
1995	27,600	20,000	136,000	22,213	14,892	67%
1996	27,600	20,125	137,000	21,440	12,643	59%
1997	25,800	25,800		25,176	17,747	70%
1998	25,800	25,800	134,000	25,531	19,318	76%
1999	32,860	32,860	129,000	20,562	14,080	68%
2000	31,360	31,360	71,500	26,108	18,877	72%
2001	33,600	26,500	69,000	27,178	20,570	76%
2002	39,100	30,825	78,900	28,619	21,279	74%
2003	43,300	32,309	81,100	26,150	19,419	74%
2004	46,810	27,205	81,150	29,637	22,462	76%
2005	53,860	29,000	87,920	29,505	22,982	78%
2006	58,882	29,000	89,404	26,798	19,992	75%
2007	68,800	37,355	91,700	26,668	18,558	70%
*2008	78,100	50,000	104,000	21,340	15,167	71%

Sources: Other species ABC, TAC, OFL and 1992-2002 Other species catch from AKRO website.
BSAI skate catch 1992-1996 from Fritz 1996, 1997, 1997-2002 from Gaichas et al. 2004.
2003-2007 Other Species and BSAI skate catch from AKRO CAS. *2008 data current as of 10/3/2008.

Table 4. Estimated catch (t) of all skate species combined by target fishery, gear, and area, 1997-2002. Source: Gaichas AFSC.

Target fishery	gear	1997	1998	1999	2000	2001	2002
Arrowtooth	hook n line		0.65	9.72	1.31		0.49
	trawl	1.62	117.64	17.74	43.02	89.98	81.55
Arrowtooth Total		1.62	118.29	27.46	44.33	89.98	82.04
Atka mackerel	trawl	110.51	130.81	126.66	71.50	80.57	73.30
Flatheadsole	trawl	777.22	1,867.59	1,215.15	1,655.80	1,752.36	1,530.37
Other	hook n line		10.42	26.07	52.48	70.43	31.17
	trawl						8.82
Other Total			10.42	26.07	52.48	70.43	39.98
OtherFlats	trawl	39.18	103.15	69.22	115.16	20.09	58.48
Pacific cod	hook n line	13,298.81	13,534.64	9,651.09	12,975.65	14,116.58	14,059.10
	pot	1.50	0.01	0.11	0.06	0.10	0.00
	trawl	715.23	770.48	984.30	1,053.86	631.91	1,400.41
Pacific cod Total		14,015.53	14,305.12	10,635.50	14,029.56	14,748.59	15,459.51
Pollock	trawl	349.73	405.67	375.87	598.19	627.58	807.04
Rock sole	trawl	679.20	558.69	322.21	334.28	820.60	836.61
Rockfish	hook n line	110.27	6.73	0.69	1.70	4.42	0.84
	trawl	30.05	39.94	53.61	50.53	47.67	78.14
Rockfish Total		140.32	46.67	54.30	52.23	52.09	78.99
Sablefish	hook n line	266.00	110.10	109.54	115.86	194.11	233.13
	pot			0.09	0.01	0.06	0.01
	trawl		0.06			1.24	
Sablefish Total		266.00	110.16	109.63	115.87	195.41	233.14
Turbot	hook n line	140.82	280.84	319.92	317.36	187.07	120.80
	pot			1.22			
	trawl	16.13	18.67	17.34	23.92	16.66	7.76
Turbot Total		156.95	299.51	338.48	341.28	203.73	128.57
Unknown	hook n line	0.11	2.00	1.16	0.95	0.21	
	trawl		1.09		0.01	0.11	
Unknown Total		0.11	3.09	1.16	0.95	0.32	
Yellowfinsole	trawl	1,210.99	1,358.70	778.11	1,464.90	1,908.69	1,950.67
Grand Total		17,747.37	19,317.86	14,079.84	18,876.53	20,570.46	21,278.69

FMP area	area	1997	1998	1999	2000	2001	2002
AI	541	569.98	640.25	462.61	501.96	540.77	288.88
	542	200.87	369.17	239.96	608.31	422.64	217.74
	543	86.30	119.02	99.79	698.20	1,546.14	188.84
AI Total		857.15	1,128.45	802.36	1,808.47	2,509.56	695.46
EBS	509	1,920.87	2,317.12	2,033.62	2,830.27	3,092.09	3,112.51
	512	0.92		14.33		91.68	132.82
	513	2,572.53	2,605.18	1,993.53	2,641.56	2,726.15	4,036.76
	514	134.61	40.86	203.65	101.55	83.42	223.02
	516	74.26	73.35	199.06	122.64	249.95	336.13
	517	3,499.07	4,820.64	3,514.42	4,910.51	4,378.18	4,394.10
	518	49.00	82.65	80.14	52.09	101.80	65.00
	519	42.69	106.07	57.86	83.01	96.52	68.93
	521	7,066.94	7,205.81	4,420.95	5,724.41	6,517.25	7,327.22
	523	548.85	455.37	404.81	284.01	324.73	314.50
	524	980.48	482.36	355.11	318.01	399.14	572.23
EBS Total		16,890.22	18,189.41	13,277.48	17,068.06	18,060.90	20,583.23
BSAI Total		17,747.37	19,317.86	14,079.84	18,876.53	20,570.46	21,278.69

Table 5. Estimated catch (t) of all skate species combined by target fishery (upper) and reporting area (lower) 2003-2008. Source: AKRO CAS. *2008 data complete as of October 3, 2008.

region	target	2003	2004	2005	2006	2007	*2008
EBS	Atka	20	35	22	8	26	7
	P cod	14,954	18,000	18,975	14,459	12,713	8,867
	flatfish	3,085	2,613	2,546	3,220	3,462	2,891
	pollock	471	843	731	1,306	1,299	2,381
	rockfish	11	6	4	3	3	1
	sablefish	2	2	2	13	18	9
	other target	220	91	25	26	56	57
	EBS total	**18,764**	**21,591**	**22,305**	**19,034**	**17,578**	**14,213**
AI	Atka	74	108	118	133	127	97
	P cod	200	486	406	417	633	284
	flatfish	254	247	100	188	100	493
	pel pollock	0	0	0	<1	<1	<1
	rockfish	61	16	26	22	69	51
	sablefish	55	8	24	108	38	25
	other target	11	6	3	89	13	3
	AI total	**655**	**871**	**677**	**958**	**980**	**953**

	area	2003	2004	2005	2006	2007	*2008
EBS	508	0	<1	0	0	0	0
	509	2,009	2,170	3,226	3,335	3,572	2,670
	512	25	205	15	0	0	1
	513	2,785	2,883	4,007	2,663	2,353	1,674
	514	281	67	196	221	445	64
	516	132	417	239	252	395	281
	517	3,038	3,046	3,656	2,389	2,148	1,869
	518	25	7	3	8	1	389
	519	199	139	103	65	106	91
	521	8,948	10,310	8,467	8,334	7,088	5,733
	523	307	323	244	279	334	178
	524	1,016	2,025	2,149	1,490	1,136	1,264
	EBS total	**18,764**	**21,591**	**22,305**	**19,034**	**17,578**	**14,213**
AI	541	302	472	472	562	327	363
	542	234	260	124	329	391	430
	543	118	139	82	67	263	160
	AI total	**655**	**871**	**677**	**958**	**980**	**953**

Table 6. Observed skate catch and percent retained by species, and by region, 2003-2007. *2007 reported as of October 15, 2007 (not a complete year). Source: North Pacific Groundfish Observer Program database.

	2003		2004		2005		2006		2007	
Species	**Obs Catch (t)**	**% Retained**	**Obs Catch (t)**	**% Retained**	**Obs Catch (t)**	**% Retained**	**Obs Catch (t)**	**% Retained**	**Obs Catch (t)**	**% Retained**
Alaska	1,179	49%	4,373	36%	4,125	39%	4,956	36%	4,076	32%
Aleutian	71	28%	264	36%	304	31%	154	43%	119	28%
Bathyraja UnID	58	77%	77	8%	6,319	37%	4,586	29%	3,233	23%
Bering	43	27%	233	12%	197	10%	128	17%	79	21%
Big	26	60%	131	27%	165	19%	179	27%	84	46%
Commander	2	1%	15	18%	26	5%	16	5%	21	16%
Longnose	1	32%	15	42%	5	44%	2	48%		0%
Mud			29	7%	22	4%	6	20%	13	7%
Raja UnID					10	4%				0%
Roughtail			5	8%	2	2%	5	12%	2	3%
Skate UnID	13,024	38%	8,822	27%	3,853	28%	2,819	26%	510	14%
Whiteblotched	9	1%	153	21%	58	24%	92	28%	39	28%
Whitebrow			5	31%	7	7%	3	22%	2	21%
Other	2	1%	0	2%	0	100%	0	67%	2	14%
Total	**14,416**	**39%**	**14,123**	**30%**	**15,092**	**34%**	**12,947**	**31%**	**8,181**	**27%**

	2003		2004		2005		2006		2007	
Region	**Obs Catch (t)**	**% Retained**	**Obs Catch (t)**	**% Retained**	**Obs Catch (t)**	**% Retained**	**Obs Catch (t)**	**% Retained**	**Obs Catch (t)**	**% Retained**
AI	437	18%	590	21%	463	17%	690	21%	406	34%
EBS	13,978	39%	13,533	30%	14,629	35%	12,258	32%	7,775	27%
Total	14,416	39%	14,123	30%	15,092	34%	12,947	31%	8,181	27%

Table 7. EBS shelf bottom trawl survey estimates of Alaska skate (*Bathyraja parmifera*) biomass (metric tons). Line indicates the start year of the model. Estimates and CVs in bold (1999-2008) were obtained directly from trawl survey data when species identification was reliable. Estimates and CVs prior to 1999 were partitioned using species composition data from 1999-2008.

year	biomass	CV
1982	167,826	0.10
1983	163,970	0.10
1984	190,037	0.10
1985	158,860	0.10
1986	255,409	0.10
1987	334,132	0.10
1988	392,645	0.10
1989	395,370	0.10
1990	513,751	0.10
1991	433,529	0.10
1992	379,682	0.10
1993	370,356	0.10
1994	412,663	0.10
1995	385,126	0.10
1996	426,649	0.10
1997	402,720	0.10
1998	352,101	0.10
1999	**353,197**	**0.16**
2000	**314,565**	**0.06**
2001	**415,549**	**0.06**
2002	**411,156**	**0.06**
2003	**373,520**	**0.05**
2004	**435,061**	**0.05**
2005	**548,010**	**0.05**
2006	**438,307**	**0.05**
2007	**479,633**	**0.07**
2008	**362,127**	**0.06**

Table 8. Alaska skate EBS shelf survey length compositions, 2000-2008. Bin number is the lower limit of each 4 cm length bin; data are proportions of each bin. N = sample size.

	year								
bin	2000	2001	2002	2003	2004	2005	2006	2007	2008
4	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
8	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
12	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
16	0.000	0.000	0.001	0.000	0.000	0.000	0.000	0.000	0.000
20	0.007	0.010	0.010	0.006	0.006	0.004	0.003	0.007	0.004
24	0.037	0.039	0.023	0.033	0.016	0.019	0.024	0.017	0.017
28	0.047	0.056	0.031	0.031	0.030	0.025	0.026	0.019	0.020
32	0.047	0.058	0.037	0.047	0.032	0.033	0.031	0.028	0.027
36	0.052	0.048	0.044	0.053	0.032	0.041	0.036	0.042	0.038
40	0.051	0.048	0.042	0.048	0.049	0.047	0.045	0.051	0.052
44	0.046	0.051	0.044	0.052	0.051	0.061	0.052	0.052	0.061
48	0.052	0.044	0.041	0.053	0.058	0.057	0.057	0.062	0.061
52	0.056	0.049	0.045	0.041	0.062	0.054	0.051	0.052	0.065
56	0.052	0.043	0.037	0.036	0.051	0.057	0.057	0.055	0.062
60	0.055	0.051	0.035	0.041	0.047	0.064	0.055	0.050	0.059
64	0.045	0.043	0.033	0.039	0.042	0.053	0.053	0.060	0.060
68	0.035	0.047	0.041	0.043	0.049	0.046	0.048	0.055	0.046
72	0.038	0.046	0.035	0.041	0.043	0.048	0.049	0.053	0.051
76	0.030	0.035	0.041	0.042	0.047	0.040	0.048	0.046	0.049
80	0.040	0.030	0.035	0.047	0.038	0.040	0.037	0.041	0.043
84	0.030	0.026	0.046	0.037	0.043	0.039	0.044	0.040	0.040
88	0.034	0.033	0.069	0.044	0.044	0.052	0.038	0.045	0.043
92	0.051	0.060	0.092	0.056	0.062	0.048	0.062	0.058	0.055
96	0.070	0.071	0.094	0.088	0.081	0.062	0.068	0.063	0.057
100	0.066	0.069	0.076	0.065	0.072	0.059	0.064	0.060	0.055
104	0.043	0.031	0.037	0.043	0.034	0.035	0.038	0.028	0.026
108	0.014	0.011	0.010	0.013	0.010	0.011	0.012	0.011	0.009
112	0.002	0.002	0.003	0.002	0.003	0.002	0.002	0.003	0.001
116	0.000	0.000	0.000	0.000	0.000	0.001	0.000	0.001	0.000
120	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
124	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
128	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
132	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
N	316	354	333	332	380	370	352	362	346

Table 9. Partitioned Alaska skate catch estimates (metric tons) based on observed catch data and survey species composition. Total BSAI catch estimates for each fishery (right-most column) were used in the SS2 base model. Because 2008 catch data are incomplete, 2007 catch was used for 2008.

Year	EBS shelf Longline	EBS shelf Trawl	EBS slope Longline	EBS slope Trawl	AI Longline	AI Trawl	BSAI Longline	BSAI Trawl
1992	12,239	2,698	23	8	166	92	12,428	2,798
1993	8,822	1,945	16	6	119	67	8,958	2,017
1994	10,263	2,262	19	7	139	77	10,421	2,346
1995	10,746	2,369	20	7	145	81	10,911	2,457
1996	9,123	2,011	17	6	123	69	9,263	2,086
1997	12,907	2,845	24	8	150	84	13,081	2,937
1998	13,900	3,064	26	9	198	110	14,123	3,183
1999	9,703	2,139	19	7	141	78	9,862	2,224
2000	12,744	2,809	24	9	388	216	13,157	3,034
2001	13,973	3,080	26	9	440	245	14,438	3,334
2002	15,776	3,477	119	42	138	77	16,033	3,596
2003	13,718	3,218	30	8	102	77	13,850	3,302
2004	16,591	3,892	27	23	148	61	16,766	3,975
2005	17,673	3,366	40	4	115	70	17,827	3,441
2006	14,736	3,248	27	10	153	85	14,916	3,343
2007	13,676	3,015	25	9	172	96	13,873	3,119
2008	13,676	3,015	25	9	172	96	13,873	3,119

Table 10. Alaska skate length compositions from the EBS 2007 longline and trawl fisheries. Bin number is the lower limit of each 4 cm length interval.

	2007	
bin	**longline**	**trawl**
4	0.000	0.000
8	0.000	0.000
12	0.000	0.000
16	0.000	0.001
20	0.000	0.008
24	0.000	0.018
28	0.000	0.014
32	0.000	0.024
36	0.000	0.030
40	0.002	0.040
44	0.005	0.057
48	0.006	0.061
52	0.016	0.052
56	0.027	0.047
60	0.046	0.060
64	0.063	0.066
68	0.053	0.048
72	0.072	0.053
76	0.055	0.057
80	0.059	0.044
84	0.060	0.046
88	0.065	0.058
92	0.089	0.051
96	0.118	0.058
100	0.137	0.055
104	0.080	0.027
108	0.031	0.013
112	0.010	0.008
116	0.006	0.004
120	0.002	0.001
124	0.001	0.000
128	0.000	0.000
132	0.000	0.000
N	251	215

Table 11. Final parameter values of the base model. Where parameters were estimated freely within the model, minimum and maximum bounds are shown.

parameter		value	min	max	fix?
growth and natural mortality	natural mortality (M)	0.13			X
	length at A1 (L1)	16.4	10	30	
	length at A2 (L2)	101.6	70	120	
	von Bertalanffy coefficient (k)	0.153	0.05	0.2	
	CV of L1	0.288	0	0.5	
	ln CV of L2	-1.691	-3	1	
length-weight relationship	coefficient (a)	4.0×10^6			X
	exponent (b)	3.149			X
length at maturity	length at 50% maturity (a)	93.28			X
	slope (b)	-0.548			X
weight-fecundity relationship	coefficient (a)	0.5			X
	exponent (b)	0			X
stock-recruit relationship	ln virgin recruitment level (R0)	10.63	5	15	
	steepness (h)	1			X
	SD of R0 (σR)	0.4			X
EBS shelf survey catchability	ln catchability (Q)	0			X
longline length selectivity	peak (p1)	103.2	7.6	126	
	top (p2)	1.71	-6	4	
	ascending width (p3)	3.84	-1	9	
	descending width (p4)	5.07	-1	9	
	selectivity at first size bin (p5)	-1.80	-5	9	
	selectivity at last size bin (p6)	9.0			X
trawl length selectivity	peak (p1)	48.6	7.6	126.2	
	top (p2)	1.87	-6	4	
	ascending width (p3)	4.52	-1	9	
	descending width (p4)	6.44	-1	9	
	selectivity at first size bin (p5)	-0.609	-5	9	
	selectivity at last size bin (p6)	9	-5	9	X
survey length selectivity (logistic)	(p1)	-32.99			X
	(p2)	285.9			X
longline age selectivity (logistic)	(p1)	6.5			X
	(p2)	0.547	0	30	
trawl age selectivity (logistic)	(p1)	3.5			X
	(p2)	0.076	0	30	
survey age selectivity (logistic)	(p1)	4			X
	(p2)	0.188	0	30	
initial fishing mortality	longline fishery F	0.042	0.001	1	
	trawl fishery F	0.005	0.0004	1	

Table 12. Estimates of M based on Alaska skate life history parameters from Matta (2006). "Age mature" (T_{mat}) was given a range to estimate M by the Rikhter and Efanov method to account for uncertainty in this parameter.

Sex	Hoenig	T_{mat}	Rikhter & Efanov	Alverson & Carney	Charnov	Roff	Jensen k	Jensen T_{50}
males	0.28			0.37	0.22	**0.13**	0.19	0.18
females	0.25			0.35	0.16	0.15	0.14	0.17
both		8	0.19					
		9	0.16					
		10	**0.13**					

Table 13. Overall and component likelihoods.

overall likelihood	**246.8**
survey	-14.08
length compositions	131.5
length at age	170.6
equilibrium catch	0.00136
recruitment	-30.28
forecast recruitment	-11.0

Table 14. Time series of total (age 0+) biomass (metric tons), spawning biomass (metric tons) and the number of age 0 recruits (thousands of fish) predicted by the base model.

	total biomass (t)	female spawning biomass (t)	recruits (1000s)
1992	482,909	106,643	30,241
1993	493,992	104,411	36,861
1994	508,334	104,095	41,539
1995	518,551	104,160	41,570
1996	525,003	105,864	47,993
1997	530,193	109,527	49,978
1998	528,812	113,013	49,184
1999	525,420	116,538	57,946
2000	527,163	121,736	39,851
2001	526,321	124,082	48,257
2002	525,997	124,788	53,862
2003	526,363	123,631	38,982
2004	531,447	122,124	29,117
2005	534,239	119,369	36,532
2006	537,499	116,691	36,532
2007	544,135	115,278	36,532
2008	550,840	115,307	36,532

Table 15. Time series of exploitation rates (catch/total biomass) as estimated by the model.

year	longline	trawl	total
1992	0.026	0.006	0.032
1993	0.018	0.004	0.022
1994	0.021	0.005	0.025
1995	0.021	0.005	0.026
1996	0.018	0.004	0.022
1997	0.025	0.006	0.030
1998	0.027	0.006	0.033
1999	0.019	0.004	0.023
2000	0.025	0.006	0.031
2001	0.027	0.006	0.034
2002	0.030	0.007	0.037
2003	0.026	0.006	0.033
2004	0.032	0.007	0.039
2005	0.033	0.006	0.040
2006	0.028	0.006	0.034
2007	0.025	0.006	0.031
2008	0.025	0.006	0.031

Table 16. Summary of major results of the base model and management recommendations for Alaska skates in the BSAI.

Tier	3a
Reference mortality rates	
M	0.13
$F_{35\%}$	0.10
$F_{40\%}$	0.087
Equilibrium spawning biomass (t)	
$B_{35\%}$	65,149
$B_{40\%}$	74,457
$B_{100\%}$	186,142
Projected biomass for 2009 (t)	
Spawning (at max F_{ABC})	108,093
Total	551,439
ABC for 2009	
F_{ABC} (maximum permissible)	0.087
F_{ABC} (recommended)	0.087
ABC (t; maximum permissible)	25,854
ABC (t; recommended)	25,854
Overfishing level for 2009	
F_{OFL}	0.10
OFL (t)	30,077
Projections for 2010	
Spawning biomass (t; at max F_{ABC})	108,089
Total biomass (t)	549,000
ABC (t; maximum permissible)	25,796
OFL (t)	30,009

Table 17. Total skate biomass (metric tons) with coefficient of variation (cv) from bottom trawl surveys of the Eastern Bering Sea (EBS) shelf, EBS slope, and Aleutian Islands (AI), 1975-2008.

Year	EBS shelf		EBS slope		AI	
	biomass	cv	biomass	cv	Biomass	cv
1975	24,349	0.19				
1976						
1977						
1978						
1979	58,147	0.14	3,056	0.26		
1980					4,257	0.25
1981			2,743	0.12		
1982	164,084	0.10	2,723	0.10		
1983	161,329	0.09			9,683	0.12
1984	186,976	0.09				
1985	149,573	0.11	3,329	0.10		
1986	251,296	0.15			15,436	0.19
1987	346,679	0.10				
1988	408,242	0.11	3,271	0.21		
1989	406,007	0.08				
1990	533,837	0.11				
1991	448,054	0.09	4,031	0.25	14,967	0.17
1992	390,294	0.09				
1993	374,882	0.07				
1994	414,054	0.08			25,014	0.10
1995	391,537	0.08				
1996	403,521	0.06				
1997	391,032	0.07			28,922	0.14
1998	354,000	0.05				
1999	348,477	0.16				
2000	325,292	0.06			29,320	0.09
2001	420,313	0.06				
2002	366,315	0.07	69,275	0.50	34,413	0.11
2003	386,339	0.05				
2004	416,559	0.05	33,156	0.08	53,071	0.16
2005	481,194	0.05				
2006	442,556	0.05			54,210	0.12
2007	475,024	0.07				
2008	381,744	0.05	37,548	0.08		

Table 18. Estimates of M for the Other Skates group based on *Raja* sp. life history parameters. "Age mature" (T_{mat}) was given a range for M estimates by the Rikhter and Efanov method to account for uncertainty in this parameter. Study areas are indicated as CA (California), GOA (Gulf of Alaska), and BC (British Columbia. Life history parameter sources: Zeiner and Wolf 1993, Gburski et al. 2007, McFarlane and King 2006.

Species	Area	Sex	Hoenig	T_{mat}	Rikhter & Efanov	Alverson & Carney	Charnov	Roff
Big skate	CA	*males*	0.38					
	CA	*females*	0.35					
	CA	*both*		8	0.19			
	CA			9	0.16			
	CA			10	0.13			
	CA			11	0.12			
	CA			12	**0.10**			
	GOA	*males*	0.28			0.33	0.28	
	GOA	*females*	0.30			0.45	0.15	
	BC	*males*	0.17			0.25	0.10	0.34
	BC	*females*	0.16			0.25	0.08	0.27
	BC	*both*		5	0.32			
	BC			6	0.26			
	BC			7	0.22			
	BC			8	0.19			
Longnose skate	CA	*males*	0.32			0.31	0.44	0.23
	CA	*females*	0.35			0.45	0.29	0.03
	CA	*both*		7	0.22		0.31	
	CA			8	0.19			
	CA			9	0.16			
	CA			10	0.13			
	GOA	*males*	0.17			0.24	0.11	
	GOA	*females*	0.17			0.28	0.07	
	BC	*males*	0.18			0.25	0.13	0.21
	BC	*females*	0.16			0.22	0.11	0.12
	BC	*both*		6	0.26			
	BC			7	0.22			
	BC			8	0.19			
	BC			9	0.16			
	BC			10	0.13			

Figures

Figure 1. Skate diversity on the Bering Sea slope: five species of skate captured in a single trawl haul on the NMFS Bering sea slope survey, 2002. Species pictured include whitebrow skate (*B. minispinosa*), mud skate (*B. taranetzi*), whiteblotched skate (*B. maculata*), Aleutian skate (*B. aleutica*), and Commander skate (*B. lindbergi*). Photo credit: Gerald Hoff.



Figure 2. Distribution of skate species in Alaskan waters. These maps were created primarily using survey data, although observer records were included whenever positive species identification was possible (through voucher specimens or photographs). ***(Source: Stevenson et al. 2007)***



Figure 2 continued. Distribution of skate species in Alaskan waters. (Source: Stevenson et al. 2007)



Figure 3. Skate species composition (by weight) by habitat area, EBS shelf (upper left), EBS slope (upper right), and AI (bottom). EBS shelf data are from 2007 bottom trawl survey; AI data are from 2006 bottom trawl survey; EBS slope data are from 2004 bottom trawl survey.



Figure 4. Relative abundance of skate species in the EBS by depth. (Source: Stevenson et al. 2006.)



Figure 5. Average survey CPUE, Alaska skate (*Bathyraja parmifera*), 2001-2004. Data are from RACE Bering Sea Groundfish Surveys and are shown as average CPUE (kg/ha) for each station.



Figure 6. Average survey CPUE, Bering skate (*Bathyraja interrupta*), 2001-2004. Data are from RACE Bering Sea Groundfish Surveys and are shown as average CPUE (kg/ha) for each station.



Figure 7. Skate distribution in the AI from NMFS bottom trawl surveys. Specimens of *B. parmifera* in the western AI have now been described as a new species (see below).



Figure 8. Skate diversity in the Aleutians: a new species, the leopard skate, from the Aleutian Islands (left) formerly thought to be the same species as the extremely common Alaska skate, *B. parmifera* (from the EBS, right). Photo credits: leopard skate, Richard MacIntosh; Alaska skate, Beth Matta.



Figure 9. Map of the eastern Bering Sea with the six known skate nursery site locations and designations as a northern or southern nursery site. (See the legend for nursery site designation.) Source: Gerald Hoff, AFSC, unpublished data.



Figure 10. Embryo length composition data used in a cohort analysis of embryo development time. Figure is from G. Hoff (pers. comm.).



Figure 11. Ocean temperature versus embryo development time for 21 skate species. Dark grey circle is the Alaska skate. Equation and R^2 are the values of the fitted relationship. Figure is from G. Hoff, AFSC, pers. comm.



Figure 12. Identification of observed incidentally caught skates in AI (left) and EBS (right) groundfish fisheries, 2004 (top) and 2005 (bottom). Source: AFSC Fishery Monitoring and Analysis program database.



Figure 12 continued. Identification of observed incidentally caught skates in AI (left) and EBS (right) groundfish fisheries, 2006 (top) and 2007 (bottom). Source: North Pacific Groundfish Observer Program database. 2007 data are reported through October 15, 2007.



Figure 13. Total skate catch (all species combined) by FMP reporting area for both the EBS and the AI, 2003-2005. Source: AKRO CAS.



Figure 14. Relative proportion of Alaska skates and Other Skates in each habitat area. Graphs represent weighted averages from 1999-2008 trawl survey biomass estimates. These data were used to reconstruct catch data for the Alaska skate.



Figure 15. Estimated catch of Alaska skates (t) in the BSAI from 1992 to 2007. Data were obtained from the Blend system and AKRO CAS. Data from 2008 are incomplete and not shown.



Fig. 16. Fishery length compositions by quarter (unbinned data) for Alaska skates during 2007.



Figure 17. Observed size at age data from Alaska skates collected in the 2003 EBS shelf trawl survey, sexes combined (n=182). The three year embryonic development period included in the base model is represented by the shaded area.



Figure 18. Observed size at age data from Alaska skates collected in the 2005 longline fishery, sexes combined (n=208). The three year embryonic development period included in the base model is represented by the shaded area.



Figure 19. The relationship between total length (TL) and total body weight (W) for the Alaska skate, both sexes combined (n=526).



Figure 20. Simplified schematic depiction of population dynamics model used in the Alaska skate assessment. Blue diamonds indicate physical quantities, red circles indicate rates. R_a = recruitment in year a, M = natural mortality, SSB = spawning biomass, BH indicates that a Beverton-Holt stock-recruit relationship is applied to SSB to estimate recruitment.



Figure 21. Female Alaska skate maturity-at-length data shown with fitted logistic curve from Matta (2006) (n=642).



Figure 22. Length-based capture probability for skates (*Bathyraja* spp.) in the EBS shelf bottom trawl survey, based on data from Kotwicki and Weinberg (2005).



Figure 23. Observed biomass (circles) from EBS shelf surveys 1992-2008, with approximate confidence intervals (± 2 SE), and predicted survey biomass from the model (black line).



Figure 24. EBS shelf survey length compositions from 2000-2008. Grey bars = observed proportions; black line with circles = model predictions. Red column indicates 96-99 cm length bin.



Figure 24 continued. EBS shelf survey length compositions from 2000-2008. Grey bars = observed proportions; black line with circles = model predictions. Red column indicates 96-99 cm length bin.



Figure 24 continued. EBS shelf survey length compositions from 2000-2008. Grey bars = observed proportions; black line with circles = model predictions. Red column indicates 96-99 cm length bin.





Figure 25. Observed and model-predicted length compositions from the 2007 longline (top) and trawl (bottom) fisheries, with model predictions. Grey bars = observed values, black line with circles = predicted values. Red column indicates the 96-99 cm length bin.



Figure 26. Observed and model-predicted length-at-age from the 2003 (upper panel) and 2007 (lower panel EBS shelf survey.



Figure 27. Observed and model-predicted length-at-age from the 2005 longline fishery.



Figure 28. Observed length-at-age from the three datasets used in the model, with the population estimate of length-at-age (blue line) superimposed.



Figure 29. Length selectivities of the longline fishery, trawl fishery, and EBS shelf trawl survey.



Figure 30. Time series of expected recruitment (in thousands of age 0 fish), with the time series of individual year class estimates predicted by the model and the expected Beverton-Holt stock-recruit relationship with a steepness of 1.0.



Figure 31. Relationship between the number of age 0 recruits (in thousands of fish) and female spawning biomass (t). Time series of individual year class estimates from SS2 is shown with a Beverton-Holt stock-recruit relationship with a steepness of 1.0.



Figure 32. Likelihood profile analyses for natural mortality (M; upper panel) and age at 50% selectivity for the trawl survey (lower panel). Yellow circles indicates values used in the model.



Figure 33. Observed survey biomass (grey circles) and model fit (black line) produced when age at 50% selectivity for the survey is fixed at 4.5 years.



Figure 34. Likelihood-profile analysis for sigma R. Note that survey likelihood is on a separate axis.





Figure 35. Examples of improved fit to survey length composition (upper panel) and length-at-age (lower panel) data produced by the alternative model.



Figure 36. Selectivity curves produced by the alternative model.



Figure 37. Population estimate of length-at-age (blue line) produced by the alternative model superimposed on observed length-at-age from the three datasets used in the model.



Figure 38. Time series of model estimates for total (age 0+) biomass (t) and female spawning biomass (t).



Figure 39. Aggregated skate biomass (metric tons) estimated from RACE bottom trawl surveys in each of the three major habitat areas (1982 – 2008).



Figure 40. EBS (upper panel) and AI (lower panel) skate food webs derived from mass balance ecosystem models, with skate species aggregated in each area. (Source: K. Aydin, AFSC, code available upon request.)



Figure 41. Comparative density (upper panels) and exploitation rate (lower panels) of Alaska (left panels) and all other *Bathyraja* (right panels) skates in the AI, EBS, and GOA (early 1990s, before fishery in GOA). (Alaska skates are a very small component of skate biomass in the GOA, and are therefore not modeled separately.) Note that the Other skates plot does not include the most common species in that region, the big skate and longnose skate—see the GOA skate SAFE for information on those skates. Biomass density plots are from trawl survey data; exploitation rate plots are derived from catch and biomass estimates and from assumed estimates of skate productivity (approximated from Frisk et al. 2001).



Figure 42. Mortality sources and consumption of skates in the EBS—mortality pie (upper panels) and estimates of annual consumption by predators (lower panels) for EBS Alaska skates (left panels) and all other EBS skates (right panels). Model outputs were derived from diet compositions, production rates, and consumption rates of skate predators, and from skate catch data.



Figure 43. Mortality sources and consumption of skates in the AI—mortality pie (upper panels) and estimates of annual consumption by predators (lower panels) for AI (former) Alaska skate (left panels) and AI Other Skates (right panels). Model outputs were derived from diet compositions, production rates, and consumption rates of skate predators, and from skate catch data.



Figure 44. Diet composition (upper panels) and annual estimated prey consumption by skates (lower panels) for EBS Alaska skates (left panels) and Other Skates (right panels). Results were generated from stomach content collections occurring during RACE trawl surveys.



Figure 45. Diet composition (upper panels) and annual estimated prey consumption by skates (lower panels) for AI Alaska skates (left panels) and Other Skates (right panels). Consumption rates were estimated using published diet data from the Kuril Islands (Orlov 1998, 1999) and estimated prey densities.





Figure 46. Diet composition (by weight) for the other two biomass-dominant skate species in the Aleutian Islands (which are included in the "Other Skates" group in the previous figure): whiteblotched skate (top) and Aleutian skate (bottom). Results were generated from stomach content collections occurring during trawl surveys, and are described in more detail in Yang (2007).