

1. Assessment of the Walleye Pollock Stock in the Gulf of Alaska

Martin Dorn, Kerim Aydin, Steven Barbeaux, Michael Guttormsen,
Bernard Megrey, Kally Spalinger, and Mark Wilkins

Alaska Fisheries Science Center
National Marine Fisheries Service

Executive Summary

Summary of major changes

Relative to last year's assessment, the following changes have been made in the current assessment.

New Input data

1. Fishery: 2008 total catch and catch at age.
2. Shelikof Strait EIT survey: 2009 biomass and age composition.
3. NMFS bottom trawl survey: 2009 biomass and length composition.
3. ADF&G crab/groundfish trawl survey: 2009 biomass and length composition, 2008 age composition.

Assessment model

The age-structured assessment model developed using ADModel Builder (a C++ software language extension and automatic differentiation library) and used for assessments in 1999-2008 was used again for this year's assessment.

Assessment results

The model estimate of spawning biomass in 2010 is 184,567 t, which is 29.8% of unfished spawning biomass (based on average post-1977 recruitment) and below $B_{40\%}$ (248,000 t). New NMFS bottom trawl, ADF&G crab/groundfish and Shelikof Strait EIT surveys were conducted in 2009, and all showed increases in biomass, although the magnitude of the increases were not consistent between surveys. Other spawning aggregations surveyed acoustically in winter of 2009 also increased, but remained close to historical low levels, with the exception of the Chirikof area, where only trace quantities of spawning pollock were detected. Spawning biomass is projected to increase in 2010 and in subsequent years. The initial estimate of the 2007 year class is 1.7 times average recruitment, but the author's ABC recommendation is based on setting the 2007 year class equal to the post-1977 average due to uncertainty in this estimate.

The author's 2010 ABC recommendation for pollock in the Gulf of Alaska west of 140° W lon. (W/C/WYK) is 77,150 t, an increase of 78% from the 2009 ABC. This recommendation is based on an average 2007 year class, and a more conservative alternative to the maximum permissible F_{ABC} introduced in the 2001 SAFE. The OFL in 2010 is 103,210 t. In 2011, the recommended ABC and OFL are 101,510 t and 135,010 t, respectively.

For pollock in southeast Alaska (East Yakutat and Southeastern areas), the ABC recommendations for 2010 and 2011 in Appendix A are 9,245 t and the OFL is 12,326 t (the same for both years). These

recommendations are based on estimated biomass in the southeast Alaska from the 2009 NMFS bottom trawl survey.

Summary

Natural mortality = 0.3

Tier: 3b

2010 harvests

Maximum permissible ABC:	$F_{40\%} \text{ (adjusted)} = 0.17$	Yield = 89,800 t
Recommended ABC:	$F_{40\%} \text{ (author's adjusted)} = 0.14$	Yield = 77,150 t
Overfishing (OFL):	$F_{35\%} \text{ (adjusted)} = 0.19$	Yield = 103,210 t

2011 harvests

Maximum permissible ABC:	$F_{40\%} \text{ (adjusted)} = 0.18$	Yield = 117,750 t
Recommended ABC:	$F_{40\%} \text{ (author's adjusted)} = 0.16$	Yield = 101,510 t
Overfishing (OFL):	$F_{35\%} \text{ (adjusted)} = 0.21$	Yield = 135,010 t

Equilibrium female spawning biomass

$B_{100\%} = 620,000$ t

$B_{40\%} = 248,000$ t

$B_{35\%} = 217,000$ t

Projected 2010 biomass

Age 3+ biomass = 756,550 t

Female spawning biomass = 184,567 t

Responses to Comments of the Scientific and Statistical Committee (SSC)

There were no comments in the December 2008 minutes that were specific to the Gulf of Alaska pollock assessment. As a general recommendation for all Tier 3 and lower stocks, the SSC recommended that the probability of the spawning stock biomass falling below B20% be estimated for a 3-5 year projection. These estimates have been routinely provided in the Gulf of Alaska pollock assessment since 2002.

Introduction

Walleye pollock (*Theragra chalcogramma*) is a semi-pelagic schooling fish widely distributed in the North Pacific Ocean. Pollock in the Gulf of Alaska are managed as a single stock independently of pollock in the Bering Sea and Aleutian Islands. The separation of pollock in Alaskan waters into eastern Bering Sea and Gulf of Alaska stocks is supported by analysis of larval drift patterns from spawning locations (Bailey et al. 1997), genetic studies of allozyme frequencies (Grant and Utter 1980), mtDNA variability (Mulligan et al. 1992), and microsatellite allele variability (Bailey et al. 1997).

The results of studies of stock structure in the Gulf of Alaska are equivocal. There is evidence from allozyme frequency and mtDNA that spawning populations in the northern part of the Gulf of Alaska (Prince William Sound and Middleton Island) may be genetically distinct from the Shelikof Strait spawning population (Olsen et al. 2002). However significant variation in allozyme frequency was found between Prince William Sound samples in 1997 and 1998, indicating a lack of stability in genetic structure for this spawning population. Olsen et al. (2002) suggest that interannual genetic variation may be due to variable reproductive success, adult philopatry, source-sink population structure, or utilization of the same spawning areas by genetically distinct stocks with different spawning timing. Peak spawning at the two major spawning areas in the Gulf of Alaska occurs at different times. In the Shumagin Island area, peak spawning apparently occurs between February 15- March 1, while in Shelikof Strait peak spawning occurs later, typically between March 15 and April 1. It is unclear whether the difference in timing is genetic, or a response to differing environmental conditions in the two areas.

Fishery

The commercial fishery for walleye pollock in the Gulf of Alaska started as a foreign fishery in the early 1970s (Megrey 1989). Catches increased rapidly during the late 1970s and early 1980s (Table 1.1). A large spawning aggregation was discovered in Shelikof Strait in 1981, and a fishery developed for which pollock roe was an important product. The domestic fishery for pollock developed rapidly in the Gulf of Alaska with only a short period of joint venture operations in the mid-1980s. The fishery was fully domestic by 1988.

The fishery for pollock in the Gulf of Alaska is entirely shore-based with approximately 90% of the catch taken with pelagic trawls. During winter, fishing effort targets pre-spawning aggregations in Shelikof Strait and near the Shumagin Islands (Fig. 1.1). Fishing in summer is less predictable, but typically occurs on the east side of Kodiak Island and in nearshore waters along the Alaska Peninsula.

Incidental catch in the Gulf of Alaska directed pollock fishery is low. For tows classified as pollock targets in the Gulf of Alaska between 2004 and 2008, about 94% of the catch by weight consisted of pollock (Table 1.2). Nominal pollock targets are defined by the dominance of pollock in the catch, and may include tows where other species were targeted, but where pollock were caught instead. The most common managed species in the incidental catch are arrowtooth flounder, Pacific cod, flathead sole, Pacific ocean perch, miscellaneous flatfish, and the shorttraker/rougheye rockfish complex. The most common non-target species are squid, eulachon, various shark species (e.g., Pacific sleeper sharks, spiny dogfish, salmon shark), and grenadiers. Bycatch estimates for prohibited species over the period 2004-2008 are given in Table 1.3.

Kodiak is the major port for pollock in the Gulf of Alaska, with 63% of the 2004-2008 landings. In the western Gulf of Alaska, Sand Point, Dutch Harbor, King Cove, and Akutan are important ports, sharing 37% of 2004-2008 landings. Secondary ports, including Alitak Bay, Homer, Ninilchik, Seward, and Sitka account for less than 1% of the 2004-2008 landings.

Since 1992, the Gulf of Alaska pollock TAC has been apportioned spatially and temporally to reduce potential impacts on Steller sea lions. The details of the apportionment scheme have evolved over time, but the general objective is to allocate the TAC to management areas based on the distribution of surveyed biomass, and to establish three or four seasons between mid-January and autumn during which some fraction of the TAC can be taken. The Steller Sea Lion Protection Measures implemented in 2001 established four seasons in the Central and Western GOA beginning January 20, March 10, August 25, and October 1, with 25% of the total TAC allocated to each season. Allocations to management areas 610, 620 and 630 are based on the seasonal biomass distribution as estimated by groundfish surveys. In addition, a new harvest control rule was implemented that requires suspension of directed pollock fishing when spawning biomass declines below 20% of the reference unfished level.

Data Used in the Assessment

The data used in the assessment model consist of estimates of annual catch in tons, fishery age composition, NMFS summer bottom trawl survey estimates of biomass and age composition, echo integration trawl (EIT) survey estimates of biomass and age composition in Shelikof Strait, egg production estimates of spawning biomass in Shelikof Strait, ADF&G bottom trawl survey estimates of biomass and length and age composition, and historical estimates of biomass and length and age composition from surveys conducted prior to 1984 using a 400-mesh eastern trawl. Binned length composition data are used in the model only when age composition estimates are unavailable, such as the fishery in the early part of the modeled time period and the most recent survey. The FOCI year class prediction is used qualitatively along with other information to evaluate the likely strength of incoming year classes.

Total Catch

Estimated catch was derived by the NMFS Regional Office from shoreside electronic logbooks and observer estimates of at-sea discards (Table 1.4). Increases in discard in 2008 (primarily in area 610) are pollock taken as bycatch in Pacific cod fishery, and are not from the directed pollock fishery. Catches include the state-managed pollock fishery in Prince William Sound. Since 1996 the pollock Guideline Harvest Level (GHL) for the PWS fishery has been deducted from the Acceptable Biological Catch (ABC) by the NPFMC Gulf of Alaska Plan Team for management purposes.

Fishery Age Composition

Estimates of fishery age composition were derived from at-sea and port sampling of the pollock catch for length and ageing structures (otoliths). Pollock otoliths collected during the 2008 fishery were aged using the revised criteria described in Hollowed et al. (1995), which involved refinements in the criteria to define edge type. Catch age composition was estimated using methods described by Kimura and Chikuni (1989). Age samples were used to construct age-length keys by sex and stratum. These keys were applied to sex and stratum specific length frequency data to estimate age composition, which were then weighted by the catch in numbers in each stratum to obtain an overall age composition. Age and length samples from the 2008 fishery were stratified by half year and statistical area as follows:

Time strata		Shumagin-610	Chirikof-620	Kodiak-630	W. Yakutat and PWS-640 and 649
1st half (A and B seasons)	No. ages	395	463	252	83
	No. lengths	2137	5299	1620	452
	Catch (t)	6,110	15,576	5,132	1798
2nd half (C and D seasons)	No. ages	251	173	434	---
	No. lengths	1863	1216	3724	---
	Catch (t)	11,146	3,483	9,255	---

In the first half of 2008, the 2000 and 1999 year classes (now age-8 and age-9 fish) were still present as a secondary mode in the fishery age composition, but were much less common in the second half of the year (Fig. 1.2). Age-3 and age-4 fish (2005 and 2004 year classes) were the dominant mode in the fishery age composition in both seasons and in all years. Age-4 fish (2004 year class) were present in significant quantities only in area 620.

Fishery catch at age in 1976-2008 is presented in Table 1.5 (See also Fig. 1.3). Sample sizes for ages and lengths are given in Table 1.6.

Gulf of Alaska Bottom Trawl Survey

Trawl surveys have been conducted by Alaska Fisheries Science Center (AFSC) every three years (beginning in 1984) to assess the abundance of groundfish in the Gulf of Alaska (Table 1.7). Starting in 2001, the survey frequency was increased to every two years. The survey uses a stratified random design, with 49 strata based on depth, habitat, and management area (Martin 1997). Area-swept biomass estimates are obtained using mean CPUE (standardized for trawling distance and mean net width) and stratum area. The survey is conducted from chartered commercial bottom trawlers using standardized poly-Northeastern high opening bottom trawls rigged with roller gear. In a typical survey, 800 tows are completed. On average, 70% of these tows contain pollock (Table 1.8).

The time series of pollock biomass used in the assessment model is based on the surveyed area in the Gulf of Alaska west of 140° W lon., obtained by adding the biomass estimates for the Shumagin, Chirikof, Kodiak INPFC areas, and the western portion of Yakutat INPFC area. Biomass estimates for 1990, 1993, 1996, 1999, 2003, 2005, 2007 and 2009 for the west Yakutat region were obtained by splitting strata and survey CPUE data at 140° W lon. (M. Martin, AFSC, Seattle, WA, pers. comm. 2007). For surveys in 1984 and 1987, the average percent in West Yakutat in the 1990-99 surveys was used. The average was also used in 2001, when West Yakutat was not surveyed.

An adjustment was made to the survey time series to account for unsurveyed pollock in Prince William Sound. This adjustment was derived from an area-swept biomass estimate for PWS from a trawl survey conducted by ADF&G in 1999, using a standard ADF&G 400 mesh eastern trawl. The 1999 biomass estimate for PWS was 6,304 t ± 2,812 t (95% CI) (W. Bechtol, ADF&G, 1999, pers. comm.). The PWS biomass estimate should be considered a minimum estimate because ADF&G survey gear is less effective at catching pollock compared to the triennial survey gear (von Szalay and Brown 2001). For 1999, the biomass estimates for the NMFS bottom trawl survey and the PWS survey were simply added to obtain a total biomass estimate. The adjustment factor for the 1999 survey, (PWS + NMFS)/NMFS, was applied to other triennial surveys, and increased biomass by 1.05%.

The Alaska Fisheries Science Center's (AFSC) Resource Assessment and Conservation Engineering (RACE) Division conducted the eleventh comprehensive bottom trawl survey since 1984 during the summer of 2009. The spatial distribution of pollock shows some differences from previous surveys, with higher CPUEs around the entrance to the Shelikof sea valley and in Amatuli Trench (Fig. 1.4). Other area of higher CPUE was the east side of Kodiak Island, nearshore along the Alaska Peninsula, and immediately north of Dixon Entrance in Southeast Alaska, which are areas where pollock CPUE is typically higher (Fig. 1.4). The 2009 gulfwide biomass estimate of pollock was 703,644 t, representing an increase of 123% from the 2007 gulfwide biomass estimate, and the largest estimate since 1996. The biomass estimate for the portion of the Gulf of Alaska west of 140° W long. is 666,505 t.

Bottom Trawl Age and Length Composition

Estimates of numbers at age from the bottom trawl survey were obtained from random otolith samples and length frequency samples (Table 1.9). Numbers at age were estimated for three strata: Western GOA (Shumagin INPFC area), Central GOA (Chirikof and Kodiak INPFC areas), Eastern GOA (Yakutat and Southeastern INPFC areas) using age-length keys and CPUE-weighted length frequency data. The combined Western and Central age composition was used in the assessment model. Ages are not yet available for the 2009 survey, and instead lengths were used in the assessment model. Length composition by statistical area showed a mode of age-1 fish in all areas that increased in size from the Shumagin area to the Southeast area, most likely due to seasonal growth during the course of the survey (Fig. 1.5). This pattern has been seen in previous bottom trawl surveys. There was a mode of very large fish (~60 cm) in the Shumagin area that was not seen elsewhere. There was a broad length distribution in the central Gulf of Alaska (Chirikof and Kodiak areas) that included both juveniles and adults.

Shelikof Strait Echo Integration Trawl Survey

Echo integration trawl surveys to assess the biomass of pollock in the Shelikof Strait area have been conducted annually since 1981 (except 1982 and 1999). Survey methods and results for 2009 are presented in a NMFS processed report (Guttormsen et. al. in review). Biomass estimates using the Simrad EK echosounder from 1992 onwards were re-estimated to take into account recently published work of eulachon acoustic target strength (Gauthier and Horne 2004). Previously, acoustic backscatter was attributed to eulachon based on the percent composition of eulachon in trawls, and it was assumed that eulachon had the same target strength as pollock. Since Gauthier and Horne (2004) determined that the target strength of eulachon was much lower than pollock, the acoustic backscatter could be attributed entirely to pollock even when eulachon were known to be present. In 2008, the noise-reduced *R/V Oscar Dyson* became the designated survey vessel for acoustic surveys in the Gulf of Alaska. In winter of 2007, a vessel comparison experiment was conducted between the *R/V Miller Freeman* and the *R/V Oscar Dyson*, which obtained an OD/MF ratio of 1.132 in Shelikof Strait.

The 2009 biomass estimate for Shelikof Strait is 265,971 t, an increase of 15% from the 2008 biomass. Biomass of pollock ≥ 43 cm (a proxy for spawning biomass) increased by 60% from the 2007 estimate, apparently due to above average recruitment to the spawning population (Fig. 1.6).

Additional EIT surveys in winter 2009 covered the Shumagin Islands spawning area, Sanak Gully, Chirikof, and Marmot Bay. Estimates from these areas are given below. An exploratory survey along the shelf break from Sanak Island west to Unimak Island did not detect significant quantities of pollock.

2009 EIT survey results

		<i>Sanak</i>	<i>Shumagin</i>	<i>Shelikof</i>	<i>Chirikof</i>	<i>Marmot Bay</i>	<i>Total</i>
Total	Tons	31,435	63,337	265,971	396	19,759	380,898
	Percent	9%	18%	74%	0%	5%	
Biomass ≥43 cm	Tons	31,409	17,980	100,170	296	8789	158,644
	Percent	21%	12%	67%	0%	6%	

In comparison to 2008, biomass estimates were higher with the exception of Chirikof, where very few pollock were found (Fig. 1.7). In Sanak Gully, there was 22% increase, while in the Shumagin area there was 58% increase. For all areas surveyed, there was a 39% increase from the areas surveyed in 2008. However, biomass in all areas surveyed remains much lower than estimates from the early 2000s.

Since the assessment model only includes age 2 and older pollock, the biomass of age-1 fish in the 1995, 2000, 2005, and 2008 surveys was subtracted from the total biomass for those years, reducing the biomass by 15%, 13%, 5% and 9% respectively (Table 1.7). In all other years, the biomass of age-1 fish was less than 2% of the total EIT biomass estimate.

Echo Integrated Trawl Survey Length Frequency

Annual biomass distributions by length from the Shelikof Strait EIT survey show the progression of strong year classes through the population (Fig. 1.8). In the 2009 survey, the age-2 fish from the 2007 year class were both numerically dominant, and dominant in the biomass distribution by length. Since age composition estimates were already available from the 2008 survey, size composition data were not used in the assessment model.

Echo Integrated Trawl Survey Age Composition

Estimates of numbers at age from the Shelikof Strait EIT survey (Table 1.10) were obtained from random otolith samples and length frequency samples. Otoliths collected during the 1994 - 2009 EIT surveys were aged using the criteria described in Hollowed et al. (1995). Sample sizes for ages and lengths are given Table 1.11.

Egg Production Estimates of Spawning Biomass

Estimates of spawning biomass in Shelikof Strait based on egg production methods were included in the assessment model. A complete description of the estimation process is given in Picquelle and Megrey (1993). The estimates of spawning biomass in Shelikof Strait show a pattern similar to the acoustic survey (Table 1.7). The annual egg production spawning biomass estimate for 1981 is questionable because of sampling deficiencies during the egg surveys for that year (Kendall and Picquelle 1990). Coefficients of variation (CV) associated with these estimates were included in the assessment model. Egg production estimates were discontinued because the Shelikof Strait EIT survey provided similar information.

Alaska Department of Fish and Game Crab/Groundfish Trawl Survey

The Alaska Department of Fish and Game (ADF&G) has conducted bottom trawl surveys of nearshore areas of the Gulf of Alaska since 1987. Although these surveys are designed to monitor population trends of Tanner crab and red king crab, walleye pollock and other fish are also sampled. Standardized survey methods using a 400-mesh eastern trawl were employed from 1987 to the present. The survey is designed

to sample a fixed number of stations from mostly nearshore areas from Kodiak Island to Unimak Pass, and does not cover the entire shelf area. The average number of tows completed during the survey is 360. Details of the ADF&G trawl gear and sampling procedures are in Blackburn and Pengilly (1994).

The 2009 biomass estimate for pollock for the ADF&G crab/groundfish survey was 145,438 t, up 43% from the 2008 biomass estimate (Table 1.7).

ADF&G Survey Length Frequency

Pollock length-frequencies for the ADF&G survey in 1989-2009 (excluding 1991 and 1995) typically show a mode at lengths greater than 45 cm (Fig. 1.9). The predominance of large fish in the ADF&G survey may result from the selectivity of the gear, or because of greater abundance of large pollock in the areas surveyed. Length composition in 2009 is similar to previous surveys, with a mean length of approximately 50 cm.

ADF&G Survey Age Composition

Ages were determined by age readers in the AFSC age and growth unit from samples of pollock otoliths collected during the 2000, 2002, 2004, 2006, and 2008 ADF&G surveys (N = 559, 538, 591,588, and 597). Comparison with fishery age composition shows that older fish (> age-8) are more common in the ADF&G crab/groundfish survey. This is consistent with the assessment model, which estimates a domed-shaped selectivity pattern for the fishery, but an asymptotic selectivity pattern for the ADF&G survey.

Pre-1984 bottom trawl surveys

Considerable survey work was carried out in the Gulf of Alaska prior to the start of the NMFS triennial bottom trawl surveys in 1984. Between 1961 and the mid-1980s, the most common bottom trawl used for surveying was the 400-mesh eastern trawl. This trawl (or minor variants thereof) was used by IPHC for juvenile halibut surveys in the 1960s, 1970s, and early 1980s, and by NMFS for groundfish surveys in the 1970s.

Comparative work using the ADF&G 400-mesh eastern trawl and the NMFS poly-Nor' eastern trawl produced estimates of relative catchability (von Szalay and Brown 2001), making it possible to evaluate trends in pollock abundance from these earlier surveys in the pollock assessment. Von Szalay and Brown (2001) estimated a fishing power correction (FPC) for the ADF&G 400-mesh eastern trawl of 3.84 (SE = 1.26), indicating that 400-mesh eastern trawl CPUE for pollock would need to be multiplied by this factor to be comparable to the NMFS poly-Nor' eastern trawl.

In most cases, earlier surveys in the Gulf of Alaska were not designed to be comprehensive, with the general strategy being to cover the Gulf of Alaska west of Cape Spencer over a period of years, or to survey a large area to obtain an index for group of groundfish, i.e., flatfish or rockfish. For example, Ronholt et al. (1978) combined surveys for several years to obtain gulfwide estimates of pollock biomass for 1973-6. There are several difficulties with such an approach, including the possibility of double-counting or missing a portion of the stock that happened to migrate between surveyed areas.

An annual gulfwide index of pollock abundance was obtained using generalized linear models (GLM). Based on examination of historical survey trawl locations, four index sites were identified (one per INPFC area) that were surveyed relatively consistently during the period 1961-1983, and during the triennial survey time series (1984-99). The index sites were designed to include a range of bottom depths from nearshore to the continental slope. A generalized linear model (GLM) was fit to pollock CPUE data with year, site, depth strata (0-100 m, 100-200 m, 200-300 m, >300 m), and a site-depth interaction as factors. Both the pre-1984 400-mesh eastern trawl data and post-1984 triennial trawl survey data were

used. For the earlier period, analysis was limited to sites where at least 20 trawls were made during the summer (May 1-Sept 15).

Pollock CPUE data consist of observations with zero catch and positive values otherwise, so a GLM model with Poisson error and a logarithmic link was used (Hastie and Tibshirani 1990). This form of GLM has been used in other marine ecology applications to analyze trawl survey data (Smith 1990, Swartzman et al. 1992). The fitted model was used to predict mean CPUE by site and depth for each year with survey data. Predicted CPUEs (kg km^{-2}) were multiplied by the area within the depth strata (km^2) and summed to obtain proxy biomass estimates by INPFC area. Since each INPFC area contained only a single non-randomly selected index site, these proxy biomass estimates are potentially biased and would not incorporate the variability in relationship between the mean CPUE at an index site and the mean CPUE for the entire INPFC area. A comparison between these proxy biomass estimates by INPFC area and the actual NMFS triennial survey estimates by INPFC area for 1984-99 was used to obtain correction factors and variance estimates. Correction factors had the form of a ratio estimate (Cochran 1977), in which the sum of the NMFS survey biomass estimates for an INPFC area for 1984-99 is divided by the sum of the proxy biomass estimates for the same period.

Variances were obtained by bootstrapping data within site-depth strata and repeating the biomass estimation algorithm. A parametric bootstrap assuming a lognormal distribution was used for the INPFC area correction factors. Variance estimates do not reflect the uncertainty in the FPC estimate. In the assessment model, the FPC is not applied to the biomass estimates, but instead include the information about FPC estimate (mean and variance) was used as a likelihood component for relative survey catchability,

$$\log L = \frac{(q_1/q_2 - \hat{FPC})^2}{2 \sigma_{FPC}^2},$$

where q_1 is the catchability of the NMFS bottom trawl survey, q_2 is the catchability of historical 400-mesh eastern trawl surveys, \hat{FPC} is the estimated fishing power correction (= 3.84), and σ_{FPC} is the standard error of the FPC estimate (= 1.26).

Estimates of pollock biomass were very low (<300,000 t) between 1961 and 1971, increased by at least a factor of ten in 1974 and 1975, and then declined to approximately 900,000 t in 1978 (Table 1.12). No trend in pollock abundance is noticeable since 1978, and biomass estimates during 1978-1982 are in the same range as the post-1984 triennial survey biomass estimates. The coefficients of variation (CV) for GLM-based biomass estimates range between 0.24 and 0.64, and, as should be anticipated, are larger than the triennial survey biomass estimates, which range between 0.12 and 0.38.

Results were generally consistent with the multi-year combined survey estimates published previously (Table 1.12), and indicate a large increase in pollock biomass in the Gulf of Alaska occurred between the early 1960s (~200,000 t) and the mid 1970s (>2,000,000 t). Increases in pollock biomass between the 1960s and 1970s were also noted by Alton et al. (1987). In the 1961 survey, pollock were a relatively minor component of the groundfish community with a mean CPUE of 16 kg/hr (Ronholt et al. 1978). Arrowtooth flounder was the most common groundfish with a mean CPUE of 91 kg/hr. In the 1973-76 surveys, the CPUE of arrowtooth flounder was similar to the 1961 survey (83 kg/hr), but pollock CPUE had increased 20-fold to 321 kg/hr, and was by far the dominant groundfish species in the Gulf of Alaska. Meuter and Norcross (2002) also found that pollock was low in the relative abundance in 1960s, became the dominant species in Gulf of Alaska groundfish community in the 1970s, and subsequently declined in relative abundance.

Questions concerning the comparability of pollock CPUE data from historical trawl surveys with later surveys probably can never be fully resolved. However, because of the large magnitude of the change in CPUE between the surveys in the 1960s and the early 1970s using similar trawling gear, the conclusion that there was a large increase in pollock biomass seems robust. Model results suggest that population biomass in 1961, prior to large-scale commercial exploitation of the stock, may have been lower than at any time since then. Early speculation about the rise of pollock in the Gulf of Alaska in the early 1970s implicated the large biomass removals of Pacific ocean perch, a potential competitor for euphausiid prey (Somerton et al. 1979, Alton et al. 1987). More recent work has focused on role of climate change (Anderson and Piatt 1999, Bailey 2000). The occurrence of large fluctuations in pollock abundance without large changes in direct fishing impacts suggests a need for precautionary management. If pollock abundance is controlled primarily by the environment, or through indirect ecosystem effects, it may be difficult to reverse population declines, or to achieve rebuilding targets should the stock become depleted. Reliance on sustained pollock harvests in the Gulf of Alaska, whether by individual fishermen, processing companies, or fishing communities, may be difficult over the long-term.

Qualitative trends

To assess qualitatively recent trends in abundance, each survey time series was standardized by dividing the annual estimate by the average since 1987. Shelikof Strait EIT survey estimates prior to 2008 were rescaled to be comparable to subsequent surveys conducted by the *R/V Oscar Dyson*. Although there is considerable variability in each survey time series, a fairly clear downward trend is evident to 2000, followed by a stable, though variable, trend (Fig. 1.10). New surveys in 2009 show strongly contrasting trends, with both the NMFS and ADFG bottom trawl survey showing a steep increase to above the long-term average, while the Shelikof Strait EIT survey is close to 60% of the long-term average in 2009.

Indices derived from fisheries catch data were also evaluated for trends in biological characteristics (Fig. 1.11). The percent of females in the catch is close to 50-50, but shows a slight, though non-significant, downward trend, which may be related to changes in the seasonal distribution of the catch. The percent female increased to 54% in 2007 and remained at a similar level in 2008. The mean age shows interannual variability due to strong year classes passing through the population, but no downward trends that would suggest excessive mortality rates. The percent of old fish in the catch (nominally defined as age 8 and older) is also highly variable due to variability in year class strength. The percent of old fish increased to a peak in 1997, declined due to weaker recruitment in the 1990s and increases in total mortality (both from fishing and predation), but increased from 2005 to 2008 as the large 1999 and 2000 year classes entered the age-8 plus group. Under a constant $F_{40\%}$ harvest rate, the mean percent of age 8 and older fish in the catch is approximately 17%. An index of catch at age diversity was computed using the Shannon-Wiener information index,

$$-\sum p_a \ln p_a ,$$

where p_a is the proportion at age. Increases in fishing mortality would tend to reduce age diversity, but year class variability would also influence age diversity. The index of age diversity is relatively stable during 1976-2008 (Fig. 1.11).

McKelvey Index

McKelvey (1996) found a significant correlation between the abundance of age-1 pollock in the Shelikof Strait EIT survey and subsequent estimates of year-class strength. The McKelvey index is defined as the estimated abundance of 9-16 cm fish in the Shelikof Strait EIT survey, and is an index of recruitment at age 2 in the following year (Table 1.13). The relationship between the abundance of age-1 pollock in the

Shelikof Strait EIT survey and year-class strength provides a recruitment forecast for the year following the most recent Shelikof Strait EIT survey. The 2009 Shelikof EIT survey age-1 estimate is 0.33 billion (9th highest in abundance out of 26 surveys), which suggests that recruitment of the 2008 year class is likely to be close to the median level of abundance. However, the acoustic survey of the Shumagin area in 2009 produced an estimate of 2.17 billion age-1 pollock, which, if had occurred in the Shelikof Strait survey, would have been the third-largest estimate.

Analytic Approach

Model description

An age-structured model covering the period from 1961 to 2009 (49 yrs) was used to assess Gulf of Alaska pollock. Population dynamics were modeled using standard formulations for mortality and fishery catch (e.g. Fournier and Archibald 1982, Deriso et al. 1985, Hilborn and Walters 1992). Year- and age-specific fishing mortality was modeled as a product of a year effect, representing the full-recruitment fishing mortality, and an age effect, representing the selectivity of that age group to the fishery. The age effect was modeled using a double-logistic function with time-varying parameters (Dorn and Methot 1990, Sullivan et al. 1997). The model was fit to time series of catch biomass, survey indices of abundance, and estimates of age and length composition from the fishery and surveys. Details of the population dynamics and estimation equations are presented in an appendix.

Model parameters were estimated by maximizing the log likelihood of the data, viewed as a function of the parameters. Lognormal likelihoods were used for survey biomass and total catch estimates, and multinomial likelihoods were used for age and length composition data.

<i>Likelihood component</i>	<i>Statistical model for error</i>	<i>Variance assumption</i>
Fishery total catch (1964-2009)	Log-normal	CV = 0.05
POP fishery length comp. (1964-71)	Multinomial	Sample size = 60
Fishery age comp. (1972-2008)	Multinomial	Year-specific sample size = 60-400
Shelikof EIT survey biomass (1981-2009)	Log-normal	Survey-specific CV = 0.10-0.35
Shelikof EIT survey age comp. (1981-2009)	Multinomial	Sample size = 60
NMFS bottom trawl survey biom. (1984-2009)	Log-normal	Survey-specific CV = 0.12-0.38
NMFS bottom trawl survey age comp. (1984-2007)	Multinomial	Survey-specific sample size = 38-74
NMFS bottom trawl survey length comp. (2009)	Multinomial	Survey-specific sample size = 60
Egg production biomass (1981-92)	Log-normal	Survey specific CV = 0.10-0.25
ADF&G trawl survey biomass (1989-2009)	Log-normal	CV = 0.25
ADF&G survey age comp. (2000,2002,2004,2006, 2008)	Multinomial	Sample size = 10
ADF&G survey length comp. (1989-2009)	Multinomial	Sample size = 10
Historical trawl survey biomass (1961-1982)	Log-normal	Survey-specific CV = 0.24-0.64
Historical trawl survey age comp. (1973)	Multinomial	Sample size = 60
Historical trawl survey length comp. (1961-1982)	Multinomial	Sample size = 10
Fishery selectivity random walk process error	Log-normal	Slope CV = 0.10 (0.001 for 1961-71)
	Normal	Inflection age SD = 0.40 (0.004 for 1961-71)
Recruit process error (1961-1968,2009)	Log-normal	$\sigma_R = 1.0$

Recruitment

In most years, year-class abundance at age 2 was estimated as a free parameter. A prior constraint was imposed on recruitment at the start of the modeled time period to improve parameter estimability. Instead of estimating the abundance of each age of the initial age composition independently, we parameterized the initial age composition with mean log recruitment plus a log deviation from an equilibrium age structure based on that mean initial recruitment. A penalty was added to the log likelihood so that the log deviations would have the same variability as recruitment during the assessment period ($\sigma_R = 1.0$). We also used the same constraint for log deviations in recruitment for 1961-68, and in 2009. Log deviations were estimated as free parameters in other years. These relatively weak constraints were sufficient to obtain fully converged parameter estimates while retaining an appropriate level of uncertainty (e.g. the CV of recruitment in 2009 $\approx \sigma_R$).

Modeling fishery data

To accommodate changes in selectivity during the development of the fishery, we allowed the parameters of the double logistic function to vary according to a random walk process (Sullivan et al. 1997). This approach allows selectivity to vary from one year to the next, but restricts the amount of variation that can occur. The resulting selectivity patterns are similar to those obtained by grouping years, but transitions between selectivity patterns occur gradually rather than abruptly. Constraining the selectivity pattern for a group of years to be similar can be done simply by reducing the year-specific standard deviation of the process error term. Since limited data are available from the Pacific ocean perch fishery years (1964-71) and in 2009, the process error standard deviation for those years was assumed to be very small, so that annual changes in selectivity are very restricted during these years.

Modeling survey data

Survey abundance was assumed to be proportional to total abundance as modified by the estimated survey selectivity pattern. Expected population numbers at age for the survey were based on the mid-date of the survey, assuming constant fishing and natural mortality throughout the year. Standard deviations in the log-normal likelihood were set equal to the sampling error CV (coefficient of variation) associated with each survey estimate of abundance (Kimura 1991).

Survey catchability coefficients can be fixed or freely estimated. The NMFS bottom trawl survey catchability was fixed at one in this and previous assessments as a precautionary constraint on the total biomass estimated by the model. A likelihood profile on trawl catchability showed that the maximum likelihood estimate of trawl catchability was approximately 0.7. This result is reasonable because pollock are known to form pelagic aggregations and occur in nearshore areas not well sampled by the NMFS bottom trawl survey. Catchability coefficients for other surveys were estimated as free parameters. Egg production estimates of spawning stock biomass were included in the model by setting the age-specific selectivity equal to the estimated percent mature at age estimated by Hollowed et al. (1991).

The Simrad EK acoustic system has been used to estimate biomass since 1992. Earlier surveys (1981-91) were obtained with an older Biosonics acoustic system (Table 1.7). Biomass estimates similar to the Biosonics acoustic system can be obtained using the Simrad EK when a volume backscattering (S_v) threshold of -58.5 dB is used (Hollowed et al. 1992). Because of the newer system's lower noise level, abundance estimates since 1992 have been based on a S_v threshold of -70 dB. The Shelikof Strait EIT survey time series was split into two periods corresponding to the two acoustic systems, and separate survey catchability coefficients were estimated for each period. For the 1992 and 1993 surveys, biomass estimates using both noise thresholds were used to provide information on relative catchability.

A vessel comparison (VC) experiment was conducted in March 2007 during the Shelikof Strait acoustic-trawl survey. The VC experiment involved the *R/V Miller Freeman* (MF, the survey vessel used to

conduct Shelikof Strait surveys since the mid-1980s), and the *R/V Oscar Dyson* (OD), a noise-reduced survey vessel designed to conduct surveys that have traditionally been done with the *R/V Miller Freeman*. The vessel comparison experiment was designed to collect data either with the two vessels running beside one another at a distance of 0.7 nmi., or with one vessel following nearly directly behind the other at a distance of about 1 nmi. The methods were similar to those used during the 2006 Bering Sea VC experiment (De Robertis et al. 2008). Results indicate that the ratio of 38 kHz pollock backscatter from the *R/V Oscar Dyson* relative to the *R/V Miller Freeman* was significantly greater than one (1.13), as would be expected if the quieter OD reduced the avoidance response of the fish. Because this difference was significant, several methods were evaluated in the 2008 assessment for incorporating this result in the assessment model. The method that was adopted was to treat the MF and the OD time series as independent survey time series, and to include the vessel comparison results directly in the log likelihood of the assessment model. This likelihood component is given by

$$\log L = -\frac{1}{2(\sigma_S^2)} [\log(q_{OD}) - \log(q_{MF}) - \delta_{OD:MF}]^2,$$

where $\log(q_{OD})$ is the log catchability of the *R/V Oscar Dyson*, $\log(q_{MF})$ is the log catchability of the *R/V Oscar Dyson*, $\delta_{OD:MF} = 0.1240$ is the mean of log scale paired difference in backscatter, $\text{mean}[\log(s_{AOD}) - \log(s_{AMF})]$ obtained from the vessel comparison, and $\sigma_S = 0.0244$ is the standard error of the mean.

Ageing error

An ageing error conversion matrix is used in the assessment model to translate model population numbers at age to expected fishery and survey catch at age (Table 1.14). Dorn et al. (2003) estimated this matrix using an ageing error model fit to the observed percent reader agreement at ages 2 and 9. Mean percent agreement is close to 100% at age 1 and declines to 40% at age 10. Annual estimates of percent agreement are variable, but show no obvious trend, hence a single conversion matrix for all years in the assessment model was adopted. The model is based on a linear increase in the standard deviation of ageing error and the assumption that ageing error is normally distributed. The model predicts percent agreement by taking into account the probability that both readers are correct, both readers are off by one year in the same direction, and both readers are off by two years in the same direction (Methot 2000). The probability that both agree and were off by more than two years was considered negligible. A recent study evaluated pollock ageing criteria using radiometric methods and found them to be unbiased (Kastelle and Kimura 2006).

Length frequency data

The assessment model was fit to length frequency data from various sources by converting predicted age distributions (as modified by age-specific selectivity) to predicted length distributions using an age-length conversion matrix. Because seasonal differences in pollock length at age are large, several conversion matrices were used. For each matrix, unbiased length distributions at age were estimated for several years using age-length keys, then averaged across years. A conversion matrix estimated by Hollowed et al. (1998) was used for length-frequency data from the early period of the fishery. A conversion matrix was estimated using 1992-98 Shelikof Strait EIT survey data and used for winter survey length frequency data. The following length bins were used: 17 - 27, 28 - 35, 36 - 42, 43 - 50, 51 - 55, 56 - 70 (cm). Finally, a conversion matrix was estimated using second and third trimester fishery age and length data during the years (1989-98) and was used for the ADF&G survey length frequency data. The following length bins were used: 25 - 34, 35 - 41, 42 - 45, 46 - 50, 51 - 55, 56 - 70 (cm), so that the first three bins would capture most of the summer length distribution of the age-2, age-3 and age-4 fish, respectively. Bin definitions were different for the summer and the winter conversion matrices to account for the seasonal growth of the younger fish (ages 2-4).

Parameter estimation

A large number of parameters are estimated when using this modeling approach. More than half of these parameters are year-specific deviations in fishery selectivity coefficients. Parameters were estimated using ADModel Builder, a C++ software language extension and automatic differentiation library. Parameters in nonlinear models are estimated in ADModel Builder using automatic differentiation software extended from Greiwank and Corliss (1991) and developed into C++ class libraries. The optimizer in ADModel builder is a quasi-Newton routine (Press et al. 1992). The model is determined to have converged when the maximum parameter gradient is less than a small constant (set to 1×10^{-4}). ADModel builder includes post-convergence routines to calculate standard errors (or likelihood profiles) for any quantity of interest.

A list of model parameters is shown below:

<i>Population process modeled</i>	<i>Number of parameters</i>	<i>Estimation details</i>
Initial age structure	Ages 3-10 = 8	Estimated as log deviances from the log mean; constrained by random deviation process error from an equilibrium unfished age structure
Recruitment	Years 1961-2009 = 49	Estimated as log deviances from the log mean; recruitment in 1961-68, and 2009 constrained by random deviation process error.
Natural mortality	Age- and year-invariant = 1	Not estimated in the model
Fishing mortality	Years 1961-2009 = 49	Estimated as log deviances from the log mean
Mean fishery selectivity	4	Slope parameters estimated on a log scale, intercept parameters on an arithmetic scale
Annual changes in fishery selectivity	4 * (No. years -1) = 194	Estimated as deviations from mean selectivity and constrained by random walk process error
Survey catchability	No. of surveys + 1 = 7	AFSC bottom trawl survey catchability not estimated, other catchabilities estimated on a log scale. Two catchability periods were estimated for the EIT survey.
Survey selectivity	10 (EIT survey: 2, BT survey: 4, ADF&G survey: 2, Historical 400-mesh eastern trawls: 2)	Slope parameters estimated on a log scale. The egg production survey uses a fixed selectivity pattern equal to maturity at age.
Total	126 primary parameters + 192 process error parameters + 2 fixed parameters = 320	

Parameters Estimated Independently

Pollock life history characteristics, including natural mortality, growth, and maturity, were estimated independently. These parameters are used in the model to estimate spawning and population biomass and obtain predictions of fishery and survey biomass. Pollock life history parameters include:

- Natural mortality (M)
- Proportion mature at age

- Weight at age and year by fishery and by survey

Natural mortality

Hollowed and Megrey (1990) estimated natural mortality using a variety of methods including estimates based on: a) growth parameters (Alverson and Carney 1975, and Pauly 1980), b) GSI (Gunderson and Dygert, 1988), c) monitoring cohort abundance, and d) estimation in the assessment model. These methods produced estimates of natural mortality that ranged from 0.24 to 0.30. The maximum age observed was 22 years. For the assessment modeling, natural mortality was assumed to be 0.3 for all ages.

Hollowed et al. (2000) developed a model for Gulf of Alaska pollock that accounted for predation mortality. The model suggested that natural mortality declines from 0.8 at age 2 to 0.4 at age 5, and then remains relatively stable with increasing age. In addition, stock size was higher when predation mortality was included. In a theoretical study, Clark (1999) evaluated by the effect of an erroneous M on both estimated abundance and target harvest rates for a simple age-structured model. He found that “errors in estimated abundance and target harvest rate were always in the same direction, with the result that, in the short term, extremely high exploitation rates can be recommended (unintentionally) in cases where the natural mortality rate is overestimated and historical exploitation rates in the catch-at-age data are low.” He proposed that this error could be avoided by using a conservative (low) estimate of natural mortality. This suggests that the current approach of using a potentially low but still credible estimate of M for assessment modeling is consistent with the precautionary approach. However, it should be emphasized that the role of pollock as prey in the Gulf of Alaska ecosystem cannot be fully evaluated using a single species assessment model (Hollowed et al. 2000).

Maturity at age

In the 2002 assessment, maturity at age for Gulf of Alaska pollock was estimated using maturity stage data collected during winter EIT surveys in the Gulf of Alaska during 1983-2002. These estimates replaced a maturity at age vector estimated by Hollowed et al. (1991) who used maturity stage data collected during 1983-89. Maturity stages for female pollock describe a continuous process of ovarian development between immature and post-spawning. For the purposes of estimating a maturity vector (the proportion of an age group that has been or will be reproductively active during the year) for stock assessment, all fish greater than or equal to a particular maturity stage are assumed to be mature, while those less than that stage are assumed to be immature. Maturity stages in which ovarian development had progressed to the point where ova were distinctly visible were assumed to be mature. Maturity stages are qualitative rather than quantitative, so there is subjectivity in assigning stages, and a potential for different technicians to apply criteria differently. Because the link between pre-spawning maturity stages and eventual reproductive activity later in the season is not well established, the division between mature and immature stages is problematic. Changes in the timing of spawning could also affect maturity at age estimates. Merati (1993) compared visual maturity stages with ovary histology and a blood assay for vitellogenin and found general consistency between the different approaches. Merati (1993) noted that ovaries classified as late developing stage (i.e., immature) may contain yolked eggs, but it was unclear whether these fish would spawn later in the year. The average sample size of female pollock maturity stage data per year since 2000 from winter EIT surveys in the Gulf of Alaska is 360 (Table 1.15).

Estimates of maturity at age in 2009 from winter EIT surveys were below the long-term average for age 4 through age-6 pollock, but close to the long-term average for the older ages (Fig. 1.12). These inter-annual changes may reflect environmental conditions, pollock population biology, effect of strong year classes moving through the population, or simply ageing error. Because there did not appear to be an objective basis for excluding data, the 1983-2008 average maturity at age was used in the assessment.

Logistic regression (McCullagh and Nelder 1983) was also used to estimate the age and length at 50% maturity at age for each year. Annual estimates of age at 50% maturity are highly variable and range from 3.7 years in 1984 to 6.1 years in 1991, with an average of 4.9 years. Length at 50% mature is less variable than the age at 50% mature, suggesting that at least some of the variability in the age at maturity can be attributed to changes in length at age (Fig 1.13). Changes in year-class dominance could also potentially affect estimates of maturity at age. There is less evidence of trends in the length at 50% mature, with only the 1983 and 1984 estimates as unusually low values. The average length at 50% mature for all years is approximately 43 cm.

Weight at age

Year-specific weight-at-age estimates are used in the model to obtain expected catches in biomass. Where possible, year and survey-specific weight-at-age estimates are used to obtain expected survey biomass. For each data source, unbiased estimates of length at age were obtained using year-specific age-length keys. Bias-corrected parameters for the length-weight relationship, $W = a L^b$, were also estimated. Weights at age were estimated by multiplying length at age by the predicted weight based on the length-weight regressions.

Model evaluation

Model fit to age composition data was evaluated using plots of observed and predicted age composition in the fishery (Fig. 1.14), Shelikof Strait EIT survey (Fig. 1.15), and the NMFS trawl survey (Fig. 1.16). Model fits to fishery age composition data are good in most years. The fit of Shelikof Strait EIT survey age composition shows large residuals at age 2 and age 3 in 2006-2009 due to inconsistencies between the initial estimates of abundance and subsequent information about the magnitude of these year classes.

General trends in survey time series fit reasonably well and model fits to survey biomass estimates are similar to previous assessments (Dorn et al. 2008) (Figs. 1.17-1.19). The discrepancy between the NMFS trawl survey and the Shelikof Strait EIT survey biomass estimates in the 1980s accounts for the poor model fit to both time series during those years. It was not possible for the model to fit all the 2009 survey estimates simultaneously. Both the NMFS bottom trawl survey and the ADF&G surveys showed large increases in biomass in 2009 (123% increase for the NMFS bottom trawl survey, 43% increase for the ADF&G crab/groundfish survey), while the Shelikof Strait EIT showed only a slight increase (15%) and remains close to historically low levels. For a pollock population to increase by the amount indicated by the NMFS bottom trawl survey, recruitment to population would have to have been very large. Available information (including the length information from the NMFS and the ADF&G surveys) does not indicate that extremely large year classes have recruited to the pollock population. Model results show a modest increase in survey biomass in 2009, which results in a strongly positive residual for the NMFS and ADF&G bottom trawl surveys in 2009, and a strongly negative residual for the Shelikof Strait EIT survey.

A likelihood profile for NMFS trawl survey catchability shows that the likelihood is higher for models with catchability equal to 0.70 (Fig. 1.20), compared to the estimate of 0.74 in the 2008 assessment. The change in log likelihood is small (about 2.0) between models with fixed and estimated catchability, indicating that despite the large change in biomass, there is little objective basis for choosing one model over the other. These results are similar to previous assessments. Consequently we used a base model with fixed trawl survey catchability of 1.0 to be consistent with recommendations in previous assessments.

Assessment Model Results

Parameter estimates and model output are presented in a series of tables and figures. Estimated survey selectivity and fishery selectivity for different periods given in Table 1.16 (see also Figure 1.21). Table 1.17 gives the estimated population numbers at age for the years 1961-2009. Table 1.18 gives the estimated time series of age 3+ population biomass, age-2 recruitment, and harvest rate (catch/3+ biomass) for 1977-2008 (see also Fig. 1.22). Stock size peaked in the early 1980s at approximately 1.1 times the proxy for unfished stock size ($B_{100\%}$ = mean 1979-2008 recruitment multiplied by the spawning biomass per recruit in the absence of fishing ($SPR@F=0$)). In 1998, the stock dropped below the $B_{40\%}$ for the first time since the 1970s, reached a minimum in 2003 of 21% of unfished stock size, increased to 33% of unfished in 2006. The stock then declined slightly, dropping to 29% of unfished stock size in 2009.

Retrospective comparison of assessment results

A retrospective comparison of assessment results for the years 1996-2009 indicates the current estimated trend in spawning biomass for 1990-2009 is consistent with previous estimates (Fig. 1.23, top panel). All time series show a similar pattern of decreasing spawning biomass in the 1990s followed by a period of greater stability in 2000s. There appear to be no consistent pattern of bias in estimates of ending year biomass, but assessment errors are clearly correlated over time, such that there are runs of over estimates and under estimates. The estimated 2009 age composition from the current assessment is similar to projected 2009 age composition in the 2008 assessment (Fig. 1.23, bottom panel). The largest discrepancy is the estimate of the age-3 fish (2006 year class), which is about half the size of last year's assessment. In addition, the estimate of the age-2 fish (2007 year class) is 1.2 billion fish, compared to a mean recruitment of 0.7 billion in the 2008 assessment.

Stock and recruitment

Recruitment of Gulf of Alaska pollock is more variable ($CV = 1.06$) than Eastern Bering Sea pollock ($CV = 0.62$). Other North Pacific groundfish stocks, such as sablefish and Pacific ocean perch, also have high recruitment variability. However, unlike sablefish and Pacific ocean perch, pollock have a short generation time (<10 yrs), so that large year classes do not persist in the population long enough to have a buffering effect on population variability. Because of these intrinsic population characteristics, the typical pattern of biomass variability for Gulf of Alaska pollock will be sharp increases due to strong recruitment, followed by periods of gradual decline until the next strong year class recruits to the population. Gulf of Alaska pollock is more likely to show this pattern than any other groundfish stock in the North Pacific due to the combination of a short generation time and high recruitment variability.

Since 1980, strong year classes have occurred every four to six years (Fig. 1.22). Because of high recruitment variability, the functional relationship between spawning biomass and recruitment is difficult to estimate despite good contrast in spawning biomass. Strong and weak year classes have been produced at high and low level of spawning biomass. The 1972 year class (one of the largest on record) was produced by an estimated spawning biomass close to current levels, suggesting that the stock has the potential to produce strong year classes. Spawner productivity is higher at low spawning biomass compared to high spawning biomass, indicating that survival of eggs to recruitment is density-dependent (Fig. 1.24). However, this pattern of density-dependent survival only emerges on a decadal scale, and could be confounded with environmental variability on the same temporal scale. These decadal trends in spawner productivity have produced the pattern of increase and decline in the GOA pollock population. The last two decades have been a period of relatively low spawner productivity.

We summarize information on recent year classes in the table below. The high estimates of age-1 fish in both the Shumagin and Shelikof Strait EIT surveys in 2008 suggests that the 2007 year class is both

abundant and widely distributed. The model estimate of the 2007 year class in 1.2 billion, approximately twice as large as an average year class. For the 2008 year class, estimates of age-1 abundance were very high in the Shumagin EIT survey, but not in the Shelikof Strait EIT survey.

Year of recruitment	2009	2010	2011
Year class	2007	2008	2009
FOCI prediction	<i>Average</i>	<i>Average</i>	<i>Average</i>
Survey information	2008 Shelikof EIT survey age-1 estimate is 1.4 billion (5th in abundance out of 25 surveys) 2008 Shumagin EIT survey age-1 estimate is 1.5 billion	2009 Shelikof EIT survey age-1 estimate is 0.33 billion (9th in abundance out of 26 surveys) 2009 Shumagin EIT survey age-1 estimate is 2.2 billion	

Projections and Harvest Alternatives

Reference fishing mortality rates and spawning biomass levels

Since 1997, Gulf pollock have been managed under Tier 3 of NPFMC harvest guidelines. In Tier 3, reference mortality rates are based on the spawning biomass per recruit (SPR), while biomass reference levels are estimated by multiplying the SPR by average recruitment. Estimates of the F_{SPR} harvest rates were obtained using the life history characteristics of Gulf of Alaska pollock (Table 1.19). Spawning biomass reference levels were based on mean 1979-2008 recruitment (689 million), which slightly lower than the post-1979 mean in the 2008 assessment. The average did not include the recruitment in 2009 (2007 year class) due to uncertainty in the estimate of year class strength. Spawning was assumed to occur on March 15th, and female spawning biomass was calculated using mean weight at age for the Shelikof Strait EIT surveys in 2005-2009 to estimate current reproductive potential. The SPR at $F=0$ was estimated as 0.899 kg/recruit. This estimate represents an 8% increase from the 2008 estimate primarily due to increases in weight at age in the 2009 Shelikof Strait EIT survey. F_{SPR} rates depend the selectivity pattern of the fishery. Selectivity in the Gulf of Alaska pollock fishery changed as the fishery evolved from a foreign fishery occurring along the shelf break to a domestic fishery on spawning aggregations and in nearshore waters (Fig. 1.1). For SPR calculations, we used a selectivity pattern based on an average for 2004-2008 to reflect current selectivity patterns. Gulf of Alaska pollock F_{SPR} harvest rates are given below:

F_{SPR} rate	Fishing mortality	<i>Equilibrium under average 1979-2008 recruitment</i>				
		<i>Avg. Recr. (Million)</i>	<i>Total 3+ biom. (1000 t)</i>	<i>Female spawning biom. (1000 t)</i>	<i>Catch (1000 t)</i>	<i>Harvest rate</i>
100.0%	0.000	689	1721	620	0	0.0%
50.0%	0.169	689	1085	310	139	12.8%
45.0%	0.197	689	1018	279	152	14.9%

40.0%	0.229	689	949	248	165	17.4%
35.0%	0.267	689	878	217	178	20.2%

The $B_{40\%}$ estimate of 248,000 t represents a 4% increase from the $B_{40\%}$ estimate of 237,000 t in the 2008 assessment, and reflects both the increase in mean weight at age during spawning and a decrease in average recruitment. The model estimate of spawning biomass in 2010 is 184,567 t, which is 29.8% of unfished spawning biomass and below $B_{40\%}$ (248,000 t), thereby placing Gulf of Alaska pollock in sub-tier “b” of Tier 3. In sub-tier “b” the OFL and maximum permissible ABC fishing mortality rates are adjusted downwards as described by the harvest guidelines (see SAFE Summary Chapter).

2010 acceptable biological catch

The definitions of OFL and maximum permissible F_{ABC} under Amendment 56 provide a buffer between the overfishing level and the intended harvest rate, as required by NMFS national standard guidelines. Since estimates of stock biomass from assessment models are uncertain, the buffer between OFL and ABC provides a margin of safety so that assessment error will not result in the OFL being inadvertently exceeded. For Gulf of Alaska pollock, the maximum permissible F_{ABC} harvest rate is 86.0% of the OFL harvest rate. In the 2001 assessment, based on an analysis that showed that the buffer between the maximum permissible F_{ABC} and OFL decreased when the stock is below approximately $B_{50\%}$, we developed a more conservative alternative that maintains a constant buffer between ABC and F_{ABC} at all stock levels (Table 1.20). While there is always some probability of exceeding F_{OFL} due to imprecise stock assessments, it seemed unreasonable to reduce safety margin as the stock declines.

This alternative is given by the following

$$\text{Define } B^* = B_{40\%} \frac{F_{35\%}}{F_{40\%}}$$

$$\text{Stock status: } B / B^* > 1, \text{ then } F = F_{40\%}$$

$$\text{Stock status: } 0.05 < B / B^* \leq 1, \text{ then } F = F_{40\%} \times (B / B^* - 0.05) / (1 - 0.05)$$

$$\text{Stock status: } B / B^* \leq 0.05, \text{ then } F = 0$$

This alternative has the same functional form as the maximum permissible F_{ABC} ; the only difference is that it declines linearly from B^* ($= B_{47\%}$) to $0.05B^*$ (Fig. 1.25).

Projections for 2010 for F_{OFL} , the maximum permissible F_{ABC} , and an adjusted $F_{40\%}$ harvest rate with a constant buffer between F_{ABC} and F_{OFL} are given in Table 1.21.

ABC recommendation

There were three new surveys in 2009: the NMFS bottom trawl survey, the ADF&G crab/groundfish survey, and the Shelikof Strait EIT survey. While all surveys showed an increase in pollock biomass in 2009, results from these surveys were not consistent with each other. Biomass was steeply higher in the NMFS and the ADF&G surveys, while the Shelikof Strait EIT showed a relatively modest increase.

Length data collected during the surveys does not suggest extremely strong recruitment to population, as would be necessary for the population to double over two years. Winter EIT surveys in other areas of the Gulf of Alaska also showed increases, but remain close to historically low levels.

When this information is incorporated into the assessment, the model is unable to achieve a satisfactory fit to any survey in 2009, but produces estimates of stock size that are between those that would be obtained if each of the three surveys were used individually. The estimated abundance of mature fish in 2010 stock size is 17% higher than projected in the 2008 assessment, and is projected to increase further over the next five years.

The initial estimate 2007 year class is 1.7 times average recruitment, and was abundant in both Shumagin area and Shelikof Strait in the 2008 EIT surveys. However initial estimates of year-class strength are highly uncertain, and there have been several instances recently when an initial estimate of year class size decreased as more information accumulated (for example the 1994 and 1999 year classes). For the 1999 year class, a decision was made set the year class equal to average recruitment until a stable estimate of year class abundance was obtained. In retrospect, this decision appears to have been appropriate, though of course there is no guarantee that this will always be the case. The *ad hoc* nature of this approach is problematic, and refinements may be possible through Management Strategy Evaluations (MSE). Here we provide both a projection for the estimated 2007 year class, and a projection where the 2007 year class has been set to the average.

Although the surveys indicate an upwards trend in stock abundance, the lack of consistency between surveys is a major concern. In addition, the continued low spawning biomass in Shelikof Strait and other spawning areas in the Gulf of Alaska in 2009 is a concern. In previous years, concern over the decline in spawning activity in Shelikof Strait was mitigated by the additional winter surveying efforts which in aggregate resulted in an estimate of spawning biomass that was close to the model estimate. In 2009, the aggregate spawning biomass was 54% of the model estimate, so this was not the case in 2009.

Based on these considerations, we considered that a more cautious approach is warranted for 2010, and used the model with an average 2007 year class and an adjusted $F_{40\%}$ harvest rate for the author's recommended 2010 ABC of 77,150 t. The elements of risk-aversion in this recommendation relative to using the point estimate of the model and the maximum permissible F_{ABC} are the following: 1) fixing trawl catchability at 1.0; 2) applying a more conservative harvest rate than the maximum permissible F_{ABC} , 3) setting the 2007 year class equal to the average. These risk-averse elements reduce the recommended ABC to approximately 50% of the model point estimate. In 2011, the ABC based an adjusted $F_{40\%}$ harvest rate is 101,510 t (Table 1.22). The OFL in 2010 is 103,210 t, and the OFL in 2011 if the recommended ABC is taken in 2010 is 135,010 t.

To evaluate the probability that the stock will drop below the $B_{20\%}$ threshold, we projected the stock forward for five years and removed catches based on the spawning biomass in each year and the author's recommended fishing mortality schedule. This projection incorporates uncertainty in stock status, uncertainty in the estimate of $B_{20\%}$, and variability in future recruitment. We then sampled from the likelihood of future spawning biomass using Markov chain Monte Carlo (MCMC) (Fig. 1.26). A chain of 1,000,000 samples was thinned by selecting every 200th sample. Analysis of the thinned MCMC chain indicates that probability of the stock dropping below $B_{20\%}$ will be negligible in all years.

Projections and Status Determination

A standard set of projections is required for stocks managed under Tier 3 of Amendment 56. This set of projections encompasses seven harvest scenarios designed to satisfy the requirements of Amendment 56, the National Environmental Protection Act, and the Magnuson-Stevens Fishery Conservation and

Management Act (MSFCMA). For each scenario, the projections begin with the 2009 numbers at age as estimated by the assessment model and assume the 2009 catch will be 43,865 t (Tom Pearson pers. comm., October 16, 2009). In each year, the fishing mortality rate is determined by the spawning biomass in that year and the respective harvest scenario. Recruitment is drawn from an inverse Gaussian distribution whose parameters consist of maximum likelihood estimates determined from recruitments during 1979-2008 as estimated by the assessment model. Spawning biomass is computed in each year based on the time of peak spawning (March 15) using the maturity and weight schedules in Table 1.19. This projection scheme is run 1000 times to obtain distributions of possible future stock sizes, fishing mortality rates, and catches.

Five of the seven standard scenarios are used in an Environmental Assessment prepared in conjunction with the final SAFE. These five scenarios, which are designed to provide a range of harvest alternatives that are likely to bracket the final TAC for 2010, are as follows (“ $max F_{ABC}$ ” refers to the maximum permissible value of F_{ABC} under Amendment 56):

Scenario 1: In all future years, F is set equal to $max F_{ABC}$. (Rationale: Historically, TAC has been constrained by ABC, so this scenario provides a likely upper limit on future TACs.)

Scenario 2: In all future years, F is set equal to the F_{ABC} recommended in the assessment.

Scenario 3: In all future years, F is set equal to the five-year average F (2004-2008). (Rationale: For some stocks, TAC can be well below ABC, and recent average F may provide a better indicator of F_{TAC} than F_{ABC} .)

Scenario 4: In all future years, F is set equal to $F_{75\%}$. (Rationale: This scenario represents a very conservative harvest rate and was requested by the Regional Office based on public comment.)

Scenario 5: In all future years, F is set equal to zero. (Rationale: In extreme cases, TAC may be set at a level close to zero.)

Two other scenarios are needed to satisfy the MSFCMA’s requirement to determine whether a stock is currently in an overfished condition or is approaching an overfished condition. These two scenarios are as follow (for Tier 3 stocks, the MSY level is defined as $B_{35\%}$):

Scenario 6: In all future years, F is set equal to F_{OFL} . (Rationale: This scenario determines whether a stock is overfished. If the stock is expected to be 1) above its MSY level in 2009 or 2) above 1/2 of its MSY level in 2009 and above its MSY level in 2019 under this scenario, then the stock is not overfished)

Scenario 7: In 2010 and 2011, F is set equal to $max F_{ABC}$, and in all subsequent years, F is set equal to F_{OFL} . (Rationale: This scenario determines whether a stock is approaching an overfished condition. If the stock is expected to be 1) above its MSY level in 2012, or 2) above 1/2 of its MSY level in 2012 and above its MSY level in 2022 under this scenario, then the stock is not approaching an overfished condition.)

Results from scenarios 1-5 are presented in Table 1.21. A separate table (Table 1.22) is given in which the 2007 year class (age-2 fish in 2009) is assumed equal to the 1979-2008 average recruitment at age 2. Under all harvest policies, mean spawning biomass is projected to increase gradually over the next five years (Fig. 1.27). Plots of individual projection runs are highly variable (Fig. 1.28), and may provide a more realistic view of potential pollock abundance in the future.

Under the MSFCMA, the Secretary of Commerce is required to report on the status of each U.S. fishery with respect to overfishing. This report involves the answers to three questions: 1) Is the stock being subjected to overfishing? 2) Is the stock currently overfished? 3) Is the stock approaching an overfished condition?

The catch estimate for the most recent complete year (2008) is 52,500 t, which is less than the 2008 OFL of 72,110 t. Therefore, the stock is not being subject to overfishing.

Scenarios 6 and 7 are used to make the MSFCMA's other required status determination as follows:

Spawning biomass is estimated to be 163,748 t in 2009, which is less than $B_{35\%}$ (217,000 t), but greater than $\frac{1}{2}$ of $B_{35\%}$. Under scenario 6, the projected mean spawning biomass in 2019 is 242,595 t, 112% of $B_{35\%}$. Therefore, Gulf of Alaska pollock are not currently overfished.

Under scenario 7, projected mean spawning biomass in 2012 is 241,264 t, which is 111% of $B_{35\%}$. Therefore, Gulf of Alaska pollock is not approaching an overfished condition.

Ecosystem considerations

Prey of pollock

An ECOPATH model was assembled to characterize food web structure in Gulf of Alaska using diet data and population estimates during 1990-93. We use ECOPATH here simply as a tool to integrate diet data and stock abundance estimates in a consistent way to evaluate ecosystem interactions. We focus primarily on first-order trophic interactions: prey of pollock and the predators of pollock.

Pollock trophic interactions occur primarily in the pelagic pathway in the food web, which leads from phytoplankton through various categories of zooplankton to planktivorous fish species such as capelin and sandlance (Fig. 1.29); the primary prey of pollock are euphausiids. Pollock also consume shrimp, which are more associated with the benthic pathway, and make up approximately 18% of age 2+ pollock diet. All ages of GOA pollock are primarily zooplanktivorous during the summer growing season (>80% by weight zooplankton in diets for juveniles and adults; Fig 1.30). While there is an ontogenetic shift in diet from copepods to larger zooplankton (primarily euphausiids) and fish (Fig. 1.30), cannibalism is not as prevalent in the Gulf of Alaska as in the Eastern Bering Sea, and fish consumption is low even for large pollock (Yang and Nelson 2000).

There are no extended time series of zooplankton abundance for the shelf waters of the Gulf of the Alaska. Brodeur and Ware (1995) provide evidence that biomass of zooplankton in the center of the Alaska Gyre was twice as high in the 1980s than in the 1950s and 1960s, consistent with a shift to positive values of the PDO since 1977. The percentage of zooplankton in diets of pollock is relatively constant throughout the 1990s (Fig. 1.30). While indices of stomach fullness exist for these survey years, a more detailed bioenergetics modeling approach would be required to examine if feeding and growth conditions have changed over time, especially given the fluctuations in GOA water temperature in recent years (Fig. 15, Ecosystem Considerations Appendix), as water temperature has a considerable effect on digestion and other energetic rates.

Predators of pollock

Initial ECOPATH model results show that the top five predators on pollock >20 cm by relative importance are arrowtooth flounder, Pacific halibut, Pacific cod, Steller sea lion (SSL), and the directed pollock fishery (Fig. 1.31). For pollock less than 20cm, arrowtooth flounder represent close to 50% of

total mortality. All major predators show some diet specialization, and none depend on pollock for more than 50% of their total consumption (Fig. 1.32). Pacific halibut is most dependent on pollock (48%), followed by SSL (39%), then arrowtooth flounder (24% for juvenile and adult pollock combined), and lastly Pacific cod (18%). It is important to note that although arrowtooth flounder is the largest single source of mortality for both juvenile and adult pollock (Fig 1.31), arrowtooth depend less on pollock in their diets than do the other predators.

Arrowtooth consume a greater number of smaller pollock than do Pacific cod or Pacific halibut, which consume primarily adult fish. However, by weight, larger pollock are important to all three predators (Fig. 1.33). Length frequencies of pollock consumed by the western stock of Steller sea lions tend towards larger fish, and generally match the size frequencies of cod and halibut (Zeppelin et al. 2004). The diet of Pacific cod and Pacific halibut are similar in that the majority of their diet besides pollock is from the benthic pathway of the food web. Alternate prey for Steller sea lions and arrowtooth flounder are similar, and come primarily from the pelagic pathway.

Predation mortality, as estimated by ECOPATH, is extremely high for GOA pollock >20cm. Estimates for the 1990-1993 time period indicate that known sources of predation sum to 90%-120% of the total production of walleye pollock calculated from 2004 stock assessment growth and mortality rates; estimates greater than 100% may indicate a declining stock (as shown by the stock assessment trend in the early 1990s; Fig 1.34, top), or the use of mortality rates which are too low. Conversely, as >20cm pollock include a substantial number of 2-year olds, it may be that mortality rate estimates for this age range is low. In either case, predation mortality for pollock in the GOA is much greater a proportion of pollock production than as estimated by the same methods for the Bering Sea, where predation mortality (primarily pollock cannibalism) was up to 50% of total production.

Aside from long-recognized decline in Steller sea lion abundance, the major predators of pollock in the Gulf of Alaska are stable to increasing, in some cases notably so since the 1980s (Fig. 1.34, top). This high level of predation is of concern in light of the declining trend of pollock with respect to predator increases. To assess this concern, it is important to determine if natural mortality may have changed over time (e.g. the shifting control hypothesis; Bailey 2000). To examine predator interactions more closely than in the initial model, diet data of major predators in trawl surveys were examined in all survey years since 1990.

Trends in total consumption of walleye pollock were calculated by the following formula:

$$Consumption = \sum B_{pred, size, subregion} \cdot DC_{pred, size, subregion} \cdot WLF_{pred, size, GOA} \cdot Ration_{pred, size}$$

where B(pred, size, subregion) is the biomass of a predator size class in the summer groundfish surveys in a particular survey subregion; DC is the percentage by weight of pollock in that predator group as measured from stomach samples, WLF is the weight frequency of pollock in the stomachs of that predator group pooled across the GOA region, calculated from length frequencies in stomachs and length-weight relationships from the surveys. Finally, ration is an applied yearly ration for that predator group calculated by fitting weight-at-age to the generalized von Bertalanffy growth equations as described in Essington et al. (2001). Ration is assumed fixed over time for a given size class of predator.

Fig. 1.34 (bottom) shows annual total estimates of consumption of pollock (all age classes) in survey years by the four major fish predators. Other predators, shown as constant, are taken from ECOPATH modeling results and displayed for comparison. Catch is shown as reported in Table 1.1. In contrast, the line in the figure shows the historical total production (tons/year) plus yearly change in biomass (positive or negative) from the stock assessment results. In a complete accounting of pollock mortality, the height

of the bars should match the height of the line. As shown, estimates of consumption greatly surpass estimates of production; fishing mortality is a relatively small proportion of total consumption. Overestimates in consumption rates could arise through seasonal differences in diets; while ration is seasonally adjusted, diet proportions are based on summer data. Also, better energetic estimates of consumption would improve these estimates. In terms of the stock assessment, underestimates of production could result from underestimating natural mortality, especially at ages 2-3, underestimating the rate of decline which occurred between 1990-present, or underestimates of the total biomass of pollock; this analysis should be revisited using higher mortality at younger ages than assumed in the current stock assessment.

To better judge natural mortality, consumption was calculated for two size groups of pollock, divided at 30cm fork length. This size break, which differs from the break in the ECOPATH analysis, is based on finding minima between modes of pollock in predator diets (Fig. 1.33). This break is different from the conversion matrices used in the stock assessment; perhaps due to differences in size selection between predators and surveys. For this analysis, it is assumed that pollock <30cm are ages 0-2 while pollock \geq 30cm are age 3+ fish.

Consumption of age 0-2 pollock per unit predator biomass (using survey biomass) varied considerably through survey years, although within a year all predators had similar consumption levels (Fig. 1.35, top). Correlation coefficients of consumption rates were 0.98 between arrowtooth and halibut, and 0.90 for both of these species with pollock. Correlation coefficients of these three species with cod were \sim 0.55 for arrowtooth and halibut and \sim 0.20 with pollock. The majority of this predation by weight occurred on age 2 pollock.

Plotted against age 2 pollock numbers calculated from the stock assessment, consumption/biomass and total consumption by predator shows a distinct pattern (Fig. 1.35, lower two graphs). In “low” recruitment years consumption is consistently low, while in high recruitment years consumption is high, but does not increase linearly, rather consumption seems to level out at high numbers of juvenile pollock, resembling a classic “Type II” functional response. This suggests the existence bottom-up control of juvenile consumption, in which strong year classes of pollock “overwhelm” feeding rates of predators, resulting in potentially lower juvenile mortality in good recruitment years which may amplify the recruitment. However, this result should be examined iteratively within the stock assessment, as the back-calculated numbers at age 2 assume a constant natural mortality rate. Assuming a lower mortality rate due to predator satiation would lead to lower estimates of age 2 numbers, which would make the response appear more linear.

Consumption of pollock \geq 30cm shows a different pattern over time. A decline of consumption per unit biomass is evident for halibut and cod (Fig. 1.36, top). Arrowtooth shows an insignificant decline; it is possible that the noise in the arrowtooth trend, mirroring the consumption of <30cm fish, is due to the choice of 30cm as an age cutoff. As a function of age 3+ assessment biomass, consumption per unit biomass and total consumption remained constant as the stock declined, and then fell off rapidly at low biomass levels in recent years (Fig. 1.36, middle and bottom). Again, this result should be approached iteratively, but it suggests increasing predation mortality on age 3+ pollock between 1990-2005, possibly requiring increased foraging effort from predators.

There has been a marked decline in Pacific halibut weight at age since the 1970s that Clark et al. (1999) attributed to the 1977 regime shift without being able to determine the specific biological mechanisms that produced the change. Possibilities suggested by Clark et al. (1999) include the physiological effect of an increase in temperature, intra- and interspecific competition for prey, or a change in prey quality. The two species most dependent on pollock in the early 1990s (Pacific halibut and Steller sea lion) have both shown an exceptional biological response during the post-1977 period consistent with a reduction in

carrying capacity (growth for Pacific halibut, survival for Steller sea lions). In contrast, the dominant predator on pollock in the Gulf of Alaska (arrowtooth flounder) has increased steadily in abundance over the same period and shows no evidence of decline in size at age. Given that arrowtooth flounder has a range of potential prey types to select from during periods of low pollock abundance (Fig. 1.32), we do not expect that arrowtooth would decline simply due to declines in pollock.

Taken together, Figs. 1.35 and 1.36 suggest that recruitment remains bottom-up controlled even under the current estimates of high predation mortality, and may lead to strong year classes. However, top-down control seems to have increased on age 3+ pollock in recent years, perhaps as predators have attempted to maintain constant pollock consumption during a period of declining abundance. It is possible that natural mortality on adult pollock will remain high in the ecosystem in spite of decreasing pollock abundance.

Ecosystem modeling

To examine the relative role of pollock natural versus fishing mortality within the GOA ecosystem, a set of simulations were run using the ECOPATH model shown in Fig. 1.29. Following the method outlined in Aydin et al. (2005), 20,000 model ecosystems were drawn from distributions of input parameters; these parameter sets were subjected to a selection/rejection criteria of species persistence resulting in approximately 500 ecosystems with nondegenerate parameters. These models, which did not begin in an equilibrium state, were projected forward using ECOSIM algorithms until equilibrium conditions were reached. For each group within the model, a perturbation experiment was run in all acceptable ecosystems by reducing the species survival (increasing mortality) by 10%, or by reducing gear effort by 10%, and reporting the percent change in equilibrium of all other species or fisheries catches. The resulting changes are reported as ranges across the generated ecosystems, with 50% and 95% confidence intervals representing the distribution of percent change in equilibrium states for each perturbation.

Fig. 1.37 shows the changes in other species when simulating a 10% decline in adult pollock survival (top graph), a 10% decline in juvenile pollock survival (middle graph), and a 10% decline in pollock trawl effort. Fisheries in these simulations are governed by constant fishing mortality rates rather than harvest control rules. Only the top 20 effects are shown in each graph; note the difference in scales between each graph.

The model results indicate that the largest effects of declining adult pollock survival would be declines in halibut and Steller sea lion biomass. Declines in juvenile survival would have a range of effects, including halibut and Steller sea lions, but also releasing a range of competitors for zooplankton including rockfish and shrimp. The pollock trawl itself has a lesser effect throughout the ecosystem (recall that fishing mortality is small in proportion to predation mortality for pollock); the strongest modeled effects are not on competitors for prey but on incidentally caught species (Table 1.2), with the strongest effects being on sharks.

The results presented above are taken from Gulfwide weighted averages of consumption; Steller sea lions and the fishing fleet are central place foragers, making foraging trips from specific locations (ports in the case of the fishing fleet, and rookeries or haulouts for Steller sea lions). Foraging bouts (or trawl sets) begin at the surface, and foragers attack their prey from the top down. For such species, directed and local changes in fishing may have a disproportionate effect compared to the results shown here.

In contrast, predation by groundfish is not as constrained geographically, and captures are likely to occur when the predator swims upwards from the bottom. Changes in the vertical distribution of pollock may tend to favor one mode of foraging over another. For example, if pollock move deeper in the water column due to surface warming, foraging groundfish might obtain an advantage over surface foragers.

Alternatively, pollock may respond adaptively to predation risks from groundfish or surface foragers by changing its position in the water column.

Of species affecting pollock (Fig. 1.38), arrowtooth have the largest impact on adult pollock, while bottom-up processes (phytoplankton and zooplankton) have the largest impact on juvenile pollock. It is interesting to note that the link between juvenile and adult pollock is extremely uncertain (wide error bars) within these models.

Finally, of the four major predators of pollock (Fig 1.39), all are affected by bottom-up forcing; Steller sea lions, Pacific cod, and Pacific halibut are all affected by pollock perturbations, while pollock effects on arrowtooth are much more minor.

Pair-wise correlations in predator trends were examined for consistent patterns (Fig. 1.40). For each pair-wise comparison, we used the maximum number of years available. Time series for Steller sea lions and Pacific cod begin in mid 1970s, while other time series extend back to the early 1960s. We make no attempt to evaluate statistical significance (biomass trends are highly autocorrelated), and emphasize that correlation does not imply causation. If two populations are strongly correlated in time, there are many possible explanations: both populations are responding to similar forcing, one or other is causative agent, etc.

Pollock abundance, fishery catches, and Steller sea lions are positively correlated (Fig. 1.40). Since the harvest policy for pollock is modified fixed harvest rate strategy, a positive correlation between catch and abundance would be expected. The Steller sea lion trend is more strongly correlated with pollock abundance than pollock catches, but this correlation is based on data since 1976, and does not include earlier years of low pollock abundance. The only strong inverse correlation is between arrowtooth flounder and Steller sea lions. A strong positive correlation exists between Pacific cod and Pacific halibut, and, from the 1960s to the present, between Pacific halibut and arrowtooth flounder.

Several patterns are apparent in abundance trends and the diet data. First, the two predators with alternate prey in the benthic pathway, Pacific cod and Pacific halibut, covary and have been relatively stable in the post-1977 period. Second, the long term increases in both Pacific halibut and arrowtooth flounder (with quite different diets apart from pollock) may be linked to similarities in their reproductive behavior. Both spawn offshore in late winter, and conditions that enhance onshore advection, such as El Niños, may play an important role in recruitment to nursery areas for these species (Bailey and Picquelle 2002).

Finally, it is apparent that the potential for competition between Steller sea lions and arrowtooth flounder is underappreciated, perhaps because arrowtooth flounder seem poorly designed to compete as forager in the pelagic zone. However, arrowtooth flounder consume both the primary prey of Steller sea lions (pollock), and alternate pelagic prey also utilized by Steller sea lions (capelin, herring, sandlance, salmon). Arrowtooth predation on pollock occurs at a smaller size than pollock targeted by Steller sea lions. The arrowtooth flounder population is nearly unexploited, is increasing in abundance, may be increasing its per unit consumption of pollock, and shows no evidence of density-dependent growth. And lastly, since 1976 there has been a strong inverse correlation between arrowtooth flounder and Steller sea lion abundance that is at least consistent with competition between these species.

Literature Cited

Alton, M., S. Hughes, and G. Hirschhorn. 1977. Gulf of Alaska pollock—its fisheries and resource potential. Unpubl. Manusc., 25 p. Alaska Fisheries Science Center. National Marine Fisheries Service National Oceanic and Atmospheric Administration. 7600 Sand Point Way, NE Seattle, Washington 98115-6349. Submitted to the International North Pacific Fisheries Commission

in 1977 as INPFC doc. 2019.

Alton, M. S., M. O. Nelson, and B. A. Megrey. 1987. Changes in the abundance and distribution of walleye pollock (*Theragra chalcogramma*) in the western Gulf of Alaska. *Fish. Res.* 5: 185-197.

Alverson, D. L. And M. J. Carney. 1975. A graphic review of the growth and decay of population cohorts. *Cons. int. Explor. Mer.* 133-143.

Anderson, P. J. and J. F. Piatt 1999. Community reorganization in the Gulf of Alaska following ocean climate regime shift. *Mar. Ecol. Prog. Ser.* 189:117-123.

Aydin, K., G.A. McFarlane, JR. King, B.A. Megrey, and K.W. Myers. 2005. Linking oceanic food webs to coastal production and growth rates of Pacific salmon (*Oncorhynchus* spp.), using models on three scales. *Deep-sea Res.* II. 52: 757-780.

Bailey, K.M., P.J. Stabeno, and D.A. Powers. 1997. The role of larval retention and transport features in mortality and potential gene flow of walleye pollock. *J. Fish. Biol.* 51(Suppl. A):135-154.

Bailey, K.M., T.J. Quinn II, P. Bentzen, and W.S. Grant. 1999. Population structure and dynamics of walleye pollock, *Theragra chalcogramma*. *Advances in Mar. Biol.* 37: 179-255.

Bailey, K.M. 2000. Shifting control of recruitment of walleye pollock *Theragra chalcogramma* after a major climatic and ecosystem change. *Mar. Ecol. Prog. Ser.* 198:215-224.

Bailey, K. M and S. J. Picquelle. 2002. Larval distribution of offshore spawning flatfish in the Gulf of Alaska: potential transport pathways and enhanced onshore transport during ENSO events. *Mar. Ecol. Prog. Ser.* 236:205-217.

Baranov, F.I. 1918. On the question of the biological basis of fisheries. *Nauchn. Issed. Ikhtiologicheskii Inst. Izv.* 1:81-128.

Blackburn, J. and D. Pengilly. 1994. A summary of estimated population trends of seven most abundant groundfish species in trawl surveys conducted by Alaska Department of Fish and Game in the Kodiak and Alaska Peninsula areas, 1988 through 1993. Alaska Department of Fish and Game, Regional Information Report No. 4K94-31. 19p.

Brodeur, R. D. and Ware, D.M. 1995. Interdecadal variability in distribution and catch rates of epipelagic nekton in the Northeast Pacific Ocean. pp. 329-356 in R. J. Beamish [Ed.] *Climate change and northern fish populations*. Canadian Special Publication of Fisheries and Aquatic Sciences 121. National Research Council of Canada, Ottawa.

Kastelle, C. R. and D. K. Kimura. 2006. Age validation of walleye pollock (*Theragra chalcogramma*) from the Gulf of Alaska using the disequilibrium of Pb-210 and Ra-226. *ICES Journal of Marine Science* 63:1520-1529.

Clark, W.G. 1999. Effects of an erroneous natural mortality rate on a simple age-structured model. *Can. J. Fish. Aquat. Sci.* 56:1721-1731.

Clark, W. G., S. R. Hare, A. M. Parma, P. J. Sullivan, and R. J. Trumble. 1999. Decadal changes in growth and recruitment of Pacific halibut (*Hippoglossus stenolepis*). *Can. J. Fish. Aquat. Sci.* 56(2): 242-252.

Cochran, W. G. 1977. *Sampling Techniques*. John Wiley and Sons. New York. 428 p.

Deriso, R.B., T.J. Quinn II, and P.R. Neal. 1985. Catch-age analysis with auxiliary information. *Can. J. Fish. Aquat. Sci.* 42: 815-824.

De Robertis, A., Hjellvik, V., Williamson, N. J., and Wilson, C. D. 2008. Silent ships do not always encounter more fish: comparison of acoustic backscatter recorded by a noise-reduced and a conventional research vessel. – *ICES Journal of Marine Science*, 65: 623–635.

Dorn, M. W. 2004. Extending separable age-structured assessment models to evaluate trends in juvenile mortality of walleye pollock in the Gulf of Alaska. *International Council for the Exploration of the Sea*, CM 2004/ FF:31.

Dorn, M. W., and R. D. Methot. 1990. Status of the coastal Pacific whiting resource in 1989 and recommendation to management in 1990. U.S. Dep. Commer., NOAA Tech. Memo. NMFS F/NWC-182, 84 p.

Dorn, M.W. , A.B. Hollowed., E. Brown, B.A. Megrey, C. Wilson and J. Blackburn. 1999. Walleye pollock. *In* Stock Assessment and Fishery Evaluation Report for Groundfish Resources of the Gulf of Alaska. Prepared by the Gulf of Alaska

Groundfish Plan Team, North Pacific Fishery Management Council, P.O. Box 103136, Anchorage, AK 99510. North Pacific Fisheries Management Council, Anchorage, AK.

Dorn, M.W., Hollowed, A.B., E. Brown, B. Megrey, C. Wilson, and J. Blackburn. 2001. Assessment of the walleye pollock stock in the Gulf of Alaska. *In* Stock Assessment and Fishery Evaluation Report for Groundfish Resources of the Gulf of Alaska. Prepared by the Gulf of Alaska Groundfish Plan Team, North Pacific Fishery Management Council, P.O. Box 103136, Anchorage, AK 99510. North Pacific Fisheries Management Council, Anchorage, AK.

Dorn, M.W., S. Barbeaux, B. M. Guttormsen, B. Megrey, A. Hollowed, M. Wilkins, and K. Spalinger. 2003. Assessment of the walleye pollock stock in the Gulf of Alaska. *In* Stock Assessment and Fishery Evaluation Report for Groundfish Resources of the Gulf of Alaska. Prepared by the Gulf of Alaska Groundfish Plan Team, North Pacific Fishery Management Council, P.O. Box 103136, Anchorage, AK 99510. North Pacific Fisheries Management Council, Anchorage, AK.

Doubleday, W.G. 1976. A least-squares approach to analyzing catch at age data. *Res. Bull. Int. Comm. Northw. Atl. Fish.* 12:69-81.

Fournier, D. and C. P. Archibald. 1982. A general theory for analyzing catch at age data. *Can. J. Fish. Aquat. Sci.* 39:1195-1207.

Fritz, L. W. 1993. Trawl locations of walleye pollock and Atka mackerel fisheries in the Bering Sea, Aleutian Islands, and Gulf of Alaska from 1977-92. AFSC Processed Report 93-08. NMFS, AFSC, 7600 Sand Point Way, NE, Seattle, WA 98115. 162 p.

Gauthier, S. and J. K. Horne 2004. Acoustic characteristics of forage fish species in the Gulf of Alaska and Bering Sea. *Can. J. Aquat. Fish. Sci.* 61: 1839-1850.

Grant, W.S. and F.M. Utter. 1980. Biochemical variation in walleye pollock *Theragra chalcogramma*: population structure in the southeastern Bering Sea and Gulf of Alaska. *Can. J. Fish. Aquat. Sci.* 37:1093-1100.

Griewank, A., and G.F. Corliss (eds.) 1991. Automatic differentiation of algorithms: theory, implementation and application. Proceedings of the SIAM Workshop on the Automatic Differentiation of Algorithms, held Jan. 6-8, Breckenridge, CO. Soc. Indust. and Applied Mathematics, Philadelphia.

Gunderson, D. R. and P. H. Dygert. 1988. Reproductive effort as a predictor of natural mortality rate. *J. Cons. int. Mer.* 44:200-209.

Guttormsen, M. A., C. D. Wilson, and S. Stienessen. 2001. Echo integration-trawl survey results for walleye pollock in the Gulf of Alaska during 2001. *In* Stock Assessment and Fishery Evaluation Report for Gulf of Alaska. Prepared by the Gulf of Alaska Groundfish Plan Team, North Pacific Fishery Management Council, P.O. Box 103136, Anchorage, AK 99510. North Pacific Fisheries Management Council, Anchorage, AK.

Guttormsen, M. A., C. D. Wilson, and S. Stienessen. 2002. Echo integration-trawl survey results for walleye pollock in the Gulf of Alaska during February and March 2002. *In* Stock Assessment and Fishery Evaluation Report for Gulf of Alaska. Prepared by the Gulf of Alaska Groundfish Plan Team, North Pacific Fishery Management Council, P.O. Box 103136, Anchorage, AK 99510. North Pacific Fisheries Management Council, Anchorage, AK.

Guttormsen, M. A., C. D. Wilson, and S. Stienessen. 2003. Results of the February and March 2003 Echo integration-trawl surveys of walleye pollock (*Theragra chalcogramma*) conducted in the Gulf of Alaska, Cruises MF2003-01 and MF2003-05. *In* Stock Assessment and Fishery Evaluation Report for Gulf of Alaska. Prepared by the Gulf of Alaska Groundfish Plan Team, North Pacific Fishery Management Council, P.O. Box 103136, Anchorage, AK 99510. North Pacific Fisheries Management Council, Anchorage, AK.

Guttormsen, M. A. 2004. Results of the March-April 2004 echo integration-trawl surveys of walleye pollock (*Theragra chalcogramma*) conducted in the Gulf of Alaska, Cruise MF0403. *In* Stock Assessment and Fishery Evaluation Report for Gulf of Alaska. Prepared by the Gulf of Alaska Groundfish Plan Team, North Pacific Fishery Management Council, P.O. Box 103136, Anchorage, AK 99510. North Pacific Fisheries Management Council, Anchorage, AK.

Guttormsen, M.A., A. McCarthy, and D. Jones. In review. Results of the February-March 2008 Echo Integration-Trawl Surveys of Walleye Pollock (*Theragra chalcogramma*) Conducted in the Gulf of Alaska, Cruises MF2008-01 and MF2008-04. AFSC Processed Rep. Alaska Fish. Sci. Cent., NOAA, Natl. Mar. Fish. Serv., 7600 Sand Point Way NE, Seattle WA 98115.

Hastie, T., and R. Tibshirani. 1990. Generalized additive models. Chapman and Hall, London. 289 pp.

- Heifetz, J., D. Anderl, N.E. Maloney, and T.L. Rutecki. 1999. Age validation and analysis of ageing error from marked and recaptured sablefish, *Anoplopoma fimbria*. Fish. Bull. 97:256-263.
- Hilborn, R. and C.J. Walters. 1992. Quantitative fisheries stock assessment: choice, dynamics, and uncertainty. Chapman and Hall, New York, N.Y. 570 p.
- Hollowed, A.B. and B.A. Megrey. 1990. Walleye pollock. In Stock Assessment and Fishery Evaluation Report for the 1991 Gulf of Alaska Groundfish Fishery. Prepared by the Gulf of Alaska Groundfish Plan Team, North Pacific Fishery Management Council, P.O. Box 103136, Anchorage, AK 99510.
- Heino, M., U. Dieckmann, and O. R. Godø. 2002a. Estimation of reaction norms for age and size at maturation with reconstructed immature size distributions: a new technique illustrated by application to Northeast Arctic cod. ICES Journal of Marine Science 59:562-575.
- Hollowed, A.B., B.A. Megrey, P. Munro, and W. Karp. 1991. Walleye pollock. In Stock Assessment and Fishery Evaluation Report for the 1992 Gulf of Alaska Groundfish Fishery. Prepared by the Gulf of Alaska Groundfish Plan Team, North Pacific Fishery Management Council, P.O. Box 103136, Anchorage, AK 99510.
- Hollowed, A. B., B.A. Megrey, and W. Karp. 1992. Walleye pollock. In Stock Assessment and Fishery Evaluation Report for the 1993 Gulf of Alaska Groundfish Fishery. Prepared by the Gulf of Alaska Groundfish Plan Team, North Pacific Fishery Management Council, P.O. Box 103136, Anchorage, AK 99510.
- Hollowed, A.B., C. Wilson, E. Brown, and B.A. Megrey. 1994. Walleye pollock. In Stock Assessment and Fishery Evaluation Report for the 1995 Gulf of Alaska Groundfish Fishery. Prepared by the Gulf of Alaska Groundfish Plan Team, North Pacific Fishery Management Council, P.O. Box 103136, Anchorage, AK 99510.
- Hollowed, A.B., E. Brown, P. Livingston, B.A. Megrey and C. Wilson. 1995. Walleye pollock. In Stock Assessment and Fishery Evaluation Report for Gulf of Alaska As Projected for 1996. Prepared by the Gulf of Alaska Groundfish Plan Team, North Pacific Fishery Management Council, P.O. Box 103136, Anchorage, AK 99510. North Pacific Fisheries Management Council, Anchorage, AK.
- Hollowed, A.B., E. Brown, J. Ianelli, B.A. Megrey and C. Wilson. 1998. Walleye pollock. In Stock Assessment and Fishery Evaluation Report for Groundfish Resources of the Gulf of Alaska. Prepared by the Gulf of Alaska Groundfish Plan Team, North Pacific Fishery Management Council, P.O. Box 103136, Anchorage, AK 99510. North Pacific Fisheries Management Council, Anchorage, AK.
- Hollowed, A.B., J.N. Ianelli, P. Livingston. 2000. Including predation mortality in stock assessments: a case study for Gulf of Alaska pollock. ICES J. Mar. Sci. 57:279-293.
- Hollowed, A.B., S. R. Hare and W. S. Wooster. 2001. Pacific Basin climate variability and patterns of Northeast Pacific marine fish production. Progress In Oceanography 49: 257-282.
- Hughes, S. E. and G. Hirschhorn. 1979. Biology of walleye pollock, *Theragra chalcogramma*, in Western Gulf of Alaska. Fish. Bull., U.S. 77:263-274.
- Karp, W. A. 1990. Results of echo integration midwater-trawl surveys for walleye pollock in the Gulf of Alaska in 1990. Appendix 3 of Stock Assessment and Fishery Evaluation Report for Groundfish Resources of the Gulf of Alaska. Prepared by the Gulf of Alaska Groundfish Plan Team, North Pacific Fishery Management Council, P.O. Box 103136, Anchorage, AK 99510. North Pacific Fisheries Management Council, Anchorage, AK.
- Kendall, A.W. Jr. and S.J. Picquelle. 1990. Egg and larval distributions of walleye pollock *Theragra chalcogramma* in Shelikof Strait, Gulf of Alaska. Fish. Bull., U.S. 88:133-154.
- Kimura, D.K. 1989. Variability, tuning, and simulation for the Doubleday-Deriso catch-at-age model. Can. J. Fish. Aquat. Sci. 46:941-949.
- Kimura, D.K. 1990. Approaches to age-structured separable sequential population analysis. Can. J. Fish. Aquat. Sci. 47:2364-2374.
- Kimura, D.K. 1991. Improved methods for separable sequential population analysis. Unpublished. Alaska Fisheries Science Center, 7600 Sand Point Way NE, Seattle, Washington 98115.

- Kimura, D. K. and S. Chikuni. 1989. Variability in estimating catch-in-numbers-at-age and its impact on cohort analysis. In R.J. Beamish and G.A. McFarlane (eds.), Effects of ocean variability on recruitment and an evaluation of parameters used in stock assessment models. *Can. Spec. Publ. Fish. Aquat. Sci.* 108:57-66.
- Large, W.G., and S. Pond (1982) Sensible and latent heat flux measurement over the ocean. *J. Phys. Oceanogr.* 2: 464-482.
- Lee, Y-W, Megrey, B.A., and Macklin, S.A. in prep. Using a resampling strategy to evaluate Gulf of Alaska walleye pollock recruitment forecast performance.
- McCullagh, P., and J. A. Nelder. 1983. Generalized linear models. Chapman and Hall, London. 261 p.
- McKelvey, D. 1996. Juvenile walleye pollock, *Theragra chalcogramma*, distribution and abundance in Shelikof Strait—what can we learn from acoustic survey results? p. 25-34. In U.S. Dep. Commer. NOAA Tech. Rep. NMFS 126.
- McKelvey, D.R. 1996. Juvenile walleye pollock, *Theragra chalcogramma*, distribution and abundance in Shelikof Strait—what can we learn from acoustic surveys. Ecology of Juvenile Walleye Pollock, *Theragra chalcogramma*. NOAA Technical Report NMFS 126, p 25-34.
- Macklin, S.A., R.L. Brown, J. Gray, and R.W. Lindsay (1984) METLIB-II - A program library for calculating and plotting atmospheric and oceanic fields. NOAA Tech. Memo. ERL PMEL-54, NTIS PB84-205434, 53 pp.
- Macklin, S.A., P.J. Stabeno, and J.D. Schumacher (1993) A comparison of gradient and observed over-the-water winds along a mountainous coast. *J. Geophys. Res.* 98: 16,555–16,569.
- Megrey, B.A. 1989. Exploitation of walleye pollock resources in the Gulf of Alaska, 1964-1988: portrait of a fishery in transition. Proc. International Symp. on the Biology and Management of Walleye Pollock, Lowell Wakefield Fisheries Symp., Alaska Sea Grant Rep. 89-1, 33-58.
- Megrey, B.A. and Macklin, S.A. unpublished. Critical Analysis of FOCI Walleye Pollock Recruitment Prediction for the Gulf of Alaska, 1992-2005.
- Megrey, B.A., Lee, Y-W, and Macklin, S.A. 2005. Comparative analysis of statistical tools to identify recruitment-environment relationships and forecast recruitment strength. *ICES Journal of Marine Science* 62(7): 1256-1269.
- Merati, N. 1993. Spawning dynamics of walleye pollock, *Theragra chalcogramma*, in Shelikof Strait, Gulf of Alaska. Unpublished MS thesis. University of Washington. 134 p.
- Martin, M.H. 1997. Data Report: 1996 Gulf of Alaska bottom trawl survey. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-AFSC-82, 235 p.
- Methot, R.D. 2000. Technical description of the stock synthesis assessment program. U.S. Dept. Commer., NOAA Tech. Memo. NMFS-NWFSC-43, 46 p.
- Meuter, F.J. and B.L. Norcross. 2002. Spatial and temporal patterns in the demersal fish community on the shelf and upper slope regions of the Gulf of Alaska. *Fish. Bull.* 100:559-581.
- Mulligan, T.J., Chapman, R.W. and B.L. Brown. 1992. Mitochondrial DNA analysis of walleye pollock, *Theragra chalcogramma*, from the eastern Bering Sea and Shelikof Strait, Gulf of Alaska. *Can. J. Fish. Aquat. Sci.* 49:319-326.
- Olsen, J.B., S.E. Merkouris, and J.E. Seeb. 2002. An examination of spatial and temporal genetic variation in walleye pollock (*Theragra chalcogramma*) using allozyme, mitochondrial DNA, and microsatellite data. *Fish. Bull.* 100:752-764.
- Parada, C. Hinckley, S., Dorn, M., Hermann, A.J., Megrey, B.A. submitted. Estimating walleye pollock recruitment in the Gulf of Alaska using a biophysical model: Analysis of physical processes and comparison with stock assessment models ad data. Marine Ecology Progress Series.
- Pauly, D.. 1980. On the interrelationships between natural mortality, growth parameters, and mean environmental temperature in 175 fish stocks. *J. Cons. int. Explor. Mer.* 39(2):175-192.
- Press, W.H., S.A. Teukolsky, W.T. Vetterling, and B.P. Flannery. 1992. Numerical recipes in C. Second ed. Cambridge University Press. 994 p.

- Picquelle, S.J., and B.A. Megrey. 1993. A preliminary spawning biomass estimate of walleye pollock, *Theragra chalcogramma*, in Shelikof Strait, Gulf of Alaska, based on the annual egg production method. *Bulletin of Marine Science* 53(2):728-749.
- Ronholt, L. L., H. H. Shippen, and E. S. Brown. 1978. Demersal fish and shellfish resources of the Gulf of Alaska from Cape Spencer to Unimak Pass 1948 - 1976 (A historical review). Northwest and Alaska Fisheries Center Processed Report.
- Rothschild, B. J. and Mullin, A.J. 1985. The information content of stock-and-recruitment data and its non-parametric classification. *Journal du Conseil International pour l'Exploration de la Mer*. 42: 116-124.
- Saunders, M.W., G.A. McFarlane, and W. Shaw. 1988. Delineation of walleye pollock (*Theragra chalcogramma*) stocks off the Pacific coast of Canada. Proc. International Symp. on the Biology and Management of Walleye Pollock, Lowell Wakefield Fisheries Symp., Alaska Sea Grant Rep. 89-1, 379-402.
- Schnute, J.T. and L.J. Richards. 1995. The influence of error on population estimates from catch-age models. *Can. J. Fish. Aquat. Sci.* 52:2063-2077.
- Smith, S. J. 1990. Use of statistical models for the estimation of abundance from groundfish trawl survey data. *Can. J. Fish. Aquat. Sci.* 47:894-903.
- Somerton, D. 1979. Competitive interaction of walleye pollock and Pacific Ocean perch in the northern Gulf of Alaska. In S. J. Lipovsky and C.A. Simenstad (eds.) *Gutshop '78, Fish food habits studies: Proceedings of the second Pacific Northwest Technical Workshop, held Maple Valley, WA (USA), 10-13 October, 1978.*, Washington Sea Grant, Seattle, WA.
- StatSci. 1993. S-Plus for DOS reference manual. Statistical Sciences Inc., Seattle, Wash.
- Sullivan, P.J., A.M. Parma, and W.G. Clark. 1997. Pacific halibut assessment: data and methods. *Int. Pac. Halibut Comm. SCI. Rept.* 97. 84 p.
- Swartzman, G., C. Huang, and S. Kaluzny. 1992. Spatial analysis of Bering Sea groundfish survey data using generalized additive models. *Can. J. Fish. Aquat. Sci.* 49:1366-1378.
- von Szalay, P. G., and E. Brown. 2001. Trawl comparisons of fishing power differences and their applicability to National Marine Fisheries Service and the Alaska Department of Fish and Game trawl survey gear. *Alaska Fishery Research Bulletin* 8:85-95.
- Wilson, C. 1994. Echo integration-trawl survey results for pollock in the Gulf of Alaska during 1994. Appendix D of Stock Assessment and Fishery Evaluation Report for Groundfish Resources of the Gulf of Alaska. Prepared by the Gulf of Alaska Groundfish Plan Team, North Pacific Fishery Management Council, P.O. Box 103136, Anchorage, AK 99510. North Pacific Fisheries Management Council, Anchorage, AK.
- Wilson, C., M. Guttormsen, and D. McKelvey. 1995. Echo integration-trawl survey results for pollock in the Gulf of Alaska during 1995. Appendix D of Stock Assessment and Fishery Evaluation Report for Groundfish Resources of the Gulf of Alaska. Prepared by the Gulf of Alaska Groundfish Plan Team, North Pacific Fishery Management Council, P.O. Box 103136, Anchorage, AK 99510. North Pacific Fisheries Management Council, Anchorage, AK.
- Wilson, C.D., M.A. Guttormsen, and S.K. de Blois. 1996. Echo integration-trawl survey results for pollock in the Gulf of Alaska during 1996. In Stock Assessment and Fishery Evaluation Report for Groundfish Resources of the Gulf of Alaska. Prepared by the Gulf of Alaska Groundfish Plan Team, North Pacific Fishery Management Council, P.O. Box 103136, Anchorage, AK 99510. North Pacific Fisheries Management Council, Anchorage, AK.
- Yang, M-S. and M. W. Nelson. 2000. Food habits of the commercially important groundfishes in the Gulf of Alaska in 1990, 1993, and 1996. *U.S. Dep. Commer., NOAA Tech. Memo. NMFS-AFSC-112*, 174 p.
- Zeppelin, TK., DJ. Tollit, KA. Call, TJ. Orchard, and CJ. Gudmundson. 2004. Sizes of walleye pollock (*Theragra chalcogramma*) and Atka mackerel (*Pleurogrammus monopterygius*) consumed by the western stock of Steller sea lions (*Eumetopias jubatus*) in Alaska from 1998 to 2000. *Fish. Bull.* 102:509-521.

Table 1.1. Walleye pollock catch (t) in the Gulf of Alaska. The TAC for 2008 is for the area west of 140 ° W lon. (Western, Central and West Yakutat management areas) and includes the guideline harvest level for the state-managed fishery in Prince William Sound (1650 t). Research catches are also reported.

<i>Year</i>	<i>Foreign</i>	<i>Joint Venture</i>	<i>Domestic</i>	<i>Total</i>	<i>TAC</i>	<i>Research</i>
1964	1,126			1,126	---	
1965	2,749			2,749	---	
1966	8,932			8,932	---	
1967	6,276			6,276	---	
1968	6,164			6,164	---	
1969	17,553			17,553	---	
1970	9,343			9,343	---	
1971	9,458			9,458	---	
1972	34,081			34,081	---	
1973	36,836			36,836	---	
1974	61,880			61,880	---	
1975	59,512			59,512	---	
1976	86,527			86,527	---	
1977	117,834		522	118,356	150,000	75
1978	96,392	34	509	96,935	168,800	100
1979	103,187	566	1,995	105,748	168,800	52
1980	112,997	1,136	489	114,622	168,800	229
1981	130,324	16,857	563	147,744	168,800	433
1982	92,612	73,917	2,211	168,740	168,800	110
1983	81,358	134,131	119	215,608	256,600	213
1984	99,260	207,104	1,037	307,401	416,600	311
1985	31,587	237,860	15,379	284,826	305,000	167
1986	114	62,591	25,103	87,809	116,000	1202
1987		22,823	46,928	69,751	84,000	227
1988		152	65,587	65,739	93,000	19
1989			78,392	78,392	72,200	73
1990			90,744	90,744	73,400	158
1991			100,488	100,488	103,400	16
1992			90,857	90,857	87,400	40
1993			108,908	108,908	114,400	116
1994			107,335	107,335	109,300	70
1995			72,618	72,618	65,360	44
1996			51,263	51,263	54,810	147
1997			90,130	90,130	79,980	76
1998			125,098	125,098	124,730	64
1999			95,590	95,590	94,580	35
2000			73,080	73,080	94,960	56
2001			72,076	72,076	90,690	77
2002			51,937	51,937	53,490	78
2003			50,666	50,666	49,590	128
2004			63,934	63,934	65,660	53
2005			80,846	80,846	86,100	72
2006			71,976	71,976	81,300	63
2007			53,062	53,062	63,800	47
2008			52,500	52,500	53,590	26
2009					43,270	87
<i>Average (1977-2008)</i>				106,848	123,560	143

Sources: 1964-85--Megrey (1988); 1986-90--Pacific Fishery Information Network (PacFIN), Pacific Marine Fisheries Commission. Domestic catches in 1986-90 were adjusted for discard as described in Hollowed et al. (1991). 1991-2008 -- NMFS Alaska Regional Office.

Table 1.2. Incidental catch (t) of FMP species (upper table) and non-target species (bottom table) in the walleye pollock directed fishery in the Gulf of Alaska in 2004-2008. Incidental catch estimates include both retained and discarded catch. The "other" FMP species group in the upper table is broken down by species (or less inclusive species groupings) in the lower table.

<i>Managed species/species group</i>	<i>2004</i>	<i>2005</i>	<i>2006</i>	<i>2007</i>	<i>2008</i>
Pollock	62834.3	80097.8	69774.8	49815.5	46735.3
Arrowtooth flounder	1162.7	2313.4	2747.5	1630.0	1554.6
Pacific cod	393.5	352.3	709.8	276.4	578.7
Other (sharks, skates, squid, sculpin, octopus, but excluding skates in 2004)	368.6	924.6	1805.5	676.9	200.6
Flathead sole	270.6	180.2	594.4	329.6	414.0
Shortraker and roughey rockfish	48.6	32.6	96.5	81.4	101.5
Pacific Ocean perch	82.7	35.5	71.2	29.8	49.9
Rex sole	35.3	21.1	153.6	44.8	57.4
Miscellaneous flatfish	9.8	4.6	438.8	157.0	230.2
Atka mackerel	17.9	3.5	15.2	200.2	0.1
Sablefish	4.8	3.6	5.6	3.2	1.3
Dover sole and Greenland turbot	1.7	0.7	11.7	5.5	5.8
Pelagic shelf rockfish complex	2.0	2.1	9.0	6.4	4.1
Unidentified skate	1.8	1.2	5.0	9.4	5.9
Big and longnose skate	1.5	6.7	35.8	64.8	45.3
Northern rockfish	0.5	0.8	14.5	12.0	7.9
Other rockfish complex	0.1	1.3	2.5	2.0	4.5
Thornyheads	0.0	0.3	0.2	0.3	0.2
<i>Percent non-pollock</i>	<i>3.7%</i>	<i>4.6%</i>	<i>8.8%</i>	<i>6.6%</i>	<i>6.5%</i>

<i>Non target species/species group</i>	<i>2004</i>	<i>2005</i>	<i>2006</i>	<i>2007</i>	<i>2008</i>
Squid	144.5	631.2	1517.7	405.2	77.9
Eulachon	168.8	824.8	392.3	219.0	750.9
Other osmerids	66.1	175.3	167.9	49.2	378.4
Pacific sleeper shark	170.3	198.8	154.5	58.8	46.9
Scyphozoan jellyfish	22.5	184.4	69.0	23.9	188.7
Grenadiers	7.7	53.8	73.1	4.7	249.3
Spiny dogfish	9.2	15.8	50.0	47.5	59.8
Other sharks	11.1	35.2	40.9	13.7	4.3
Miscellaneous fish	13.9	16.5	38.4	24.1	35.2
Salmon shark	33.1	43.3	31.4	125.1	6.2
Big skate	0.8	1.7	23.0	24.0	11.0
Longnose skate	0.3	4.4	12.3	21.3	17.1
Octopus	0.0	0.1	3.4	1.5	0.0
Pandalid shrimp	1.5	7.4	3.1	1.9	0.8
Other skates	1.7	1.0	2.1	2.4	3.3
Sea star	0.0	1.1	2.0	4.7	6.6
Sculpins	0.1	0.0	2.4	21.8	15.4
Sea anemone unidentified	0.1	0.0	0.2	0.7	0.3
Capelin	68.0	2.7	0.1	0.0	0.0
Stichaeidae	0.1	0.0	0.1	0.3	0.0
Lanternfishes (myctophidae)	0.0	0.1	0.0	0.0	0.0
Invertebrate unidentified	0.0	0.0	0.0	0.2	0.0
Sea pens whips	0.0	0.3	0.0	0.0	0.0
Greenlings	0.0	0.0	0.0	0.0	0.0
Misc crabs	0.0	0.0	0.0	0.9	0.1
Eelpouts	1.3	0.1	0.0	0.0	0.0

Table 1.3. Bycatch of prohibited species for trawls in the Gulf of Alaska during 2004-2008 where pollock was the predominant species in the catch. Herring and halibut bycatch is reported in metric tons, while crab and salmon are reported in number of fish.

<i>Species/species group</i>	<i>2004</i>	<i>2005</i>	<i>2006</i>	<i>2007</i>	<i>2008</i>
Herring (t)	269.741	12.163	8.789	19.529	0.421
Halibut (t)	23.026	3.833	115.576	135.392	120.041
Bairdi Tanner Crab (nos.)	1,222	6	84,005	19,458	322
Red King Crab (nos.)	56	0	0	0	0
Chinook Salmon (nos.)	13,152	27,910	15,943	35,042	10,382
Non-chinook salmon (nos.)	585	781	1,413	982	847

Table 1.4. Catch (retained and discarded) of walleye pollock (t) by management area in the Gulf of Alaska during 1999-2008 compiled by the Alaska Regional Office.

Year	Utilization	Shumagin 610	Chirikof 620	Kodiak 630	West Yakutat 640	Prince William Sound 649 (state waters)		Southeast and East Yakutat 650 & 659	Total	Percent discard
						Sound 649 (state waters)	East Yakutat 650 & 659			
1999	Retained	22,864	37,349	29,515	1,737	2,178	1	93,643		
	Discarded	521	784	578	22	39	3	1,947	2.0%	
	Total	23,385	38,133	30,093	1,759	2,216	4	95,590		
2000	Retained	21,380	11,314	35,078	1,917	1,181	0	70,870		
	Discarded	694	443	854	191	22	4	2,209	3.0%	
	Total	22,074	11,757	35,933	2,108	1,203	4	73,080		
2001	Retained	30,298	17,186	19,942	2,327	1,590	0	71,344		
	Discarded	173	205	330	24	0	0	732	1.0%	
	Total	30,471	17,391	20,272	2,351	1,590	0	72,076		
2002	Retained	17,046	20,106	10,615	1,808	1,216	0	50,791		
	Discarded	416	425	287	10	6	2	1,146	2.2%	
	Total	17,462	20,531	10,902	1,818	1,222	2	51,937		
2003	Retained	16,347	18,972	12,225	940	1,118	0	49,603		
	Discarded	161	658	210	2	31	0	1,063	2.1%	
	Total	16,508	19,630	12,435	943	1,149	0	50,666		
2004	Retained	23,226	24,221	13,896	215	1,100	0	62,658		
	Discarded	342	438	459	11	26	0	1,276	2.0%	
	Total	23,568	24,659	14,355	226	1,127	0	63,934		
2005	Retained	30,791	27,286	18,986	1,876	740	0	79,680		
	Discarded	136	621	350	9	50	0	1,166	1.4%	
	Total	30,927	27,908	19,336	1,885	790	0	80,846		
2006	Retained	24,489	26,409	16,127	1,570	1,475	0	70,070		
	Discarded	203	750	951	2	1	0	1,906	2.6%	
	Total	24,691	27,159	17,078	1,572	1,476	0	71,976		
2007	Retained	17,694	18,846	13,777	84	NA	0	50,401		
	Discarded	262	516	701	3	NA	1	1,483	2.8%	
	Total	17,956	19,362	14,478	87	1,179	1	53,062		
2008	Retained	15,100	18,691	13,335	1,155	NA	0	48,281		
	Discarded	2,157	367	1,052	6	NA	2	3,584	6.8%	
	Total	17,257	19,058	14,387	1,161	635	2	52,500		
<i>Average (1999-2008)</i>		22,430	22,559	18,927	1,391	1,259	1	66,567		

Table 1.5. Catch at age (000,000s) of walleye pollock in the Gulf of Alaska in 1976-2008.

Year	Age															
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	Total
1976	0.00	1.91	24.21	108.69	39.08	16.37	3.52	2.25	1.91	0.31	0.00	0.00	0.00	0.00	0.00	198.25
1977	0.01	2.76	7.06	23.83	89.68	30.35	8.33	2.13	1.79	0.67	0.44	0.10	0.02	0.00	0.00	167.17
1978	0.08	12.11	48.32	18.26	26.39	51.86	12.83	4.18	1.36	1.04	0.32	0.04	0.01	0.00	0.00	176.80
1979	0.00	2.53	48.83	76.37	14.15	10.13	16.70	5.02	1.27	0.60	0.16	0.04	0.00	0.00	0.00	175.81
1980	0.25	19.01	26.50	58.31	36.63	11.31	8.61	8.00	3.89	1.11	0.50	0.21	0.08	0.03	0.00	174.42
1981	0.14	2.59	31.55	73.91	47.97	20.29	4.87	4.83	2.73	0.26	0.03	0.02	0.00	0.00	0.00	189.19
1982	0.01	10.67	55.55	100.77	71.73	54.25	10.46	1.33	0.93	0.55	0.03	0.02	0.02	0.00	0.00	306.31
1983	0.00	3.64	20.64	110.03	137.31	67.41	42.01	7.38	1.24	0.06	0.28	0.07	0.00	0.00	0.00	390.07
1984	0.34	2.37	33.00	38.80	120.80	170.72	62.55	19.31	5.42	0.10	0.07	0.03	0.03	0.00	0.00	453.54
1985	0.04	12.74	5.53	33.22	42.22	86.02	128.95	41.19	10.84	2.20	0.70	0.00	0.00	0.00	0.00	363.64
1986	0.66	8.63	20.34	10.12	19.13	7.32	8.70	9.78	2.13	0.80	0.00	0.00	0.00	0.00	0.00	87.59
1987	0.00	8.83	14.03	8.00	6.89	6.44	7.18	4.19	9.95	1.94	0.00	0.00	0.00	0.00	0.00	67.44
1988	0.17	3.05	20.80	26.95	11.94	5.10	3.45	1.62	0.34	3.21	0.00	0.00	0.00	0.00	0.00	76.62
1989	1.08	0.27	1.47	19.39	28.89	16.96	8.09	4.76	1.69	1.10	3.62	0.43	0.01	0.00	0.00	87.77
1990	0.00	2.77	2.40	2.99	9.49	40.39	13.06	4.90	1.08	0.41	0.01	0.56	0.01	0.07	0.06	78.20
1991	0.00	0.59	9.68	5.45	2.85	5.33	26.67	3.12	16.10	0.87	5.65	0.42	2.19	0.21	0.77	79.90
1992	0.05	3.25	5.57	50.61	14.13	4.02	8.77	19.55	1.02	1.49	0.20	0.73	0.00	0.00	0.00	109.41
1993	0.02	1.97	9.43	21.83	47.46	15.72	6.55	6.29	8.52	1.81	2.07	0.49	0.72	0.13	0.24	123.25
1994	0.06	1.26	4.49	9.63	35.92	31.32	12.20	4.84	4.60	6.15	1.44	1.02	0.29	0.09	0.08	113.37
1995	0.00	0.06	1.01	5.11	11.52	25.83	12.09	2.99	1.52	2.00	1.82	0.19	0.28	0.03	0.15	64.61
1996	0.00	1.27	1.37	1.12	3.50	5.11	12.87	10.60	3.14	1.53	0.80	1.43	0.35	0.23	0.16	43.48
1997	0.00	1.07	6.72	3.77	3.28	6.60	10.09	16.52	12.24	5.06	2.06	0.79	0.54	0.17	0.02	68.92
1998	0.31	0.27	26.44	36.44	15.06	6.65	7.50	11.36	14.96	10.76	3.75	0.75	0.38	0.21	0.11	134.95
1999	0.00	0.42	2.21	22.74	36.10	8.99	6.89	3.72	5.71	7.27	4.01	1.07	0.56	0.12	0.10	99.92
2000	0.08	0.98	2.84	3.47	14.65	24.63	6.24	5.05	2.30	1.24	3.00	1.52	0.30	0.14	0.04	66.48
2001	0.74	10.13	6.59	7.34	9.42	12.59	14.44	4.73	2.70	1.35	0.65	0.83	0.61	0.00	0.04	72.14
2002	0.16	12.31	20.72	6.76	4.47	8.75	5.37	6.06	1.33	0.82	0.43	0.30	0.33	0.22	0.13	68.16
2003	0.14	2.69	21.47	22.95	5.33	3.25	4.66	3.76	2.58	0.54	0.19	0.04	0.09	0.04	0.05	67.79
2004	0.85	6.28	11.91	31.84	25.09	5.98	2.43	2.63	0.77	0.22	0.25	0.00	0.00	0.00	0.00	88.24
2005	1.14	1.21	5.33	6.85	41.25	21.73	6.10	0.74	0.91	0.35	0.18	0.13	0.00	0.00	0.00	85.91
2006	2.20	7.79	4.16	2.75	5.97	27.38	12.80	2.45	0.83	0.46	0.23	0.10	0.07	0.03	0.00	67.22
2007	0.82	18.89	7.46	2.51	2.31	3.58	10.19	6.70	1.59	0.29	0.23	0.09	0.00	0.00	0.01	54.68
2008	0.32	6.29	21.94	6.76	2.15	1.16	2.27	5.60	2.84	0.87	0.36	0.21	0.06	0.04	0.02	50.89

Table 1.6. Number of aged and measured fish in the Gulf of Alaska pollock fishery used to estimate fishery age composition (1989-2008).

<i>Year</i>	<i>Number aged</i>			<i>Number measured</i>		
	<i>Males</i>	<i>Females</i>	<i>Total</i>	<i>Males</i>	<i>Females</i>	<i>Total</i>
1989	882	892	1,774	6,454	6,456	12,910
1990	453	689	1,142	17,814	24,662	42,476
1991	1,146	1,322	2,468	23,946	39,467	63,413
1992	1,726	1,755	3,481	31,608	47,226	78,834
1993	926	949	1,875	28,035	31,306	59,341
1994	136	129	265	24,321	25,861	50,182
1995	499	544	1,043	10,591	10,869	21,460
1996	381	378	759	8,581	8,682	17,263
1997	496	486	982	8,750	8,808	17,558
1998	924	989	1,913	78,955	83,160	162,115
1999	980	1,115	2,095	16,304	17,964	34,268
2000	1,108	972	2,080	13,167	11,794	24,961
2001	1,063	1,025	2,088	13,731	13,552	27,283
2002	1,036	1,025	2,061	9,924	9,851	19,775
2003	1,091	1,119	2,210	8,375	8,220	16,595
2004	1,217	996	2,213	4,446	3,622	8,068
2005	1,065	968	2,033	6,837	6,005	12,842
2006	1,127	969	2,096	7,248	6,178	13,426
2007	998	1,064	2,062	4,504	5,064	9,568
2008	961	1,090	2,051	7,430	8,536	15,966

Table 1.7. Biomass estimates (t) of walleye pollock from NMFS echo integration trawl surveys in Shelikof Strait, NMFS bottom trawl surveys (west of 140 W. long.), egg production surveys in Shelikof Strait, and ADF&G crab/groundfish trawl surveys. The biomass of age-1 fish is not included in Shelikof Strait EIT survey estimates in 1995, 2000, 2005 and 2008 (114,200, 57,300, 18,100 t and 19,090 t respectively). An adjustment of +1.05% was made to the AFSC bottom trawl biomass time series to account for unsurveyed biomass in Prince William Sound. In 2001, when the NMFS bottom trawl survey did not extend east of 147° W lon., an expansion factor of 2.7% derived from previous surveys was used for West Yakutat.

<i>EIT Shelikof Strait survey</i>							
<i>Year</i>	<i>R/V Miller Freeman</i>		<i>R/V Oscar</i>		<i>NMFS bottom</i>	<i>Shelikof Strait</i>	<i>ADF&G</i>
	<i>Biosonics</i>	<i>EK500</i>	<i>Dyson</i>	<i>140° W lon.</i>	<i>trawl west of</i>	<i>egg</i>	<i>crab/groundfish</i>
					<i>140° W lon.</i>	<i>production</i>	<i>survey</i>
1981	2,785,755					1,788,908	
1982							
1983	2,278,172						
1984	1,757,168				720,548		
1985	1,175,823					768,419	
1986	585,755					375,907	
1987					732,660	484,455	
1988	301,709					504,418	
1989	290,461					433,894	214,434
1990	374,731				825,609	381,475	114,451
1991	380,331					370,000	
1992	580,000	713,429				616,000	127,359
1993	295,785	435,753			755,786		132,849
1994		492,593					103,420
1995		649,401					
1996		777,172			666,521		122,477
1997		583,017					93,728
1998		504,774					81,215
1999					607,409		53,587
2000		391,327					102,871
2001		432,749			219,072		86,967
2002		256,743					96,237
2003		317,269			398,469		66,989
2004		330,753					99,358
2005		338,038			358,017		79,089
2006		293,609					69,044
2007		180,881			282,356		76,674
2008			188,942				83,476
2009			265,971		669,505		145,438

Table 1.8. Survey sampling effort and biomass coefficients of variation (CV) for pollock in the Gulf of Alaska bottom trawl survey. The number of measured pollock is approximate due to subsample expansions in the database, and the total number measured includes both sexed and unsexed fish.

Year	No. of tows	No. of tows with pollock	Survey biomass CV	Number aged			Number measured		
				Males	Females	Total	Males	Females	Total
1984	929	536	0.14	1,119	1,394	2,513	8,979	13,286	24,064
1987	783	533	0.20	672	675	1,347	8,101	15,654	24,608
1990	708	549	0.12	503	560	1,063	13,955	18,967	35,355
1993	775	628	0.16	879	1,013	1,892	14,496	18,692	34,921
1996	807	668	0.15	509	560	1,069	14,653	15,961	34,526
1999	764	567	0.38	560	613	1,173	10,808	11,314	24,080
2001	489	302	0.30	395	519	914	NA	NA	NA
2003	807	508	0.12	514	589	1,103	NA	NA	NA
2005	839	516	0.15	639	868	1,507	NA	NA	NA
2007	820	554	0.14	646	675	1,321	NA	NA	NA
2009	823	563	0.15	NA	NA	NA	NA	NA	NA

Table 1.9. Estimated number at age (000,000s) from the NMFS bottom trawl survey. Estimates are for the Western and Central Gulf of Alaska only (Management areas 610-630). Ages for the 2009 NMFS bottom trawl survey are not yet available.

Year	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	Total
1984	0.93	10.02	67.81	155.78	261.17	474.57	145.10	24.80	16.59	1.66	0.21	1.32	0.00	0.00	0.00	1159.96
1987	25.45	363.02	172.99	138.97	91.13	168.27	78.14	43.99	175.39	22.41	7.81	3.51	1.82	0.00	0.00	1292.88
1989	208.88	63.49	47.56	243.15	301.09	104.43	54.47	28.39	26.14	5.98	10.66	0.00	0.00	0.00	0.00	1094.23
1990	64.04	251.21	48.34	46.68	209.77	240.82	74.41	110.41	26.13	34.23	5.03	27.73	5.70	1.07	1.63	1147.19
1993	139.31	71.15	50.94	182.96	267.12	91.51	33.12	68.98	76.62	26.36	11.85	6.29	3.82	1.82	4.41	1036.25
1996	194.23	128.79	17.30	26.13	50.04	63.18	174.41	87.62	52.37	27.73	12.10	18.46	7.16	9.68	19.70	888.90
1999	109.73	19.17	20.94	66.76	118.94	56.80	59.04	47.71	56.40	81.97	65.18	9.67	8.28	2.50	0.76	723.85
2001	412.83	117.03	34.42	33.39	25.05	33.45	37.01	8.20	5.74	0.59	4.48	2.52	1.28	0.00	0.18	716.19
2003	75.46	18.40	128.41	140.74	73.27	44.72	36.10	25.27	14.51	8.61	3.23	1.79	1.26	0.00	0.00	571.77
2005	270.37	33.72	34.41	35.86	91.78	78.82	45.24	20.86	9.61	9.98	4.81	0.57	0.64	0.00	0.00	636.68
2007	174.01	95.96	88.59	37.11	19.23	18.90	54.98	31.11	6.64	3.04	2.78	1.00	1.13	0.00	0.00	534.48

Table 1.10. Estimated number at age (000,000s) from the echo integration-trawl survey in Shelikof Strait. For the acoustic survey in 1987, when total abundance could not be estimated, the percent at age is given.

Year	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	Total
1981	77.65	3,481.18	1,510.77	769.16	2,785.91	1,051.92	209.93	128.52	79.43	25.19	1.73	0.00	0.00	0.00	0.00	10,121.37
1983	1.21	901.77	380.19	1,296.79	1,170.81	698.13	598.78	131.54	14.48	11.61	3.92	1.71	0.00	0.00	0.00	5,210.93
1984	61.65	58.25	324.49	141.66	635.04	988.21	449.62	224.35	41.03	2.74	0.00	1.02	0.00	0.00	0.00	2,928.07
1985	2,091.74	544.44	122.69	314.77	180.53	347.17	439.31	166.68	42.72	5.56	1.77	1.29	0.00	0.00	0.00	4,258.67
1986	575.36	2,114.83	183.62	45.63	75.36	49.34	86.15	149.36	60.22	10.62	1.29	0.00	0.00	0.00	0.00	3,351.78
1987	7.5%	25.5%	55.8%	2.9%	1.7%	1.2%	1.6%	1.2%	2.1%	0.4%	0.1%	0.0%	0.0%	0.0%	0.0%	100.0%
1988	17.44	109.93	694.32	322.11	77.57	16.99	5.70	5.60	3.98	8.96	1.78	1.84	0.20	0.00	0.00	1,266.41
1989	399.48	89.52	90.01	222.05	248.69	39.41	11.75	3.83	1.89	0.55	10.66	1.42	0.00	0.00	0.00	1,119.25
1990	49.14	1,210.17	71.69	63.37	115.92	180.06	46.33	22.44	8.20	8.21	0.93	3.08	1.51	0.79	0.24	1,782.08
1991	21.98	173.65	549.90	48.11	64.87	69.60	116.32	23.65	29.43	2.23	4.29	0.92	4.38	0.00	0.00	1,109.32
1992	228.03	33.69	73.54	188.10	367.99	84.11	84.99	171.18	32.70	56.35	2.30	14.67	0.90	0.30	0.00	1,338.85
1993	63.29	76.08	37.05	72.39	232.79	126.19	26.77	35.63	38.72	16.12	7.77	2.60	2.19	0.49	1.51	739.61
1994	185.98	35.77	49.30	31.75	155.03	83.58	42.48	27.23	44.45	48.46	14.79	6.65	1.12	2.34	0.57	729.49
1995	10,689.87	510.37	79.37	77.70	103.33	245.23	121.72	53.57	16.63	10.72	14.57	5.81	2.12	0.44	0.00	11,931.45
1996	56.14	3,307.21	118.94	25.12	53.99	71.03	201.05	118.52	39.80	13.01	11.32	5.32	2.52	0.03	0.38	4,024.36
1997	70.37	183.14	1,246.55	80.06	18.42	44.04	51.73	97.55	52.73	14.29	2.40	3.05	0.93	0.46	0.00	1,865.72
1998	395.47	88.54	125.57	474.36	136.12	14.22	31.93	36.30	74.08	25.90	14.30	6.88	0.27	0.56	0.56	1,425.05
2000	4,484.41	755.03	216.52	15.83	67.19	131.64	16.82	12.61	9.87	7.84	13.87	6.88	1.88	1.06	0.00	5,741.46
2001	288.93	4,103.95	351.74	61.02	41.55	22.99	34.63	13.07	6.20	2.67	1.20	1.91	0.69	0.50	0.24	4,931.27
2002	8.11	162.61	1,107.17	96.58	16.25	16.14	7.70	6.79	1.46	0.66	0.35	0.34	0.15	0.13	0.00	1,424.45
2003	51.19	89.58	207.69	802.46	56.58	7.69	4.14	1.58	1.46	0.85	0.28	0.00	0.10	0.00	0.00	1,223.60
2004	52.58	93.94	57.58	159.62	356.33	48.78	2.67	3.42	3.32	0.52	0.42	0.00	0.66	0.00	0.00	779.84
2005	1,626.13	157.49	55.54	34.63	172.74	162.40	36.02	3.61	2.39	0.00	0.76	0.00	0.00	0.00	0.00	2,251.71
2006	161.69	835.96	40.75	11.54	17.42	55.98	74.97	32.25	6.90	0.83	0.75	0.53	0.00	0.00	0.00	1,239.57
2007	53.54	231.73	174.88	29.66	10.14	17.27	34.39	20.85	1.54	1.05	0.69	0.00	0.00	0.00	0.00	575.74
2008	1,368.02	391.20	249.56	53.18	12.01	2.16	4.07	10.66	6.69	2.01	0.53	0.00	0.00	0.00	0.00	2,100.10
2009	331.94	1,204.50	110.22	98.69	60.21	9.91	2.90	0.86	5.07	6.13	1.37	0.24	0.00	0.00	0.00	1,832.03

Table 1.11. Survey sampling effort and biomass coefficients of variation (CV) for pollock in the Shelikof Strait EIT survey. Survey CVs are reported for 1981-91, while relative estimation error using a geostatistical method are reported for 1992-2009.

Year	No. of midwater tows		No. of bottom trawl tows	Survey biomass		Number aged		Number measured	
	tows			CV		Males	Females	Total	Males
1981	36	18	0.12	1,921	1,815	3,736	NA	NA	NA
1983	47	1	0.16	1,642	1,103	2,745	NA	NA	NA
1984	42	0	0.18	1,739	1,622	3,361	NA	NA	NA
1985	57	0	0.14	1,055	1,187	2,242	NA	NA	NA
1986	38	1	0.22	642	618	1,260	NA	NA	NA
1987	27	0	---	557	643	1,200	NA	NA	NA
1988	26	0	0.17	537	464	1,001	NA	NA	NA
1989	21	0	0.10	757	796	1,553	NA	NA	NA
1990	25	16	0.17	988	1,117	2,105	NA	NA	NA
1991	16	2	0.35	478	628	1,106	NA	NA	NA
1992	17	8	0.04	784	765	1,549	NA	NA	NA
1993	22	2	0.05	583	624	1,207	NA	NA	NA
1994	42	12	0.05	554	633	1,187	NA	NA	NA
1995	22	3	0.05	599	575	1,174	NA	NA	NA
1996	30	8	0.04	724	775	1,499	NA	NA	NA
1997	16	14	0.04	682	853	1,535	NA	NA	NA
1998	22	9	0.04	863	784	1,647	NA	NA	NA
2000	31	0	0.05	430	370	800	NA	NA	NA
2001	15	9	0.05	314	378	692	NA	NA	NA
2002	18	1	0.07	278	326	604	NA	NA	NA
2003	17	2	0.05	294	322	616	NA	NA	NA
2004	13	2	0.09	422	315	737	NA	NA	NA
2005	22	1	0.04	543	335	878	NA	NA	NA
2006	17	2	0.04	295	487	782	NA	NA	NA
2007	9	1	0.06	NA	NA	NA	NA	NA	NA
2008	10	2	0.06	172	248	420	NA	NA	NA
2008	9	3	0.06	254	301	555	NA	NA	NA

Table 1.12. Estimates of pollock biomass obtained from GLM model predictions of pollock CPUE and INPFC area expansions. Biomass estimates were multiplied by the von Szalay and Brown (2001) FPC of 3.84 for comparison to the NMFS triennial trawl survey biomass estimates. Coefficients of variation do not reflect the variance of the FPC estimate.

<i>Year</i>	<i>Biomass (t)</i>	<i>FPC-adjusted</i>	<i>biomass (t)</i>	<i>CV</i>
1961	50,356		193,369	0.24
1962	57,496		220,783	0.30
1970	7,979		30,640	0.42
1971	4,257		16,348	0.64
1974	1,123,447		4,314,035	0.38
1975	1,501,142		5,764,384	0.52
1978	223,277		857,383	0.31
1980	146,559		562,787	0.27
1981	257,219		987,719	0.33
1982	356,433		1,368,703	0.29

Other published estimates of pollock biomass from surveys using 400-mesh eastern trawls

<i>Year</i>	<i>Biomass (t)</i>	<i>Source</i>
1961	57,449	<i>Ronholt et al. 1978</i>
1961-62	91,075	<i>Ronholt et al. 1978</i>
1973-75	1,055,000	<i>Alton et al. 1977</i>
1973-76	739,293	<i>Ronholt et al. 1978</i>
1973-75	610,413	<i>Hughes and Hirschhorn 1979</i>

Table 1.13. Predictions of Gulf of Alaska pollock year-class strength. The FOCI prediction is the prediction of year-class strength made in the natal year of the year class, and was derived from environmental indices, larval surveys, and the time series characteristics of pollock recruitment. The McKelvey index is the estimated abundance of 9-16 cm pollock from the Shelikof Strait EIT survey.

<i>Year class</i>	<i>FOCI prediction</i>	<i>Year of EIT survey</i>	<i>McKelvey index</i>	<i>Rank abundance of McKelvey index</i>
1980		1981	0.078	14
1981				
1982		1983	0.001	26
1983		1984	0.062	17
1984		1985	2.092	3
1985		1986	0.579	6
1986				
1987		1988	0.017	24
1988		1989	0.399	7
1989		1990	0.049	22
1990		1991	0.022	23
1991		1992	0.228	11
1992	Strong	1993	0.063	16
1993	Average	1994	0.186	12
1994	Average	1995	10.688	1
1995	Average-Strong	1996	0.061	18
1996	Average	1997	0.070	15
1997	Average	1998	0.395	8
1998	Average			
1999	Average	2000	4.484	2
2000	Average	2001	0.291	10
2001	Average-Strong	2002	0.008	25
2002	Average	2003	0.051	21
2003	Average	2004	0.053	20
2004	Average	2005	1.626	4
2005	Average	2006	0.162	13
2006	Average	2007	0.054	19
2007	Average	2008	1.368	5
2008	Average	2009	0.332	9
2009	Average		---	---

Table 1.14. Ageing error transition matrix used in the Gulf of Alaska pollock assessment model.

<i>True Age</i>	<i>St. dev.</i>	<i>Observed Age</i>									
		<i>1</i>	<i>2</i>	<i>3</i>	<i>4</i>	<i>5</i>	<i>6</i>	<i>7</i>	<i>8</i>	<i>9</i>	<i>10</i>
1	0.18	0.9970	0.0030	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
2	0.23	0.0138	0.9724	0.0138	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
3	0.27	0.0000	0.0329	0.9342	0.0329	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
4	0.32	0.0000	0.0000	0.0571	0.8858	0.0571	0.0000	0.0000	0.0000	0.0000	0.0000
5	0.36	0.0000	0.0000	0.0000	0.0832	0.8335	0.0832	0.0000	0.0000	0.0000	0.0000
6	0.41	0.0000	0.0000	0.0000	0.0001	0.1090	0.7817	0.1090	0.0001	0.0000	0.0000
7	0.45	0.0000	0.0000	0.0000	0.0000	0.0004	0.1333	0.7325	0.1333	0.0004	0.0000
8	0.50	0.0000	0.0000	0.0000	0.0000	0.0000	0.0012	0.1554	0.6868	0.1554	0.0012
9	0.54	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0028	0.1747	0.6450	0.1775
10	0.59	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0052	0.1913	0.8035

Table 1.15. Proportion mature at age for female pollock based on maturity stage data collected during winter EIT surveys in the Gulf of Alaska (1983-2009).

Year	2	3	4	5	6	7	8	9	10+	Sample size
1983	0.000	0.165	0.798	0.960	0.974	0.983	0.943	1.000	1.000	1333
1984	0.000	0.145	0.688	0.959	0.990	1.000	0.992	1.000	1.000	1621
1985	0.015	0.051	0.424	0.520	0.929	0.992	0.992	1.000	1.000	1183
1986	0.000	0.021	0.105	0.849	0.902	0.959	1.000	1.000	1.000	618
1987	0.000	0.012	0.106	0.340	0.769	0.885	0.950	0.991	1.000	638
1988	0.000	0.000	0.209	0.176	0.606	0.667	1.000	0.857	0.964	464
1989	0.000	0.000	0.297	0.442	0.710	0.919	1.000	1.000	1.000	796
1990	0.000	0.000	0.192	0.674	0.755	0.910	0.945	0.967	0.996	1844
1991	0.000	0.000	0.111	0.082	0.567	0.802	0.864	0.978	1.000	628
1992	0.000	0.000	0.040	0.069	0.774	0.981	0.990	1.000	0.983	765
1993	0.000	0.016	0.120	0.465	0.429	0.804	0.968	1.000	0.985	624
1994	0.000	0.007	0.422	0.931	0.941	0.891	0.974	1.000	1.000	872
1995	0.000	0.000	0.153	0.716	0.967	0.978	0.921	0.917	0.977	805
1996	0.000	0.000	0.036	0.717	0.918	0.975	0.963	1.000	0.957	763
1997	0.000	0.000	0.241	0.760	1.000	1.000	0.996	1.000	1.000	843
1998	0.000	0.000	0.065	0.203	0.833	0.964	1.000	1.000	0.989	757
2000	0.000	0.012	0.125	0.632	0.780	0.579	0.846	1.000	0.923	356
2001	0.000	0.000	0.289	0.308	0.825	0.945	0.967	0.929	1.000	374
2002	0.000	0.026	0.259	0.750	0.933	0.974	1.000	1.000	1.000	499
2003	0.000	0.029	0.192	0.387	0.529	0.909	0.750	1.000	1.000	301
2004	0.000	0.000	0.558	0.680	0.745	0.667	1.000	1.000	1.000	444
2005	0.000	0.000	0.706	0.882	0.873	0.941	1.000	1.000	1.000	321
2006	0.000	0.000	0.043	0.483	0.947	0.951	0.986	1.000	1.000	476
2007	0.000	0.000	0.333	0.667	0.951	0.986	0.983	1.000	1.000	313
2008	0.000	0.000	0.102	0.241	0.833	1.000	0.968	0.952	1.000	240
2009	0.000	0.000	0.140	0.400	0.696	1.000	1.000	1.000	1.000	296
<i>Average</i>										
<i>All years</i>	0.001	0.019	0.260	0.550	0.814	0.910	0.961	0.984	0.991	
<i>2000-2009</i>	0.000	0.007	0.275	0.543	0.811	0.895	0.950	0.988	0.992	
<i>2005-2009</i>	0.000	0.000	0.265	0.535	0.860	0.976	0.987	0.990	1.000	

Table 1.16. Estimated selectivity at age for Gulf of Alaska pollock fisheries and surveys. The fisheries and surveys were modeled using double logistic selectivity functions with random walk process error for the fishery logistic parameters. Fishery selectivity at age reported below is the average of the annual selectivity for the indicated time period, rescaled so that the maximum is one.

Age	Historical					400-mesh		
	POP fishery (1961-71)	Foreign (1972-84)	domestic (1985-2001)	Recent domestic (2002-2008)	EIT survey	Bottom trawl survey	ADF&G bottom trawl	eastern trawl 1961-82
2	0.001	0.040	0.039	0.224	0.986	0.199	0.060	0.120
3	0.021	0.256	0.142	0.435	0.972	0.331	0.136	0.395
4	0.415	0.749	0.392	0.679	0.943	0.516	0.282	0.758
5	1.000	1.000	0.718	0.864	0.889	0.733	0.496	0.938
6	0.949	0.928	0.932	0.962	0.793	0.921	0.713	0.986
7	0.706	0.686	1.000	1.000	0.649	1.000	0.866	0.997
8	0.364	0.352	0.944	0.978	0.472	0.935	0.947	0.999
9	0.131	0.130	0.742	0.687	0.301	0.766	0.984	1.000
10	0.039	0.042	0.328	0.165	0.172	0.566	1.000	1.000

Table 1.17. Total estimated abundance at age (numbers in 000,000s) of Gulf of Alaska pollock from the age-structured assessment model.

	<i>Age</i>								
	2	3	4	5	6	7	8	9	10
1961	361	186	114	71	53	37	27	20	16
1962	398	267	138	85	52	39	27	20	27
1963	426	295	198	102	63	39	29	20	34
1964	96	316	219	147	76	47	29	21	41
1965	247	71	234	162	108	56	34	21	46
1966	131	183	53	172	118	79	41	25	50
1967	326	97	135	38	122	84	57	30	55
1968	382	241	72	99	27	87	60	41	63
1969	663	283	178	52	70	19	63	44	77
1970	310	491	209	124	33	45	13	44	89
1971	660	230	363	151	86	23	32	9	98
1972	1,222	489	170	264	106	61	17	23	79
1973	933	905	362	121	176	71	42	12	75
1974	3,047	691	669	257	80	118	49	30	64
1975	625	2,257	511	471	164	52	79	35	69
1976	397	463	1,649	356	324	114	37	57	77
1977	1,872	293	333	1,136	244	225	80	26	99
1978	2,629	1,384	213	228	766	166	157	57	92
1979	2,467	1,940	992	145	155	525	117	113	110
1980	3,492	1,823	1,403	680	99	106	369	84	164
1981	1,778	2,578	1,322	980	468	68	74	263	182
1982	433	1,314	1,875	920	669	321	47	53	326
1983	488	317	936	1,299	634	463	226	34	280
1984	203	358	225	631	863	423	318	162	232
1985	463	148	246	141	377	516	266	221	290
1986	1,574	337	102	150	77	201	284	169	371
1987	538	1,149	237	67	94	48	127	196	398
1988	157	395	828	163	44	61	31	84	433
1989	370	115	287	580	109	29	39	20	378
1990	1,604	273	84	204	393	70	18	25	290
1991	1,000	1,185	201	61	140	250	43	11	230
1992	399	739	870	144	42	92	160	27	158
1993	237	294	537	607	95	27	58	102	132
1994	142	174	213	375	402	61	17	37	160
1995	214	105	127	150	252	264	40	11	135
1996	835	158	77	90	103	171	178	27	104
1997	396	617	116	55	64	71	117	121	91
1998	169	292	451	82	37	40	44	72	136
1999	155	123	204	290	49	21	23	25	127
2000	215	114	88	137	177	28	12	13	95
2001	885	159	83	62	88	106	16	7	72
2002	827	647	113	56	39	53	64	10	54
2003	125	602	460	77	36	25	34	40	44
2004	99	91	427	315	51	24	16	22	59
2005	89	70	62	284	206	33	16	11	59
2006	305	63	48	41	180	129	21	10	50
2007	715	219	44	32	26	113	80	13	43
2008	308	514	153	30	21	17	74	53	40
2009	1,156	222	363	106	20	14	11	50	65
<i>Average</i>	746	536	388	265	177	117	78	54	132

Table 1.18. Estimates of population biomass, recruitment, and harvest of Gulf of Alaska pollock from the age-structured assessment model. The harvest rate is the catch in biomass divided by the total biomass of age 3+ fish at the start of the year.

Year	3+ total			Age 2			2008 Assessment results		
	biomass (1,000 t)	Female spawn. biom. (1,000 t)	Age 2 recruits (million)	Catch (t)	Harvest rate	3+ total biomass	Female spawn. biom.	Age 2 recruits	Harvest rate
1977	1,909	445	1,872	118,356	6%	2,145	502	2,043	6%
1978	2,067	480	2,629	96,935	5%	2,319	545	2,820	4%
1979	2,559	488	2,467	105,748	4%	2,840	555	2,591	4%
1980	3,040	542	3,492	114,622	4%	3,333	613	3,636	3%
1981	3,719	450	1,778	147,744	4%	4,020	502	1,840	4%
1982	3,872	527	433	168,740	4%	4,157	580	446	4%
1983	3,287	654	488	215,608	7%	3,515	709	504	6%
1984	2,658	688	203	307,401	12%	2,846	745	210	11%
1985	1,960	619	463	284,826	15%	2,114	675	482	13%
1986	1,568	502	1,574	87,809	6%	1,709	551	1,629	5%
1987	1,636	421	538	69,751	4%	1,763	463	553	4%
1988	1,549	382	157	65,739	4%	1,659	419	160	4%
1989	1,405	372	370	78,392	6%	1,502	404	375	5%
1990	1,199	336	1,604	90,744	8%	1,277	362	1,617	7%
1991	1,323	318	1,000	100,488	8%	1,388	343	1,007	7%
1992	1,641	282	399	90,857	6%	1,696	302	400	5%
1993	1,491	319	237	108,908	7%	1,535	335	237	7%
1994	1,250	369	142	107,335	9%	1,284	383	142	8%
1995	1,050	338	214	72,618	7%	1,078	349	215	7%
1996	870	305	835	51,263	6%	891	314	830	6%
1997	887	263	396	90,130	10%	902	270	394	10%
1998	811	199	169	125,098	15%	820	204	167	15%
1999	653	181	155	95,590	15%	659	185	151	15%
2000	573	169	215	73,080	13%	577	172	206	13%
2001	543	165	885	72,076	13%	542	167	821	13%
2002	692	138	827	51,937	8%	668	139	745	8%
2003	857	131	125	50,666	6%	804	130	112	6%
2004	760	148	99	63,934	8%	706	141	93	9%
2005	643	192	89	80,846	13%	589	178	83	14%
2006	557	205	305	71,976	13%	503	185	307	14%
2007	535	186	715	53,062	10%	481	164	618	11%
2008	603	185	308	52,500	9%	537	161	600	10%
2009	652	182	1,156						
Average									
1977-2009	1,479	339	798	105,149	8%	1,589	367	814	8%
1979-2008			689					706	

Table 1.19. Gulf of Alaska pollock life history and fishery vectors used to estimate spawning biomass per recruit (F_{SPR}) harvest rates. Population weight at age is based on a average for the bottom trawl survey conducted in June to August. Spawning weight at age is based on an average from the Shelikof Strait EIT survey conducted March. Proportion mature females is the average from winter EIT survey specimen data for 1983-2009.

	<i>Natural mortality</i>	<i>Fishery selectivity (Avg. 2004-2008)</i>	<i>Weight at age (kg)</i>			<i>Proportion mature females</i>
			<i>(Avg. 2005-2009)</i>	<i>Population (Avg. 2003-2007)</i>	<i>Fishery (Avg. 2004-2008)</i>	
2	0.3	0.252	0.078	0.172	0.290	0.001
3	0.3	0.476	0.262	0.386	0.528	0.019
4	0.3	0.715	0.482	0.601	0.845	0.260
5	0.3	0.883	0.791	0.859	1.120	0.550
6	0.3	0.968	1.109	1.018	1.227	0.814
7	0.3	1.000	1.377	1.191	1.346	0.910
8	0.3	0.964	1.567	1.301	1.491	0.961
9	0.3	0.610	1.697	1.484	1.550	0.984
10+	0.3	0.124	1.907	1.608	1.806	0.991

Table 1.20. Methods used to assess Gulf of Alaska pollock, 1977-2008. The basis for catch recommendation in 1977-1989 is the presumptive method by which the TAC was determined (based on the assessment and SSC minutes). The basis for catch recommendation given in 1990-2008 is the method used by the Plan Team to derive the ABC recommendation given in the SAFE summary chapter.

<i>Year</i>	<i>Assessment method</i>	<i>Basis for catch recommendation in following year</i>	<i>B40% (t)</i>
1977-81	Survey biomass, CPUE trends, M=0.4	MSY = 0.4 * M * Bzero	---
1982	CAGEAN	MSY = 0.4 * M * Bzero	---
1983	CAGEAN	Mean annual surplus production	---
1984	Projection of survey numbers at age	Stabilize biomass trend	---
1985	CAGEAN, projection of survey numbers at age, CPUE trends	Stabilize biomass trend	---
1986	CAGEAN, projection of survey numbers at age	Stabilize biomass trend	---
1987	CAGEAN, projection of survey numbers at age	Stabilize biomass trend	---
1988	CAGEAN, projection of survey numbers at age	10% of exploitable biomass	---
1989	Stock synthesis	10% of exploitable biomass	---
1990	Stock synthesis, reduce <i>M</i> to 0.3	10% of exploitable biomass	---
1991	Stock synthesis, assume trawl survey catchability = 1	FMSY from an assumed SR curve	---
1992	Stock synthesis	Max[-Pr(SB<Threshold)+Yld]	---
1993	Stock synthesis	Pr(SB>B20)=0.95	---
1994	Stock synthesis	Pr(SB>B20)=0.95	---
1995	Stock synthesis	Max[-Pr(SB<Threshold)+Yld]	---
1996	Stock synthesis	Amendment 44 Tier 3 guidelines	289,689
1997	Stock synthesis	Amendment 44 Tier 3 guidelines	267,600
1998	Stock synthesis	Amendment 44 Tier 3 guidelines	240,000
1999	AD model builder	Amendment 56 Tier 3 guidelines (with a reduction from max permissible F _{ABC})	247,000
2000	AD model builder	Amendment 56 Tier 3 guidelines	250,000
2001	AD model builder	Amendment 56 Tier 3 guidelines (with a reduction from max permissible F _{ABC})	245,000
2002	AD model builder	Amendment 56 Tier 3 guidelines (with a reduction from max permissible F _{ABC})	240,000
2003	AD model builder	Amendment 56 Tier 3 guidelines (with a reduction from max permissible F _{ABC})	248,000
2004	AD model builder	Amendment 56 Tier 3 guidelines (with a reduction from max permissible F _{ABC} , and stairstep approach for projected ABC increase)	229,000
2005	AD model builder	Amendment 56 Tier 3 guidelines (with a reduction from max permissible F _{ABC})	224,000
2006	AD model builder	Amendment 56 Tier 3 guidelines (with a reduction from max permissible F _{ABC})	220,000
2007	AD model builder	Amendment 56 Tier 3 guidelines (with a reduction from max permissible F _{ABC})	221,000
2008	AD model builder	Amendment 56 Tier 3 guidelines (with a reduction from max permissible F _{ABC})	237,000

Table 1.21. Projections of Gulf of Alaska pollock spawning biomass, full recruitment fishing mortality, and catch for 2009-2022 under different harvest policies. All projections begin with estimated age composition in 2009 using base run model, and a projected 2009 catch of 43,865 t. The values for $B_{100\%}$, $B_{40\%}$, and $B_{35\%}$ are 620,000, 248,000, and 217,000 t, respectively.

<i>Spawning biomass (t)</i>	<i>Max F_{ABC}</i>	<i>Author's recommended F</i>	<i>Average F</i>	<i>$F_{75\%}$</i>	<i>$F = 0$</i>	<i>F_{OFL}</i>	<i>Max F_{ABC} for two years, then F_{OFL}</i>
2009	163,748	163,748	163,748	163,748	163,748	163,748	163,748
2010	184,011	184,567	184,869	186,264	187,817	183,415	184,011
2011	213,891	218,019	221,180	231,987	244,669	209,571	213,891
2012	242,608	251,387	263,728	288,464	318,933	233,466	241,264
2013	262,817	275,293	304,476	346,671	401,070	248,445	254,619
2014	266,732	279,939	327,125	385,307	463,504	248,320	252,631
2015	267,655	280,375	341,767	412,927	511,907	246,713	249,404
2016	266,679	278,442	349,569	430,004	544,838	244,397	245,903
2017	266,393	277,485	355,354	443,143	570,864	243,287	244,119
2018	265,398	275,866	357,375	449,302	584,540	242,034	242,500
2019	266,123	276,164	359,974	454,793	595,379	242,595	242,858
2020	268,336	278,138	363,900	461,154	606,067	244,558	244,701
2021	268,947	278,573	366,003	465,185	613,466	244,890	244,967
2022	267,979	277,451	365,897	466,387	617,079	243,764	243,805

<i>Fishing mortality</i>	<i>Max F_{ABC}</i>	<i>Author's recommended F</i>	<i>Average F</i>	<i>$F_{75\%}$</i>	<i>$F = 0$</i>	<i>F_{OFL}</i>	<i>Max F_{ABC} for two years, then F_{OFL}</i>
2009	0.09	0.09	0.09	0.09	0	0.09	0.09
2010	0.17	0.14	0.13	0.07	0	0.19	0.17
2011	0.20	0.17	0.13	0.07	0	0.22	0.20
2012	0.22	0.20	0.13	0.07	0	0.25	0.25
2013	0.22	0.21	0.13	0.07	0	0.25	0.25
2014	0.21	0.20	0.13	0.07	0	0.24	0.24
2015	0.21	0.20	0.13	0.07	0	0.23	0.23
2016	0.20	0.19	0.13	0.07	0	0.23	0.23
2017	0.20	0.19	0.13	0.07	0	0.23	0.23
2018	0.20	0.19	0.13	0.07	0	0.23	0.23
2019	0.20	0.19	0.13	0.07	0	0.23	0.23
2020	0.20	0.19	0.13	0.07	0	0.23	0.23
2021	0.20	0.19	0.13	0.07	0	0.23	0.23
2022	0.20	0.19	0.13	0.07	0	0.23	0.23

<i>Catch (t)</i>	<i>Max F_{ABC}</i>	<i>Author's recommended F</i>	<i>Average F</i>	<i>$F_{75\%}$</i>	<i>$F = 0$</i>	<i>F_{OFL}</i>	<i>Max F_{ABC} for two years, then F_{OFL}</i>
2009	43,865	43,865	43,865	43,865	43,865	43,865	43,865
2010	102,541	88,081	80,136	42,848	0	117,871	102,541
2011	144,202	128,123	99,033	54,918	0	160,249	144,202
2012	177,016	166,277	113,757	65,186	0	192,447	202,604
2013	179,996	177,102	119,702	70,403	0	194,007	199,508
2014	172,296	170,485	121,109	72,439	0	183,818	186,654
2015	167,236	165,273	121,380	73,441	0	177,652	178,921
2016	162,089	159,248	119,357	72,527	0	171,767	172,235
2017	159,216	155,723	116,711	70,772	0	169,000	169,070
2018	160,194	156,612	117,237	71,089	0	170,461	170,406
2019	161,192	157,446	117,712	71,389	0	171,626	171,576
2020	161,417	157,622	117,493	71,286	0	171,885	171,846
2021	159,655	155,947	116,714	70,908	0	169,981	169,953
2022	158,078	154,460	115,840	70,486	0	168,195	168,178

Table 1.22. Projections of Gulf of Alaska pollock spawning biomass, full recruitment fishing mortality, and catch for 2009-2022 under different harvest policies. All projections begin with estimated age composition in 2009 using base run model, except at recruitment in 2009 is set to mean 1979-2008 recruitment, and a projected 2009 catch of 43,865 t. The values for $B_{100\%}$, $B_{40\%}$, and $B_{35\%}$ are 620,000, 248,000, and 217,000 t, respectively.

<i>Spawning biomass (t)</i>	<i>Max F_{ABC}</i>	<i>Author's recommended F</i>	<i>Average F</i>	<i>$F_{75\%}$</i>	<i>$F = 0$</i>	<i>F_{OFL}</i>	<i>Max F_{ABC} for two years, then F_{OFL}</i>
2009	163,630	163,630	163,630	163,630	163,630	163,630	163,630
2010	182,644	183,192	183,466	184,850	186,391	182,057	182,644
2011	200,569	204,417	206,912	217,148	229,173	196,546	200,569
2012	215,668	223,276	231,396	253,272	280,295	207,885	214,615
2013	231,701	242,518	260,509	296,112	342,099	220,184	225,331
2014	244,117	256,352	287,324	336,730	403,003	229,179	232,684
2015	254,276	266,689	310,601	372,515	458,103	236,526	238,654
2016	259,778	271,680	326,738	398,751	500,668	240,002	241,135
2017	262,108	273,437	337,436	417,843	533,864	240,865	241,447
2018	262,998	273,770	344,312	430,716	557,130	240,855	241,162
2019	264,778	275,091	350,449	441,139	575,073	242,015	242,180
2020	267,591	277,608	356,956	451,122	591,024	244,279	244,365
2021	268,527	278,320	360,940	457,815	602,322	244,757	244,802
2022	267,738	277,335	362,205	460,973	608,823	243,699	243,722

<i>Fishing mortality</i>	<i>Max F_{ABC}</i>	<i>Author's recommended F</i>	<i>Average F</i>	<i>$F_{75\%}$</i>	<i>$F = 0$</i>	<i>F_{OFL}</i>	<i>Max F_{ABC} for two years, then F_{OFL}</i>
2009	0.09	0.09	0.09	0.09	0	0.09	0.09
2010	0.17	0.14	0.13	0.07	0	0.19	0.17
2011	0.18	0.16	0.13	0.07	0	0.21	0.18
2012	0.20	0.17	0.13	0.07	0	0.22	0.23
2013	0.20	0.18	0.13	0.07	0	0.22	0.23
2014	0.20	0.19	0.13	0.07	0	0.22	0.23
2015	0.20	0.19	0.13	0.07	0	0.22	0.23
2016	0.20	0.19	0.13	0.07	0	0.22	0.22
2017	0.20	0.19	0.13	0.07	0	0.22	0.23
2018	0.20	0.19	0.13	0.07	0	0.23	0.23
2019	0.20	0.19	0.13	0.07	0	0.23	0.23
2020	0.20	0.19	0.13	0.07	0	0.23	0.23
2021	0.20	0.19	0.13	0.07	0	0.23	0.23
2022	0.20	0.19	0.13	0.07	0	0.23	0.23

<i>Catch (t)</i>	<i>Max F_{ABC}</i>	<i>Author's recommended F</i>	<i>Average F</i>	<i>$F_{75\%}$</i>	<i>$F = 0$</i>	<i>F_{OFL}</i>	<i>Max F_{ABC} for two years, then F_{OFL}</i>
2009	43,865	43,865	43,865	43,865	43,865	43,865	43,865
2010	89,804	77,151	70,749	37,858	0	103,206	89,804
2011	114,358	101,512	83,699	46,468	0	127,122	114,358
2012	139,412	128,790	97,031	55,511	0	151,375	159,706
2013	153,448	148,022	106,496	62,321	0	165,370	170,216
2014	156,829	153,677	111,365	66,106	0	168,294	170,639
2015	159,387	156,772	114,602	68,763	0	170,535	171,477
2016	159,844	157,014	116,373	70,364	0	170,468	170,732
2017	159,478	156,105	116,210	70,401	0	169,876	169,849
2018	160,563	157,128	116,872	70,816	0	171,181	171,086
2019	161,467	157,812	117,446	71,189	0	172,069	171,999
2020	161,629	157,865	117,299	71,139	0	172,141	172,098
2021	159,775	156,070	116,573	70,800	0	170,123	170,097
2022	158,140	154,579	115,737	70,407	0	168,266	168,252




Figure 1.1 Pollock catch in 2008 by 20 X 20 km blocks by season in the Gulf of Alaska as determined by observer-recorded haul retrieval locations. Blocks with less than 1.0 t of pollock catch are not shown. The size of the circle is proportional to the catch.




Figure 1.2. 2008 catch age composition by half year and statistical area.




Figure 1.3. Gulf of Alaska pollock catch age composition (1976-2008). The diameter of the circle is proportional to the catch. Diagonal lines show strong year classes (1972, 1975, 1976, 1977, 1978, 1979, 1984, 1988, 1994, 1995, 1999, 2000, 2004, and 2005).




Figure 1.4. Pollock CPUE for the 2009 NMFS bottom trawl survey.




Figure 1.5. Size composition of pollock by statistical area for the 2009 NMFS bottom trawl survey.





Figure 1.6. Biomass estimates of juvenile pollock (top) and adult pollock (bottom) from 1986-2008 Shelikof Strait EIT surveys. Bottom panel also shows the model estimate of total spawning biomass.




Figure 1.7. Trends in biomass estimates of from winter acoustic surveys of pre-spawning aggregations of pollock in the Gulf of Alaska.




Figure 1.8. Biomass by length for pollock in the Shelikof Strait EIT survey (1981-2009, except 1982,1987 and 1999).




Figure 1.9. Length frequency of pollock in the ADF&G crab/groundfish trawl survey (1989-2009, except 1991 and 1995).




Figure 1.10. Relative trends in pollock biomass since 1987 for the Shelikof Strait EIT survey, the NMFS bottom trawl survey, and the ADF&G crab/groundfish trawl survey. Each survey biomass estimate is standardized to the average since 1987. Shelikof Strait EIT surveys prior to 2008 were re-scaled to be comparable to the surveys conducted from 2008 onwards by the *R/V Oscar Dyson*.




Figure 1.11. Gulf of Alaska pollock catch characteristics.




Figure 1.12. Estimates of the proportion mature at age from visual maturity data collected during 2005-2009 winter EIT surveys in the Gulf of Alaska and long-term average proportion mature at age (1983-2009).




Figure 1.13. Age at 50% mature (top) and length at 50% mature (bottom) from annual logistic regressions for female pollock from winter EIT survey data in the Gulf of Alaska, 1983-2009.




Figure 1.14. Observed and predicted fishery age composition for Gulf of Alaska pollock from the base model. Continuous lines are model predictions and lines with + symbol are observed proportions at age.




Figure 1.15. Observed and predicted Shelikof Strait EIT survey age composition for Gulf of Alaska pollock from the base model. Continuous lines are model predictions and lines with + symbol are observed proportions at age.




Figure 1.16. Observed and predicted NMFS bottom trawl age composition for Gulf of Alaska pollock from the base model. Continuous lines are model predictions and lines with + symbol are observed proportions at age.





Figure 1.17. Model predicted and observed survey biomass for the Shelikof Strait EIT survey. The Shelikof EIT survey is modeled with three catchability periods corresponding to the two acoustic systems used on the *R/V Miller Freeman* (MF), with an additional catchability period for the *R/V Dyson* (DY) in 2008 and 2009. Error bars indicate plus and minus two standard deviations.




Figure 1.18. Model predicted and observed survey biomass for the NMFS bottom trawl survey (top), and the ADFG crab/groundfish survey (bottom). Error bars indicate plus and minus two standard deviations. Since variance estimates are unavailable for ADFG biomass estimates, an assumed CV of 0.25 is used in the assessment model.





Figure 1.19. Model predicted and observed survey biomass for the historical 400-mesh eastern trawl surveys (top), and the egg production survey (bottom). Error bars indicate plus and minus two standard deviations.




Figure 1.20. Uncertainty in the catchability coefficient for the NMFS trawl survey from a likelihood profile for the base model.




Figure 1.21. Estimates of time-varying fishery selectivity for Gulf of Alaska pollock. The maximum selectivity in each year is 1.0.





Figure 1.22. Estimated time series of Gulf of Alaska pollock spawning biomass (million t, top) and age-2 recruitment (billions of fish, bottom) from 1961 to 2009. Vertical bars represent two standard deviations. The B35% and B40% lines represent the current estimate of these benchmarks.





Figure 1.23. Retrospective plot of estimated Gulf of Alaska pollock female spawning biomass for stock assessments in the years 1997-2009 (top). For this figure, the time series of female spawning biomass for the 2007 assessment was calculated using the weight and maturity at age used in previous assessments to facilitate comparison. The bottom panel shows the estimated age composition in 2009 from the 2008 and 2009 assessments.




Figure 1.24. Gulf of Alaska pollock spawner productivity $\log(R/S)$ in 1961-2008 (top). A five-year running average is also shown. Spawner productivity in relation to female spawning biomass (bottom). The Ricker stock-recruit curve is linear in a plot of spawner productivity against spawning biomass. Horizontal lines indicate the mean spawner productivity for each decade within the range of spawning biomass indicated by the endpoints of the lines.




Figure 1.25. Gulf of Alaska pollock spawning biomass relative to the unfished level and fishing mortality relative to F_{MSY} (1961-2009). The ratio of fishing mortality to F_{MSY} is calculated using the estimated selectivity pattern in that year. Estimates of $B_{100\%}$ spawning biomass are based on current estimates of maturity at age, weight at age, and mean recruitment. Because these estimates change as new data become available, this figure can only be used in a general way to evaluate management performance relative to biomass and fishing mortality reference levels.




Figure 1.26. Uncertainty in spawning biomass in 2010-2014 based on a thinned MCMC chain from the joint marginal likelihood for the base model where catch is set to the author's recommended FABC.





Figure 1.27. Projected spawning biomass and catches in 2009-14 under different management strategies.





Figure 1.28. Variability in projected catch and spawning biomass in 2009-2022 under the author's recommended FABC.




Figure 1.29. Gulf of Alaska food web showing demersal (red) and pelagic (blue) pathways. Walleye pollock is shown in green. Pollock consumers stain green according to the importance of pollock in their diet.





Figure 1.30. Diet (percent wet weight) of GOA walleye pollock juveniles (top) and adults (bottom) from summer food habits data collected on NMFS bottom trawl surveys, 1990-2005.





Figure 1.31. Sources of mortality for walleye pollock juveniles (top) and adults (bottom) from an ECOPATH model of the Gulf of Alaska. Pollock less than 20cm are considered juveniles.




Figure 1.32. Diet diversity of major predators of walleye pollock from an ECOPEATH model for Gulf of Alaska during 1990-94.




Figure 1.33. Length frequencies and percent by weight of each length class of pollock prey (cm fork length) in stomachs of four major groundfish predators, from AFSC bottom-trawl surveys 1987-2005. Length of prey is uncorrected for digestion state.





Figure 1.34. (Top) Historical trends in GOA walleye pollock, Pacific cod, Pacific halibut, arrowtooth flounder, and Steller Sea Lions, from stock assessment data. (Bottom) Total catch and consumption of walleye pollock in survey years (bars) and production + biomass change as calculated from the current stock assessment results (line). See text for calculation methods.






Figure 1.35. (Top) Consumption per unit predator survey biomass of GOA walleye pollock <30cm fork length in diets, shown for each survey year. (Middle and bottom) Normalized consumption/biomass and normalized total consumption of pollock <30cm fork length, plotted against age 2 pollock numbers reported in Table 1.16.






Figure 1.36. (Top) Consumption per unit predator survey biomass of GOA walleye pollock $\geq 30\text{cm}$ fork length in diets, shown for each survey year. (Middle and bottom) Normalized consumption/biomass and normalized total consumption of pollock $\geq 30\text{cm}$ fork length, plotted against age 3+ pollock biomass reported in Table 1.17.




Figure 1.37. Ecosystem model output (percent change at future equilibrium of indicated groups) resulting from reducing adult pollock survival by 10% (top graph), reducing juvenile pollock survival by 10% (middle graph), and reducing pollock trawl effort by 10%. Dark bars indicate biomass changes of modeled species, while light bars indicate changes in fisheries catch (landings+discards) assuming a constant fishing rate within the indicated fishery. Graphs show 50% and 95% confidence intervals (bars and lines respectively) summarized over 20,000 ecosystems drawn from error ranges of input parameters (see Aydin et al. 2005 for methodology). Only the top 20 effects, sorted by median, are shown for each perturbation.




Figure 1.38. Ecosystem model output, shown as percent change at future equilibrium of adult pollock (top) and juvenile pollock, resulting from independently lowering the indicated species' survival rates by 10% (dark bars) or by reducing fishing effort of a particular gear by 10% (light bars). Graphs show 50% and 95% confidence intervals (bars and lines respectively) summarized over 20,000 ecosystems drawn from error ranges of input parameters (see Aydin et al. 2005 for methodology). Only the top 20 effects, sorted by median, are shown for each perturbation.




Figure 1.39. Ecosystem model output, shown as percent change at future equilibrium of four major predators on walleye pollock, resulting from independently lowering the indicated species' survival rates by 10% (dark bars) or by reducing fishing effort of a particular gear by 10% (light bars). Graphs show 50% and 95% confidence intervals (bars and lines respectively) summarized over 20,000 ecosystems drawn from error ranges of input parameters (see Aydin et al. 2005 for methodology). Only the top 20 effects, sorted by median, are shown for each perturbation.





Figure 1.40. Pair-wise Spearman rank correlation between abundance trends of walleye pollock, pollock fishery catches, Steller sea lions, arrowtooth flounder, Pacific halibut, and Pacific cod in the Gulf of Alaska. Rank correlations are based on the years in which abundance estimates are available for each pair.

Appendix A: Southeast Alaska pollock

Bottom trawl surveys indicate a substantial reduction in pollock abundance east of 140° W. lon. Stock structure in this area is poorly understood. Bailey et al. (1999) suggest that pollock metapopulation structure in southeast Alaska is characterized by numerous fiord populations. In the 2009 bottom trawl survey, higher pollock CPUE in southeast Alaska occurred primarily from Cape Ommaney to Dixon Entrance, where the shelf is more extensive. Pollock length composition in the 2009 bottom trawl survey is dominated by mode smaller fish (~22 cm) which likely represent age-1 pollock (Appendix Fig. 1.1). There is also broad range of larger fish that likely represent several year classes. Juveniles in this area are unlikely to influence the population dynamics of pollock in the central and western Gulf of Alaska. Ocean currents are generally northward in this area, suggesting that juvenile settlement is a result of spawning further south. Spawning aggregations of pollock have been reported from the northern part of Dixon Entrance (Saunders et al. 1988).

Historically, there has been little directed fishing for pollock in southeast Alaska (Fritz 1993). Pollock catch the Southeast and East Yakutat statistical areas has averaged about 1 t since 2000 (Table 1.4). The ban on trawling east of 140° W. lon. prevents the development of a trawl fishery for pollock in Southeast Alaska.

Pollock biomass estimates from the bottom trawl survey are variable, in part due to year-to-year differences in survey coverage. Surveys since 1996 had the most complete coverage of shallow strata in southeast Alaska, and indicate that stock size is approximately 25-75,000 t (Appendix Figure 1.1). There are no obvious trends in biomass since 1990. We recommend placing southeast Alaska pollock in Tier 5 of NPFMC harvest policy, and basing the ABC and OFL on natural mortality (0.3) and the biomass for the 2009 survey (41,088 t). Biomass in southeast Alaska was estimated by splitting survey strata and CPUE data in the Yakutat INPFC area at 140° W. lon. and combining the strata east of the line with comparable strata in the Southeastern INPFC area. **This results in a 2010 ABC of 9,245 t (41,088 t * 0.75 M), and a 2010 OFL of 12,326 t (41,088 t * M).** These recommendations represent an increase of 12% from the 2008 and 2009 recommendations due to the higher estimated biomass in southeast Alaska in the 2009 NMFS bottom trawl survey.



Appendix Figure 1.1. Pollock length composition in 2009 (left) and biomass trend in southeast Alaska from NMFS bottom trawl surveys in 1990-2009 (right). Error bars indicate plus and minus two standard deviations.

Appendix B: Gulf pollock stock assessment model

Population dynamics

The age-structured model for pollock describes the relationships between population numbers by age and year. The modeled population includes individuals from age 2 to age 10, with age 10 defined as a “plus” group, i.e., all individuals age 10 and older. The model extends from 1961 to 2009 (49 yrs). The Baranov (1918) catch equations are assumed, so that

$$c_{ij} = N_{ij} \frac{F_{ij}}{Z_{ij}} [1 - \exp(-Z_{ij})]$$

$$N_{i+1,j+1} = N_{ij} \exp(-Z_{ij})$$

$$Z_{ij} = \sum_k F_{ik} + M$$

except for the plus group, where

$$N_{i+1,10} = N_{i,9} \exp(-Z_{i,9}) + N_{i,10} \exp(-Z_{i,10})$$

where N_{ij} is the population abundance at the start of year i for age j fish, F_{ij} = fishing mortality rate in year i for age j fish, and c_{ij} = catch in year i for age j fish. A constant natural mortality rate, M , irrespective of year and age, is assumed.

Fishing mortality is modeled as a product of year-specific and age-specific factors (Doubleday 1976)

$$F_{ij} = s_j f_i$$

where s_j is age-specific selectivity, and f_i is the annual fishing mortality rate. To ensure that the selectivities are well determined, we require that $\max(s_j) = 1$. Following previous assessments, a scaled double-logistic function (Dorn and Methot 1990) was used to model age-specific selectivity,

$$s'_j = \left(\frac{1}{1 + \exp[-\beta_1(j - \alpha_1)]} \right) \left(1 - \frac{1}{1 + \exp[-\beta_2(j - \alpha_2)]} \right)$$

$$s_j = s'_j / \max (s'_j)$$

where α_1 = inflection age, β_1 = slope at the inflection age for the ascending logistic part of the equation, and α_2 , β_2 = the inflection age and slope for the descending logistic part.

Measurement error

Model parameters were estimated by maximum likelihood (Fournier and Archibald 1982, Kimura 1989, 1990, 1991). Fishery observations consist of the total annual catch in tons, C_i , and the proportions at age in the catch, p_{ij} . Predicted values from the model are obtained from

$$\hat{C}_i = \sum_j w_{ij} c_{ij}$$

$$\hat{p}_{ij} = c_{ij} / \sum_j c_{ij}$$

where w_{ij} is the weight at age j in year i . Year-specific weights at age are used when available.

Log-normal measurement error in total catch and multinomial sampling error in the proportions at age give a log-likelihood of

$$\log L_k = -\sum_i [\log (C_i) - \log (\hat{C}_i)]^2 / 2 \sigma_i^2 + \sum_i m_i \sum_j p_{ij} \log (\hat{p}_{ij} / p_{ij})$$

where σ_i is standard deviation of the logarithm of total catch ($\sim CV$ of total catch) and m_i is the size of the age sample. In the multinomial part of the likelihood, the expected proportions at age have been divided by the observed proportion at age, so that a perfect fit to the data for a year gives a log likelihood value of zero (Fournier and Archibald 1982). This formulation of the likelihood allows considerable flexibility to give different weights (i.e. emphasis) to each estimate of annual catch and age composition. Expressing these weights explicitly as CVs (for the total catch estimates), and sample sizes (for the proportions at age) assists in making reasonable assumptions about appropriate weights for estimates whose variances are not routinely calculated.

Survey observations consist of a total biomass estimate, B_i , and survey proportions at age π_{ij} . Predicted values from the model are obtained from

$$\hat{B}_i = q \sum_j w_{ij} s_j N_{ij} \exp [\phi_i Z_{ij}]$$

where q = survey catchability, w_{ij} is the survey weight at age j in year i (if available), s_j = selectivity at age for the survey, and ϕ_i = fraction of the year to the mid-point of the survey. Although there are multiple surveys for Gulf pollock, a subscript to index a particular survey has been suppressed in the above and subsequent equations in the interest of clarity. Survey selectivity was modeled using either a double-logistic function of the same form used for fishery selectivity, or simpler variant, such as single logistic function. The expected proportions at age in the survey in the i th year are given by

$$\hat{\pi}_{ij} = s_j N_{ij} \exp[\phi_i Z_{ij}] / \sum_j s_j N_{ij} \exp[\phi_i Z_{ij}]$$

Log-normal errors in total biomass and multinomial sampling error in the proportions at age give a log-likelihood for survey k of

$$\log L_k = -\sum_i [\log(B_i) - \log(\hat{B}_i)]^2 / 2 \sigma_i^2 + \sum_i m_i \sum_j \pi_{ij} \log(\hat{\pi}_{ij} / \pi_{ij})$$

where σ_i is the standard deviation of the logarithm of total biomass (~ CV of the total biomass) and m_i is the size of the age sample from the survey.

Process error

Process error refers to random changes in parameter values from one year to the next. Annual variation in recruitment and fishing mortality can be considered types of process error (Schnute and Richards 1995). In the pollock model, these annual recruitment and fishing mortality parameters are generally estimated as free parameters, with no additional error constraints. We use process error to describe changes in fisheries selectivity over time. To model temporal variation in a parameter γ , the year-specific value of the parameter is given by

$$\gamma_i = \bar{\gamma} + \delta_i$$

where $\bar{\gamma}$ is the mean value (on either a log scale or an arithmetic scale), and δ_i is an annual deviation subject to the constraint $\sum \delta_i = 0$. For a random walk where annual *changes* are normally distributed, the log-likelihood is

$$\log L_{Proc. Err.} = -\sum \frac{(\delta_i - \delta_{i+1})^2}{2 \sigma_i^2}$$

where σ_i is the standard deviation of the annual change in the parameter. We use a process error model for all four parameters of the fishery double-logistic curve. Variation in the intercept selectivity parameters is modeled using a random walk on an arithmetic scale, while variation in the slope parameters is modeled using a log-scale random walk.

The total log likelihood is the sum of the likelihood components for each fishery and survey, plus a term for process error,

$$\text{Log } L = \sum_k \text{Log } L_k + \sum_p \text{Log } L_{Proc.Err.}$$

Appendix C: Seasonal distribution and apportionment of walleye pollock among management areas in the Gulf of Alaska

Since 1992, the Gulf of Alaska pollock TAC has been apportioned between management areas based on the distribution of biomass in groundfish surveys. Both single species and ecosystem considerations provide the rationale for apportioning the TAC. From an ecosystem perspective, apportioning the TAC will spatially distribute the effects of fishing on other pollock consumers (i.e., Steller sea lions), potentially reducing the overall intensity of any adverse effects. Apportioning the TAC also ensures that no smaller component of the stock experiences higher mortality than any other. Although no sub-stock units of pollock have yet been identified in the Gulf of Alaska, it would be precautionary to manage the fishery so that if these sub-units do exist they would not be subject to high fishing mortality. Protection of sub-stock units would be most important during spawning season, when they are spatially separated. The Steller sea lion protection measures implemented in 2001 require apportionment of pollock TAC based on the seasonal distribution of biomass. Although spatial apportionment is intended to reduce the potential impact of fishing on endangered Steller sea lions, it is important to recognize that apportioning the TAC based on an inaccurate or inappropriate estimate of biomass distribution could be detrimental, both to pollock population itself, and on species that depend on pollock.

Walleye pollock in the Gulf of Alaska undergo an annual migration between summer foraging habitats and winter spawning grounds. Since surveying effort has been concentrated during the summer months and prior to spawning in late winter, the dynamics and timing of this migration are not well understood. Regional biomass estimates are highly variable, indicating either large sampling variability, large interannual changes in distribution, or, more likely, both. There is a comprehensive survey of the Gulf of Alaska in summer, but historically surveying during winter has focused on the Shelikof Strait spawning grounds. Recently there has been expanded EIT surveying effort outside of Shelikof Strait in winter, but no acoustic survey has been comprehensive, covering all areas where pollock could potentially occur.

Winter distribution

An annual acoustic survey on pre-spawning aggregations in Shelikof Strait has been conducted since 1981. Since 2000, several additional spawning areas have been surveyed multiple times, including Sanak Gully, the Shumagin Islands, the shelf break near Chirikof Island, and Marmot Bay. Although none of these spawning grounds are as important as Shelikof Strait, especially from a historical perspective, in recent years the aggregate biomass surveyed outside Shelikof Strait has been comparable to that within Shelikof Strait.

As in previous assessments, a “composite” approach was used to estimate the percent of the total stock in each management area. The estimated biomass for each survey was divided by the total biomass of pollock estimated by the assessment model in that year and then split into management areas for surveys that crossed management boundaries. The percent for each survey was added together to form a composite biomass distribution, which was then rescaled so that it summed to 100%. Model estimates of biomass at spawning took into account the total mortality between the start of the year and spawning, and used mean weight at age from Shelikof Strait surveys.

Since time series of biomass estimates for spawning areas outside of Shelikof Strait are now available, we used the four most recent surveys at each spawning area, and used a rule that a minimum of three surveys was necessary to include an area. These criteria are intended to provide estimates that reflect recent biomass distribution while at the same time providing some stability in the estimates. The biomass in these secondary spawning areas tends to be highly variable from one year to the next. Areas meeting these criteria were Shelikof Strait, the shelf break near Chirikof Island, the Shumagin area, and Sanak Gully, but excludes Morzhovoi Bay (surveyed in 2006 and 2007 with questionable timing), Barnabas and

Chiniak Gullies (surveyed once in 2001), and Marmot Bay (surveyed in 2007 and 2009). Finally, an acoustic survey in 1990 along the shelf break and on east side of Kodiak Island (Karp 1990) was used for areas not covered in any of the above surveys.

Vessel comparison experiments conducted between the *R/V Miller Freeman* and the *R/V Oscar Dyson* in Shelikof Strait in 2007 and in the Shumagin/Sanak area in 2008 found significant differences in the ratio of backscatter between the two vessels. The estimated *R/V Oscar Dyson* to *R/V Miller Freeman* ratio for the Shelikof Strait was 1.132, while the ratio for the Shumagin and Sanak areas (taken together) was 1.31. Since the *R/V Oscar Dyson* was designed to minimize vessel avoidance, biomass estimates produced by *R/V Oscar Dyson* should be considered better estimates of the true biomass than those produced by the *R/V Miller Freeman*. These results imply that the biomass in the western GOA (Sanak and Shumagin areas) has historically been underestimated relative to the central GOA. The leading hypothesis for the higher ratio in the western GOA is that the fish are distributed shallower than in Shelikof Strait, and consequently are exposed to a stronger stimulus from the vessel. When calculating the distribution of biomass by area, multipliers were applied to surveys conducted by the *R/V Miller Freeman* to make them comparable to the *R/V Oscar Dyson* (Appendix table 1.1). No vessel comparisons were conducted in the Chirikof area. A vessel specific multiplier of 1.0 was applied as differential avoidance is not expected at fish depths observed in the Chirikof area, where pollock are distributed primarily at depths greater than 300 m (e.g. in 2008 90% of pollock biomass was deeper than 275 m). No evidence for differential backscatter was found for fish deeper than 275 m in vessel comparison experiments conducted during the Shelikof and Bogoslof surveys (no pollock were observed at these depths in the Shumigans), although significant differences were observed at shallower depths.

The sum of the percent biomass for all surveys combined was 77.89%, which may reflect sampling variability, interannual variation in spawning location, or differences in echo sounder/integration systems, but also reflects the recent trend that the aggregate biomass of pollock surveyed acoustically in winter is lower than the assessment model estimates of abundance. After rescaling, the resulting average biomass distribution was 30.22%, 54.04%, 15.74% in areas 610, 620, and 630 (Appendix table 1.1). In comparison to last year's assessment, no area's percentage changed by more than two percentage points.

A-season apportionment between areas 620 and 630

In the 2002 assessment, based on evaluation of fishing patterns which suggested that the migration to spawning areas was not complete by January 20, the plan team recommended an alternative apportionment scheme for areas 620 and 630 based on the midpoint of the summer and winter distributions in area 630. This approach was not used for area 610 because fishing patterns during the A season suggested that most of the fish captured in area 610 would eventually spawn in area 610. The resulting A season apportionment using updated survey data is: 610, 30.22%; 620, 45.81%; 630, 23.97%.

Middleton Island winter EIT survey results in 2003

The apportionment for area 640, which is not managed by season, has previously been based on the summer distribution of the biomass. Fishing, however, takes places primarily in winter or early spring on a spawning aggregation near Middleton Island. During 28-29 March 2003, this area was surveyed by the NOAA ship *Miller Freeman* for the first time and biomass estimate of 6,900 t was obtained. Although maturity stage data suggested the timing of the survey was appropriate, discussions with fishing vessels contacted during the survey raised some questions about survey timing relative to peak biomass. Notwithstanding, a tier 5 calculation based on this spawning biomass gives an ABC of 1,550 t (6,901 t * 0.75 M), compared to 2,075 t for the author's 2010 ABC recommendation and an apportionment based on the summer biomass distribution. This suggests that the current approach of basing the area 640

apportionment on the gulfwide ABC and the summer biomass distribution is at least consistent with the biomass present near Middleton Island in the winter. We recommend continuing this approach until sufficient survey information during winter has accumulated to evaluate interannual variation in the biomass present in this area.

Summer distribution

The NMFS bottom trawl is summer survey (typically extending from mid-May to mid-August). Because of large shifts in the distribution of pollock between management areas one survey to the next, and the high variance of biomass estimates by management area, Dorn et al. (1999) recommended that the apportionment of pollock TAC be based upon the four most recent NMFS summer surveys. The four-survey average was updated with 2009 survey results in an average biomass distribution of 40.14%, 25.84%, 31.32%, and 2.69% in areas 610, 620, 630, and 640 (Appendix Fig. 1.2). Inclusion of the 2009 survey raised the percentage in area 620 by 5 percentage points, and decreased the percentage in 610 and 630 by 2 and 3 percentage points respectively.

Example calculation of 2010 Seasonal and Area TAC Allowances for W/C/WYK

Warning: This example is based on hypothetical ABC of 100,000 t.

1) Deduct the Prince William Sound Guideline Harvest Level.

2) Use summer biomass distribution for the 640 allowance:

$$640 \quad 0.0269 \times \text{Total TAC} = 2,694 \text{ t}$$

3) Calculate seasonal apportionments of TAC for the A, B, C, and D seasons at 25 %, 25%, 25%, and 25% of the remaining annual TAC west of 140° W lon.

$$\text{A season} \quad 0.25 \times (\text{Total TAC} - 2,694) = 24,326 \text{ t}$$

$$\text{B season} \quad 0.25 \times (\text{Total TAC} - 2,694) = 24,326 \text{ t}$$

$$\text{C season} \quad 0.25 \times (\text{Total TAC} - 2,694) = 24,326 \text{ t}$$

$$\text{D season} \quad 0.25 \times (\text{Total TAC} - 2,694) = 24,326 \text{ t}$$

4) For the A season, the allocation of TAC to areas 610, 620 and 630 is based on a blending of winter and summer distributions to reflect that pollock may not have completed their migration to spawning areas by Jan. 20, when the A season opens.

$$610 \quad 0.3022 \times 24,326 \text{ t} = 7,351 \text{ t}$$

$$620 \quad 0.4581 \times 24,326 \text{ t} = 11,145 \text{ t}$$

$$630 \quad 0.2397 \times 24,326 \text{ t} = 5,830 \text{ t}$$

5) For the B season, the allocation of TAC to areas 610, 620 and 630 is based on the composite estimate of winter biomass distribution

$$610 \quad 0.3022 \times 24,326 \text{ t} = 7,351 \text{ t}$$

$$620 \quad 0.5404 \times 24,326 \text{ t} = 13,145 \text{ t}$$

$$630 \quad 0.1574 \times 24,326 \text{ t} = 3,830 \text{ t}$$

6) For the C and D seasons, the allocation of remaining TAC to areas 610, 620 and 630 is based on the average biomass distribution in areas 610, 620 and 630 in the most recent four NMFS bottom trawl surveys of 40.14%, 25.84%, 31.32%, and 2.69%.

$$610 \quad 0.4014 / (1 - 0.0269) \times 24,326 = 10,034 \text{ t}$$

$$620 \quad 0.2584 / (1 - 0.0269) \times 24,326 = 6,461 \text{ t}$$

$$630 \quad 0.3132 / (1 - 0.0269) \times 24,326 = 7,831 \text{ t}$$


$$610 \quad 0.4014 / (1 - 0.0269) \times 24,326 = 10,034 \text{ t}$$

$$620 \quad 0.2584 / (1 - 0.0269) \times 24,326 = 6,461 \text{ t}$$

$$630 \quad 0.3132 / (1 - 0.0269) \times 24,326 = 7,831 \text{ t}$$

Appendix Table 1. Estimates of percent pollock in areas 610-630 during winter EIT surveys in the Gulf of Alaska. The biomass of age-1 pollock The biomass of age-1 fish is not included in Shelikof Strait EIT survey estimates in 2008 (19,090 t), and Shumagin survey estimates in 2006, 2008 and 2009 (12,310 t, 9,339 t and 17,407 t respectively).

<i>Survey</i>	<i>Year</i>	<i>Model estimates of total 2+ biomass at spawning</i>	<i>Survey biomass estimate</i>	<i>Multiplier from vessel comparison (OD/MF)</i>	<i>Percent</i>	<i>Percent by management area</i>		
						<i>Area 610</i>	<i>Area 620</i>	<i>Area 630</i>
Shelikof	2006	518,623	293,609	1.13	56.6%	0.0%	96.1%	3.9%
Shelikof	2007	505,673	180,881	1.13	35.8%	0.0%	97.1%	2.9%
Shelikof	2008	598,378	188,942	1.00	31.6%	0.0%	93.4%	6.6%
Shelikof	2009	673,151	265,971	1.00	39.5%	0.0%	95.6%	4.4%
Shelikof	Average				40.9%	0.0%	95.6%	4.4%
	Percent of total 2+ biomass					0.0%	39.0%	1.8%
Chirikof	2006	518,623	69,000	1.00	13.3%	0.0%	28.3%	71.7%
Chirikof	2007	505,673	35,573	1.00	7.0%	0.0%	24.0%	76.0%
Chirikof	2008	598,378	22,055	1.00	3.7%	0.0%	50.2%	49.8%
Chirikof	2009	673,151	396	1.00	0.1%	0.0%	0.0%	100.0%
Chirikof	Average				6.0%	0.0%	25.6%	74.4%
	Percent of total 2+ biomass					0.0%	1.5%	4.5%
Shumagin	2006	518,623	25,028	1.31	6.3%	92.8%	7.2%	0.0%
Shumagin	2007	505,673	20,009	1.31	5.2%	98.5%	1.5%	0.0%
Shumagin	2008	598,378	21,244	1.31	4.7%	77.2%	22.8%	0.0%
Shumagin	2009	673,151	45,357	1.00	6.7%	61.4%	38.6%	0.0%
Shumagin	Average				5.7%	82.5%	17.5%	0.0%
	Percent of total 2+ biomass					4.7%	1.0%	0.0%
Sanak	2006	518,623	127,214	1.31	32.1%	100.0%	0.0%	0.0%
Sanak	2007	505,673	60,289	1.31	15.6%	100.0%	0.0%	0.0%
Sanak	2008	598,378	19,750	1.31	4.3%	100.0%	0.0%	0.0%
Sanak	2009	673,151	31,435	1.00	4.7%	100.0%	0.0%	0.0%
Sanak	Average				17.4%	100.0%	0.0%	0.0%
	Percent of total 2+ biomass					17.4%	0.0%	0.0%
Karp (1990)	1990	986,372	78,134	1.00	7.9%	18.4%	6.3%	75.3%
	Average				7.9%	18.4%	6.3%	75.3%
	Percent of total 2+ biomass					1.5%	0.5%	6.0%
Total					77.89%	23.54%	42.09%	12.26%
Rescaled total					100.00%	30.22%	54.04%	15.74%



Appendix Figure 1.2. Percent distribution of Gulf of Alaska pollock biomass west of 140° W lon. in NMFS bottom trawl surveys in 1984-2009. The percent in West Yakutat in 1984, 1987, and 2001 was set equal to the mean percent in 1990-99.