1.B. Assessment of walleye pollock in the Bogoslof Island Region

James N. Ianelli, S. J. Barbeaux, and D. McKelvey Alaska Fisheries Science Center National Marine Fisheries Service

[NOTE: In accordance with the approved schedule, no assessment was conducted for this stock this year, however, a full stock assessment will be conducted in 2022. Until then, the values generated from the previous stock assessment (below) will be rolled over for 2022 specifications]

Summary of Results

The ABC and OFL levels using Tier 5 values and assuming the random-effects model:

The ADC and OT L levels using The .		0	eneeds model.	
	As estimated or		As estimated or	
	specified last year for:		recommended this year for:	
Quantity	2020	2021	2021	2022
M (natural mortality rate)	0.3	0.3	0.3	0.3
Tier	5	5	5	5
Biomass (t)	610,267	610,267	378,262	378,262
F _{OFL}	0.300	0.300	0.300	0.300
maxF _{ABC}	0.225	0.225	0.225	0.225
F _{ABC}	0.225	0.225	0.225	0.225
OFL (t)	183,080	183,080	113,479	113,479
maxABC (t)	137,310	137,310	85,109	85,109
ABC (t)	137,310	137,310	85,109	85,109
	As determined <i>this</i> year for:		As determined <i>this</i> year for:	
Status	2018	2019	2019	2020
Overfishing	No	n/a	No	n/a